

Week 4: (continued)

2.2 Subspaces/Spanning (continued)

2.3 Independence/Bases

2.4 Nullspaces

Linear Independence/Bases and Dimension

Vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$ in a

vector space V are **linearly dependent** if there are

scalars c_1, c_2, \dots, c_k so $\vec{0} = c_1\vec{v}_1 + c_2\vec{v}_2 + \dots + c_k\vec{v}_k$,

with at least one $c_j \neq 0$. Such a vector equation

is said to be a **relation** of linear dependence among the

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$. A collection of vectors for which the only solution of

the vector equation is the **trivial solution**, $c_1 = 0, \dots, c_k = 0$

is said to be **linearly independent**.

Examples: (1) \vec{i}, \vec{j} in R^2 ; and (2) $\{\vec{i}, \vec{j}, \vec{k}\}$ in R^3 ; are linearly independent.

Verification: $c_1\vec{i} + c_2\vec{j} + c_3\vec{k} = c_1(1, 0, 0) + c_2(0, 1, 0) + c_3(0, 0, 1) = (c_1, c_2, c_3)$
 $= (0, 0, 0) = \vec{0}$, only when $c_1 = 0, c_2 = 0, c_3 = 0$.

Our main objective is the definition of basis.

Vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$ in a vector space V that are both

(1) a spanning set for V and (2) linearly independent

are called a **basis** for V . So

(1) \vec{i}, \vec{j} are a basis of R^2 ; and (2) $\{\vec{i}, \vec{j}, \vec{k}\}$ is a basis of R^3 .

A vector space has many bases, but the **main fact**

is that if $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$ is a basis of V and $\vec{w}_1, \vec{w}_2, \dots, \vec{w}_j$

is a second basis of V , then the number of

vectors is the same: $k = j$. The number

of vectors in a basis of V is called the **dimension** of V .

We start with verifications of linear

independence in subspaces of \mathbf{R}^n

Problem

Determine whether the vectors $(1, 2, 3)$,

$(1, -1, 2)$ and $(1, -4, 1)$ are LI or LD in \mathbf{R}^3 .

If they are LD find a relation of linear dependence.

Solution [again; same method!]

Write the equation as an equation in column vectors,

$$\begin{aligned} \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} &= c_1 \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + c_3 \begin{pmatrix} 1 \\ -4 \\ 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 & 1 \\ 2 & -1 & -4 \\ 3 & 2 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix}. \end{aligned}$$

Observe that this is the homog linear system with coef matrix $A = (\vec{v}_1 \vec{v}_2 \vec{v}_3)$.

For n -by- n systems the system has a

unique solution exactly when the determinant is non-zero.

$$\text{We have } \begin{vmatrix} 1 & 1 & 1 \\ 2 & -1 & -4 \\ 3 & 2 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 1 & 1 \\ 0 & -3 & -6 \\ 0 & -1 & -2 \end{vmatrix} = 0$$

so the system has nontrivial solutions and the vectors are LD.

To find a relation we continue the reduction $A \rightarrow \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$,

so a spanning vector is $(1, -2, 1)$,

and $\vec{0} = \vec{v}_1 - 2\vec{v}_2 + \vec{v}_3$ is a nontrivial relation of LD.

For additional practice, we try 2.2 - 2c, 4c, 9 and

the text's Example 8 of a basis in \mathbf{R}^3 .

Also, see the text for the definition of **components**

and the **component vector**, which appears on the HW (this week).

Example: Show that the vectors $\vec{v}_1 = (1, 1)$

$\vec{v}_2 = (1, -1)$ are a basis of \mathbf{R}^2 .

Solution. We know that \vec{i}, \vec{j} are a basis

of \mathbf{R}^2 ; so that the dimension of \mathbf{R}^2 is

We check that $\vec{v}_1 = (1, 1)$ and $\vec{v}_2 = (1, -1)$

are linearly independent (LI) since $\begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix}$

$= -2 \neq 0$. Checking that \vec{v}_1 and \vec{v}_2 span \mathbf{R}^2

by expressing each $\vec{v} = (x, y) \in \mathbf{R}^2$ as a

linear combination (as we did for \vec{i}, \vec{j})

takes a little bit of effort, and involves fractions.

Checking that each vector equation

$\vec{v} = c_1\vec{v}_1 + c_2\vec{v}_2$ has a

solution (without finding c_1, c_2) is easier. (why?)

But the main fact has a **practical version** that saves any computation, or even very much further thought. If \vec{v}_1 and \vec{v}_2 did not span \mathbf{R}^2 there would be a 3rd vector \vec{v}_3 not in the span of \vec{v}_1, \vec{v}_2 .

But that would make $\vec{v}_1, \vec{v}_2, \vec{v}_3$ linearly independent. We could consider adding further vectors, each new one giving a larger collection of indep vectors; but we already have too many, as we've shown $\dim(\mathbf{R}^2) \geq 3$. So \vec{v}_1, \vec{v}_2 have to span (for free, given the main fact), so \vec{v}_1, \vec{v}_2 is a basis for \mathbf{R}^2 .

This is the use of the text's

Theorem 2.12.(1) If the $\dim(V) = n$ for a vector space V , then any LI set of n vectors is a basis of V .