

NAME : B. Dodson

1. In this problem, solutions to differential equations are not to include integrals.

(a) (10 points) Solve the differential equation $x \frac{dy}{dx} + y^2 = 0$, and write your solution as an explicit function $y = y(x)$.

(b) (10 points) Find the general solution of $\frac{dy}{dx} - \frac{y}{x} = 3x^3$.

Solutions:

(a) Separated equations is $-y^{-2} dy = x^{-1} dx$. Integrating,
 $\frac{1}{y} = \ln x + c$; so the explicit solution is $y(x) = \frac{1}{\ln x + c}$.

(b) The integrating factor is $f = e^{\int -\frac{1}{x} dx} = e^{-\ln x} = e^{\ln(x^{-1})} = \frac{1}{x}$.

Multiplying by the integrating factor, and re-writing the left side, $(\frac{y}{x})' = 3x^2$. Integrating, $\frac{y}{x} = x^3 + c$, so $y = x^4 + cx$.

2. (15 points) A tank whose volume is 50 L initially contains 25 g of salt dissolved in 25 L of water. A solution containing 5 g/L of salt is pumped into the tank at a rate of 3 L/min, and the well-stirred mixture flows out at a rate of 1 L/min. Find the amount of salt in the tank after 12 minutes. Include in your solution a clear statement of the differential equation satisfied by the amount of salt $A(t)$ in the tank at time t .

Solution: The volume at time t is $V(t) = 2t + 25$.

The differential equation is obtained from the rule “rate-in - rate-out” as

$$A' = r_1 c_1 - r_2 c_2 = 15 - \frac{1}{V} A,$$

with $r_1 = 3, c_1 = 5, r_2 = 1$ and $c_2 = \frac{A}{V}$. Using our solution for $V(t)$,

the differential equation is $A' + \frac{1}{2t+25} A = 15$.

The integrating factor here is $(2t + 25)^{\frac{1}{2}}$. Then

$(A(2t + 25)^{\frac{1}{2}})' = 15(2t + 25)^{\frac{1}{2}}$ gives $A(t) = 5(2t + 25) + \frac{c}{(2t + 25)^{\frac{1}{2}}}$. The initial condition $A(0) = 25$ gives $c = -500$, so $A(t) = 5(2t + 25) - \frac{500}{(2t+25)^{\frac{1}{2}}}$. Finally,

$$A(12) = (5)(49) - \frac{500}{\sqrt{49}} = 245 - \frac{500}{7}.$$

3. (25 points) Find the general solution of the following differential equations. Partial credit in parts (b) and (c) will be given for a clearly stated solution to the homogeneous equation. In this problem solutions to differential equations must be expressed only in terms of real numbers and real functions.

(a) $y''(t) + y'(t) - 6y(t) = 0$.

(b) $y''(t) + y'(t) - 2y(t) = -12e^{-2t}$.

(c) $y''(x) + 4y'(x) + 4y(x) = 3\cos(2x)$.

Solutions:

(a) $r^2 + r - 6 = (r + 3)(r - 2)$, so $r = 2, -3$ (or use the quadratic formula).

So $y = c_1e^{2x} + c_2e^{-3x}$.

(b) Checking the right side, we consider the trial function $y_p = Ae^{-2t}$. We first check y_H , and find $y_H = c_1e^x + c_2e^{-2t}$.

Since e^{-2t} occurs in y_H , the function Ae^{-2t} cannot possibly satisfy the non-homogeneous equation. So we take the “modified trial function”

$$y_p = Ate^{-2t}.$$

Differentiating twice, we find $-3A = -12$, so $y_p = 4te^{-2t}$, and

$$y = y_H + y_p = c_1e^x + c_2e^{-2t} + 4te^{-2t}.$$

(c) The solution of the homogeneous equation is $y_H = (c_1 + c_2x)e^{-2x}$.

For the trial function, we take the usual trial function, $y_p = A\cos(2x) + B\sin(2x)$.

Differentiating twice, and plugging y'_p and y''_p into the

differential equation, we equate coefficients of $\cos(2x)$ and $\sin(2x)$.

We start with $\sin(2x)$, which is the simpler coefficient, $-4B - 8A + 4B = 0$, which gives $A = 0$. Then the coefficients of $\cos(2x)$ are $8B = 3$,

using $A = 0$. We double-check, and see $y_p = \frac{3}{8}\sin(2x)$ is a solution,

$$\text{and } y = y_H + y_p = (c_1 + c_2x)e^{-2x} + \frac{3}{8}\sin(2x).$$

4. (10 points) Recall that the motion $y(x)$ of a spring-mass system is governed by

$$my''(x) + cy'(x) + ky(x) = 0,$$

where m is the mass, c is the friction constant and k is the spring constant. If $m = 2, c = 4$ and $k = 10$ find the solution to the Initial Value problem $y(0) = 3, y'(0) = -1$. In this problem solutions to differential equations must be expressed only in terms of real numbers and real functions.

Solution: Substituting the given values, and dividing by 2,

$y'' + 2y' + 5y = 0$. So the roots satisfy $r^2 + 2r + 5 = 0$, and $r = -1 \pm 2i$.

So $y = e^{-x} (c_1 \cos(2x) + c_2 \sin(2x))$.

Update: In terms of this semester's syllabus, the motion is described as "damped with oscillations". Note also problem 3b from the Spring 2006 exam, where (i) is "damped with oscillations"; (ii) is overdamped; and, (iii) is critically damped. The terminology appears in the text material of 4.5, and is summarized in problem 15 (which was assigned).