

Variance asymptotics and central limit theorems for generalized growth processes with applications to convex hulls and maximal points

T. Schreiber¹ and J. E. Yukich²

September 10, 2007

Abstract

We show that the random point measures induced by vertices in the convex hull of a Poisson sample on the unit ball, when properly scaled and centered, converge to those of a mean zero Gaussian field. We establish limiting variance and covariance asymptotics in terms of the density of the Poisson sample. Similar results hold for the point measures induced by the maximal points in a Poisson sample. The approach involves introducing a generalized spatial birth growth process allowing for cell overlap.

1 Introduction, Main Results

Given X_i , $i \geq 1$, i.i.d. random variables with values in a d -dimensional convex set S , $d \geq 2$, a classic problem in convex geometry involves determining the distribution of the number of points in the set of extreme points $\mathcal{V}(\{X_i\}_{i=1}^n)$, defined as the vertices in the convex hull of $\{X_i\}_{i=1}^n$. This problem was first considered by Rényi and Sulanke [37], with recent notable progress by Reitzner [32] - [35] and Van Vu [44].

A closely related problem involves determining, for a given $K \subset \mathbb{R}^d$, the distribution of the number of points in the set $\mathcal{M}_K(\{X_i\}_{i=1}^n)$ of K -maximal points, where a point X_j belongs to

American Mathematical Society 2000 subject classifications. Primary 60F05, Secondary 60D05

Key words and phrases. Convex hulls, maximal points, spatial birth growth processes, Gaussian limits

¹ Research supported in part by Polish Minister of Scientific Research and Information Technology grant 1 P03A 018 28 (2005-2007)

² This material is based upon work supported by the NSF under grant DMS-0203720

$\mathcal{M}_K(\{X_i\}_{i=1}^n)$ iff $(X_j \oplus K) \cap \{X_i\}_{i=1}^n = X_j$, where here and henceforth for all $B \subset \mathbb{R}^d$ and $x \in \mathbb{R}^d$ we write $x \oplus B := \{x + y : y \in B\}$. When K is $(\mathbb{R}^+)^d$ then $\mathcal{M}_K(\{X_i\}_{i=1}^n)$ is simply the set of maximal points, that is those points X_j in $\{X_i\}_{i=1}^n$ having the property that no point X_i , $i \neq j$, exceeds it in all coordinates. The limit theory for the number of maximal points in $\mathcal{M}_K(\{X_i\}_{i=1}^n)$ was first considered by Rényi [36] and Barndorff-Nielsen and Sobel [5]. Chen et al. [11] surveys the vast literature, which includes books by Ehrgott [18], Pomerol and Barba-Romero [30], and recent papers of Bai et al. [1, 2], Devroye [17], Barbour and Xia [4], and [8].

In this paper we establish convergence of the finite dimensional distributions of the re-scaled point measures induced by the random point sets $\mathcal{V}(\mathcal{P}_{\lambda\rho})$ where $\mathcal{P}_{\lambda\rho}$ denotes a Poisson point process of intensity $\lambda\rho$ on B_d , the unit radius d -dimensional ball centered at the origin and where ρ is a continuous density on B_d . For sets $K := \{(w_1, \dots, w_d) : w_d \geq (w_1^2 + \dots + w_{d-1}^2)^{\alpha/2}\}$ where $\alpha \in (0, 1]$ is fixed, we also establish convergence of the finite dimensional distributions of the point measures induced by $\mathcal{M}_K(\mathcal{P}_{\lambda\rho})$, where $\mathcal{P}_{\lambda\rho}$ denotes the Poisson point process of intensity $\lambda\rho$ on $A \times \mathbb{R}_+$, where $A \subset \mathbb{R}^{d-1}$ is compact and convex and where $\rho : A \times \mathbb{R}_+$ is continuous. These results are facilitated by introducing a generalized spatial birth growth process as a means towards obtaining explicit variance asymptotics and central limit theorems for random measures arising in convex geometry. The relevant spatial birth growth process, possibly of independent interest, modifies the classical spatial birth growth process introduced by Kolmogorov [23] as a model for crystal growth by allowing the possibility of *cell overlap*. As in [23], cells may grow at non-constant growth rates.

In the context of the set of extreme points $\mathcal{V}(\mathcal{P}_{\lambda\rho})$, the approach taken here adds to the work of Reitzner [32]-[35] and Van Vu [44] in the following ways. First, the present set-up establishes convergence of the finite dimensional distributions of the canonical point measures induced by $\mathcal{V}(\mathcal{P}_{\lambda\rho})$, whereas [32]-[35] and [44] deal with one dimensional central limit theorems. Second, we establish a formula for variance and covariance asymptotics. Third, the present paper concerns limit theory for non-uniform samples, whereas [32]-[35] and [44] treat uniform random samples.

In the context of the set of maximal points $\mathcal{M}_K(\mathcal{P}_{\lambda\rho})$, the present set-up establishes convergence of the finite dimensional distributions of the canonical point measures induced by $\mathcal{M}_K(\mathcal{P}_{\lambda\rho})$, with covariances, whereas previous work [4, 17] is concerned with one dimensional central limit theorems without a formula for covariance asymptotics and/or is limited to the case when K is a cone [8].

1.1 Terminology, ψ -growth processes

Let the function $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfy the following conditions:

($\Psi 1$) ψ is monotone and $\lim_{l \rightarrow \infty} \psi(l) = \infty$, and

($\Psi 2$) there exists $\alpha > 0$ such that $\psi(l) = l^\alpha(1 + o(1))$ for l small enough.

Let $\mathbf{0}$ denote the origin of \mathbb{R}^{d-1} , $d \geq 2$, and let $|y|$ denote the Euclidean norm of $y \in \mathbb{R}^d$. We define $K[\mathbf{0}]$ to be the ψ -hypergraph $\{(y, h) \in \mathbb{R}^{d-1} \times \mathbb{R}_+ : h \geq \psi(|y|)\}$ and, more generally, for $\bar{x} := (x, h_x) \in \mathbb{R}^{d-1} \times \mathbb{R}_+$ we define its ψ -hypergraph (or upwards cone) by

$$K[\bar{x}] := \bar{x} \oplus K[\mathbf{0}] := \{(y, h) \in \mathbb{R}^{d-1} \times \mathbb{R}_+ : h \geq h_x + \psi(|y - x|)\}. \quad (1.1)$$

Given a point set $\mathcal{X} \subseteq \mathbb{R}^{d-1} \times \mathbb{R}_+$, a point $\bar{x} \in \mathcal{X}$ is called ψ -extremal in \mathcal{X} iff $K[\bar{x}] \not\subseteq \bigcup_{\bar{y} \in \mathcal{X} \setminus \{\bar{x}\}} K[\bar{y}]$, that is to say the ψ -hypergraph of \bar{x} is not completely covered by the union of the ψ -hypergraphs of points in $\mathcal{X} \setminus \{\bar{x}\}$. Define the functional

$$\xi(\bar{x}, \mathcal{X}) := \xi(\psi; \bar{x}, \mathcal{X}) := \begin{cases} 1 & \text{if } \bar{x} \text{ is } \psi\text{-extremal in } \mathcal{X} \\ 0 & \text{otherwise.} \end{cases} \quad (1.2)$$

With D standing for some bounded domain in $\mathbb{R}^{d-1} \times \mathbb{R}_+$, we consider the version $\xi_D(\cdot, \cdot)$ of $\xi(\cdot, \cdot)$ restricted to D , by setting $\xi_D(\bar{x}, \mathcal{X})$ to be 1 iff $K[\bar{x}] \cap D \not\subseteq \bigcup_{\bar{y} \in (\mathcal{X} \setminus \{\bar{x}\}) \cap D} K[\bar{y}]$, in which case we declare \bar{x} to be ψ -extremal in $D \cap \mathcal{X}$, and otherwise we set $\xi_D(\bar{x}, \mathcal{X})$ to be zero. In case $\bar{x} \notin \mathcal{X}$ we abbreviate notation and write $\xi(\bar{x}, \mathcal{X})$ for $\xi(\bar{x}, \mathcal{X} \cup \bar{x})$ and similarly for $\xi_D(\bar{x}, \mathcal{X})$.

To provide a physical interpretation of these functionals we regard $\mathbb{R}^{d-1} \times \mathbb{R}_+$ as d -dimensional space time, with \mathbb{R}_+ standing for the time coordinate, and we interpret the graph $\partial(K[\bar{x}])$, $\bar{x} := (x, t)$ as the boundary of a $(d-1)$ -dimensional spherical particle born at x at time t (at which time it has initial radius zero) and growing thereupon with radial speed $v(t) := \frac{d}{dt}[\psi^{-1}(t)]$, provided the derivative exists. The particles (spheres) grow independently and do not exhibit exclusion, that is they may overlap or penetrate one another. A particle is *extreme* iff at some time it is not completely covered by other particles. When ψ is the identity, so that the ψ graph gives a cone, we see that ψ -extremal points coincide with maximal points [8].

In the context of this representation it should be noted that, unlike the one stated here, the classic growth process (see e.g. [23, 7, 13, 27]) assumes that particles, upon being born at random locations $x \in \mathbb{R}^{d-1}$ at random times $h_x \in \mathbb{R}^+$, form a cell by growing radially in all directions with a possibly non-constant speed, i.e., with ψ possibly non-linear. When one growing cell touches

another, it stops growing in that direction, i.e. no overlap is allowed. Furthermore, a particle born inside an existing cell is *discarded*, otherwise it is *accepted*. Letting $\hat{\xi}(\bar{x}, \mathcal{X})$ be zero or one according to whether \bar{x} is accepted or not, this paper also considers such functionals $\hat{\xi}$.

The growth process giving rise to the functional ξ will henceforth be called the *ψ -growth process with overlap*, while the process corresponding to $\hat{\xi}$ will be referred to as the *ψ -growth process without overlap*. This paper will mainly concentrate on applications of the first concept and the corresponding functional ξ , but the subsequently developed general theory also treats the latter concept in the special case of linear ψ . Throughout, let A be a compact convex subset of \mathbb{R}^{d-1} . We shall also admit the case $A := \mathbb{R}^{d-1}$ in the sequel, in which case we assume that ρ is uniformly bounded. Consider a density function ρ on $A_+ := A \times \mathbb{R}_+$, not necessarily integrable, such that

(R1) ρ is continuous on A_+ ,

(R2) there exists a constant $\delta \geq 0$ and a continuous function $\rho_0 : A \rightarrow \mathbb{R}_+$ bounded away from zero such that

$$\rho(x, h) = \rho_0(x)h^\delta(1 + o(1))$$

for h small enough and $\rho(x, h) = O(h^\delta)$ for large h uniformly in $x \in A$.

For $\lambda > 0$ we recall that $\mathcal{P}_{\lambda\rho}$ denotes the Poisson point process on A_+ with intensity measure $\lambda\rho(x, h)dx dh$. The ‘extreme point’ empirical measures $\mu_{\lambda\rho}^\xi$ and $\mu_{\lambda\rho}^{\hat{\xi}}$ generated by $\mathcal{P}_{\lambda\rho}$ are

$$\mu_{\lambda\rho}^\xi := \sum_{\bar{x} \in \mathcal{P}_{\lambda\rho}} \xi(\bar{x}, \mathcal{P}_{\lambda\rho}) \delta_{\bar{x}} \quad (1.3)$$

and

$$\mu_{\lambda\rho}^{\hat{\xi}} := \sum_{\bar{x} \in \mathcal{P}_{\lambda\rho}} \hat{\xi}(\bar{x}, \mathcal{P}_{\lambda\rho}) \delta_{\bar{x}}, \quad (1.4)$$

with δ_x standing for the unit point mass at $x \in \mathbb{R}^d$. For any random measure σ on \mathbb{R}^d we write $\bar{\sigma}$ for its centered version $\sigma - \mathbb{E}[\sigma]$, so that e.g. $\bar{\mu}_{\lambda\rho}^\xi := \mu_{\lambda\rho}^\xi - \mathbb{E}[\mu_{\lambda\rho}^\xi]$.

Notice that for small α the upwards cones $K[\bar{x}]$ have relatively narrow apertures, making it less likely that cones having apexes with a small temporal coordinate get covered by ψ -hypergraphs, i.e., one expects more ψ -extreme points as α gets smaller. Also, roughly speaking, for small δ one expects more points in $\mathcal{P}_{\lambda\rho}$ with small temporal coordinate and thus more ψ -extreme points in this case as well. One of the goals of this paper is to show (see Theorem 1.1) that the expected total mass of the extreme point empirical measures (1.3)-(1.4) is asymptotically proportional to λ^τ , where

$$\tau := \tau(d, \alpha, \delta) := \frac{d-1}{d-1+\alpha(1+\delta)}. \quad (1.5)$$

More general goals include establishing the variance asymptotics and the convergence of the finite dimensional distributions of the appropriately scaled measures (1.3)- (1.4) to Gaussian distributions (see Theorems 1.2 and 1.3) and to treat the applications to extreme and maximal points described at the outset.

Notation. Given $\alpha > 0$, put

$$\psi^{(\infty)}(l) := l^\alpha. \quad (1.6)$$

Recalling the definition of ξ , we define the functional $\xi^{(\infty)}$ by $\xi^{(\infty)}(\cdot, \cdot) := \xi(\psi^{(\infty)}; \cdot, \cdot)$ and similarly for $\hat{\xi}^{(\infty)}$. We also let \mathcal{P}_* stand for the Poisson point process in $\mathbb{R}^{d-1} \times \mathbb{R}_+$ with intensity measure $h^\delta dx dh$.

For all $\bar{x} := (x, h_x)$ and $\bar{y} := (y, h_y)$, let

$$m^{(\infty)}(\bar{x}) := \mathbb{E} [\xi^{(\infty)}(\bar{x}, \mathcal{P}_*)]$$

and

$$c_*^{(\infty)}(\bar{x}, \bar{y}) := \mathbb{E} [\xi^{(\infty)}(\bar{x}, \mathcal{P}_* \cup \bar{y}) \xi^{(\infty)}(\bar{y}, \mathcal{P}_* \cup \bar{x})] - \mathbb{E} [\xi^{(\infty)}(\bar{x}, \mathcal{P}_*)] \mathbb{E} [\xi^{(\infty)}(\bar{y}, \mathcal{P}_*)]$$

respectively denote the one and two point correlation functions for the $\psi^{(\infty)}$ growth process with overlap.

For sets A and $B \subset \mathbb{R}^d$, let $d(A, B) := \inf\{|x - y| : x \in A, y \in B\}$. Let $B_d(y, r)$ denote the d -dimensional Euclidean ball centered at $y \in \mathbb{R}^d$ with radius $r \in (0, \infty)$.

Given a subset B of \mathbb{R}^d , let $\mathcal{C}_b(B)$ denote the bounded continuous functions on B . For any signed measure μ on A_+ and $f \in \mathcal{C}_b(A_+)$, let $\langle f, \mu \rangle := \int f d\mu$. Unless otherwise specified C denotes a generic positive constant whose value may change from line to line.

1.2 Limit theory for Ψ -growth functionals

For all $f \in \mathcal{C}_b(A_+)$ with $A \subset \mathbb{R}^{d-1}$ compact and convex we define the average of the product of f and the one and two point correlation functions as follows:

$$I(f) := \int_A \int_0^\infty f(x, 0) m^{(\infty)}(\mathbf{0}, h') \rho_0^\tau(x) (h')^\delta dh' dx \quad (1.7)$$

and

$$J(f) := \int_A \int_0^\infty \int_{\mathbb{R}^{d-1}} \int_0^\infty f(x, 0) c_*^{(\infty)}((\mathbf{0}, h'), (y', h'_y)) \rho_0^\tau(x) (h'_y)^\delta (h')^\delta dh'_y dy' dh' dx. \quad (1.8)$$

The finiteness of $I(f)$ follows by Lemmas 3.2 and 3.3 (see the bound (3.11)) whereas the finiteness of $J(f)$ follows from Lemmas 3.4 and 3.5 (see the bound (3.22)) which imply rapid enough decay of two-point correlation functions.

The following are our main results. We state the results for $\mu_{\lambda\rho}^\xi$ and note that analogous results hold for $\mu_{\lambda\rho}^{\hat{\xi}}$ when ψ is linear. The first result specifies first order behavior whereas the second provides second order asymptotics.

Theorem 1.1 *We have for all $f \in \mathcal{C}_b(A_+)$*

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\tau} \mathbb{E}[\langle f, \mu_{\lambda\rho}^\xi \rangle] = I(f). \quad (1.9)$$

Theorem 1.2 *We have for all $f \in \mathcal{C}_b(A_+)$*

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\tau} \text{Var}[\langle f, \mu_{\lambda\rho}^\xi \rangle] = I(f^2) + J(f^2). \quad (1.10)$$

The next result establishes the convergence of the finite dimensional distributions of $(\lambda^{-\tau/2} \bar{\mu}_{\lambda\rho}^\xi)$.

Theorem 1.3 *The finite dimensional distributions $\lambda^{-\tau/2}(\langle f_1, \bar{\mu}_\lambda^\xi \rangle, \dots, \langle f_k, \bar{\mu}_\lambda^\xi \rangle)$, $f_1, \dots, f_k \in \mathcal{C}_b(A_+)$, of $(\lambda^{-\tau/2} \bar{\mu}_{\lambda\rho}^\xi)$ converge as $\lambda \rightarrow \infty$ to those of a mean zero Gaussian field with covariance kernel*

$$(f, g) \mapsto I(fg) + J(fg), \quad f, g \in \mathcal{C}_b(A_+). \quad (1.11)$$

Section two describes applications of ψ -growth processes with overlap, as given by the general limits of Theorems 1.1 - 1.3, to convex hulls and maximal points of i.i.d. samples.

Remarks. (i) *Applications to the ψ -growth process $\hat{\xi}$ without overlap.* The results of Theorems 1.1 - 1.3 for the functional $\hat{\xi}$ provide variance asymptotics and central limit theorems for the classic spatial birth-growth model in \mathbb{R}^{d-1} whereby seeds are born at random locations in \mathbb{R}^{d-1} and times in \mathbb{R}_+ according to the Poisson point process $\lambda\mathcal{P}_{\lambda\rho}$ on $\lambda^{1/d}A \times \mathbb{R}_+$ and grow linearly in time. Theorems 1.1 - 1.3 for $\hat{\xi}$ provide a central limit theorem for the number of seeds accepted in such models. This generalizes and extends [7, 27], which builds on work of Chiu and Quine [13], [14], Chiu [12], and Chiu and Lee [15], which do not consider convergence of finite dimensional distributions and which often restrict to models with homogeneous temporal input.

(ii) *Scaling.* The scaling $\lambda^{-\tau}$ arises in the following way. From a conceptual and analytic point of view, it is convenient to re-scale the ψ -growth process in time and space so as to obtain an equivalent growth process on Poisson points of approximately unit intensity density on a region of volume λ . The scaling is designed to asymptotically preserve the ψ -hypergraphs and the behavior of the density locally close to $h = 0$.

To achieve this we scale A_+ in the $d - 1$ spatial directions by λ^β and in the temporal direction by λ^γ . Under this temporal scaling and under **(R2)** the density ρ exhibits growth $(h\lambda^\gamma)^\delta$ for small temporal h , and we thus require $\lambda^{\beta(d-1)+\gamma(1+\delta)} = \lambda$. This scaling maps $|x|$ and h_x to $\lambda^\beta|x|$ and $\lambda^\gamma h_x$, respectively, and therefore it asymptotically preserves the ψ -hypergraphs and condition **(Ψ2)** provided $(\lambda^\beta|x|)^\alpha = \lambda^\gamma h_x(1 + o(1))$ for (x, h_x) lying on the graph of ψ , i.e. $h_x = \psi(x)$. Since $h_x = |x|^\alpha(1 + o(1))$ for such (x, h_x) , we require $\lambda^{\beta\alpha} = \lambda^\gamma$. We thus require the relations

$$\beta(d - 1) + \gamma(1 + \delta) = 1 \quad \text{and} \quad \beta\alpha = \gamma$$

which yields these values for the scaling exponents

$$\beta = \frac{\gamma}{\alpha} \quad \text{and} \quad \gamma = \frac{\alpha}{(d - 1) + \alpha(1 + \delta)}. \quad (1.12)$$

Given the re-scaled ψ -growth process on $\lambda^\beta A \times \mathbb{R}_+$ we expect that a point is ψ -extremal (i.e., $\xi = 1$) iff its time coordinate is small. Thus the functional $\mu_{\lambda\rho}^\xi(A_+)$ should exhibit growth proportional to the Lebesgue measure of $\lambda^\beta A$, that is proportional to $\lambda^{\beta(d-1)} = \lambda^\tau$. In the special case when $\delta = 0$ and the growth is linear ($\alpha = 1$) the ψ -hypergraphs are preserved by time and space scaling by $\lambda^{1/d}$, that is $\gamma = 1/d = \beta$. Thus $\tau = (d - 1)/d$ in this case.

(iii) *de-Poissonization.* In section four we de-Poissonize Theorems 1.1 - 1.3 when $\alpha \in (0, 1]$. In other words we obtain identical limit theory when $\mathcal{P}_{\lambda\rho}$ is replaced by i.i.d. random variable X_1, \dots, X_n , chosen in A_+ according to the density ρ , assumed to be integrable to 1. We expect similar de-Poissonization results for $\alpha > 1$ but are unable to prove this.

(iv) We have not tried to establish a.s. convergence in (1.9), but expect that concentration inequalities should be useful in this context.

1.3 Notation and scaling relations

Motivated by remark (ii) above, we place the ψ -growth process on its proper scale by re-scaling as follows. With β and γ as in (1.12), for a *fixed* $x \in A$ and any generic point $\bar{y} := (y, h_y) \in A_+$ we put $\bar{y}^{(\lambda)} := \bar{y}' := (y', h'_y)$ with

$$y' := y^{(\lambda)} := \lambda^\beta(y - x) \quad \text{and} \quad h'_y := h_y^{(\lambda)} := \lambda^\gamma h_y. \quad (1.13)$$

Also, for readability, in our notation *we will not explicitly indicate the dependency of the scaling in (1.13) on x* . The versions of $\psi, \rho, \mathcal{P}_{\lambda\rho}$, and ξ under this re-scaling are determined by the relations

$$\psi^{(\lambda)}(l) := \lambda^\gamma \psi(\lambda^{-\beta} l), \quad (1.14)$$

$$\rho^{(\lambda)}(y', h'_y) := \lambda^{\delta\gamma} \rho(y, h_y), \quad (1.15)$$

$$\mathcal{P}_{\lambda\rho}^{(\lambda)} := \mathcal{P}_{\lambda\rho}^{(\lambda)}[x] := \{(y', h'_y) : (y, h_y) \in \mathcal{P}_{\lambda\rho}\}, \quad (1.16)$$

and

$$\xi^{(\lambda)}((y', h'_y), \{(y'_i, h'_{y_i})\}_{i \geq 1}) := \xi((y, h_y), \{(y_i, h_{y_i})\}_{i \geq 1}) \quad (1.17)$$

and likewise for $\hat{\xi}$. Since $dy' = \lambda^{\beta(d-1)} dy$ and $dh'_y = \lambda^\gamma dh_y$, it follows that

$$\rho^{(\lambda)}(y', h'_y) dy' dh'_y = \lambda \rho(y, h_y) dy dh_y.$$

Note also that

$$\mathcal{P}_{\lambda\rho}^{(\lambda)} \stackrel{\mathcal{D}}{=} \mathcal{P}_{\rho^{(\lambda)}}. \quad (1.18)$$

Moreover, by (1.13) and (1.15) $\rho^{(\lambda)}(y', h'_y)(h'_y)^{-\delta} = \lambda^{\delta\gamma} \rho(y, h_y)(\lambda^\gamma h_y)^{-\delta}$, where $y = \lambda^{-\beta} y' + x$. Under the above re-scaling for each fixed $x \in A$ and for each (y', h'_y) we have the crucial limit

$$\lim_{\lambda \rightarrow \infty} \rho^{(\lambda)}(y', h'_y)(h'_y)^{-\delta} = \lim_{\lambda \rightarrow \infty} \rho(y, h_y)(h_y)^{-\delta} = \rho_0(x) \quad (1.19)$$

and by $(\Psi 2)$ and (1.14) for all $l \in \mathbb{R}_+$

$$\lim_{\lambda \rightarrow \infty} \psi^{(\lambda)}(l) = l^\alpha. \quad (1.20)$$

It is also worth noting that $\xi^{(\lambda)}$ could alternatively be defined by following the original definition of ξ with ψ replaced there by $\psi^{(\lambda)}$; the same applies for $\hat{\xi}^{(\lambda)}$. Observe that in fact it states approximate self-similarity of ψ -growth processes under the re-scaling given by (1.13) and (1.14). Motivated by this observation, we have already put $\psi^{(\infty)}(l) := l^\alpha$ and now we define for all $x \in A$ and for all $(y', h'_y) \in \mathbb{R}^{d-1} \times \mathbb{R}_+$

$$\rho^{(\infty)}(y', h'_y) := \rho_x^{(\infty)}(y', h'_y) := \rho_0(x)(h'_y)^\delta. \quad (1.21)$$

2 Applications

We describe here applications of the main results. We limit the discussion to

- (i) the number of vertices in the convex hull of a Poisson sample, and

(ii) the number of maximal points in a Poisson or i.i.d. sample,

but it should be emphasized that the techniques could potentially be applied to a broader scope of examples. These include for instance the variance asymptotics for Johnson-Mehl growth processes [24] with non-linear growth rates (see e.g. subsection 3.2.2 in [7] for the description of the model and the corresponding central limit theorem. Also, as observed in section 2.3 of [6], the case $\psi(l) = l^2$ (paraboloids) may figure in the limit behavior of some point processes associated with the asymptotic solutions of Burgers equation

$$\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} = \varepsilon \Delta v$$

in the inviscid limit $\varepsilon \rightarrow 0$. We will likewise not treat this example either.

2.1 Number of vertices in the convex hull of an i.i.d. sample

Recall that B_d denotes the unit radius ball centered at the origin of \mathbb{R}^d and let ∂B_d denote its boundary. Let $\rho : B_d \rightarrow \mathbb{R}_+$ be a continuous density on B_d . We shall assume that $\rho(x) = \rho_0(x/|x|)(1 - |x|)^\delta(1 + o(1))$ for some $\delta \geq 0$ and that $\rho_0 : \partial B_d \rightarrow \mathbb{R}_+$ is continuous and bounded away from 0. Let $\mathcal{P}_{\lambda\rho}$ be a Poisson point process on B_d with intensity measure $\lambda\rho(x)dx$ and let $\text{conv}(\mathcal{P}_{\lambda\rho})$ be the random polytope given by the convex hull of $\mathcal{P}_{\lambda\rho}$. Recalling that $\mathcal{V}(\mathcal{P}_{\lambda\rho})$ denotes the vertices of $\text{conv}(\mathcal{P}_{\lambda\rho})$, consider the *vertex empirical point measure*

$$\mu_{\lambda\rho} := \sum_{x \in \mathcal{V}(\mathcal{P}_{\lambda\rho})} \delta_x. \quad (2.1)$$

As will be shown in section four, Theorems 1.1 - 1.3 yield the following limit theory for $\mu_{\lambda\rho}$. Let $N(0, 1)$ denote the standard normal random variable.

Theorem 2.1 *There are constants M and V such that for all $f \in \mathcal{C}_b(B_d)$*

$$\lim_{\lambda \rightarrow \infty} \lambda^{-(d-1)/(d-1+2(1+\delta))} \mathbb{E}[\langle f, \mu_{\lambda\rho} \rangle] = M \int_{\partial B_d} f(s) \rho_0^{(d-1)/(d-1+2(1+\delta))}(s) ds \quad (2.2)$$

and

$$\lim_{\lambda \rightarrow \infty} \lambda^{-(d-1)/(d-1+2(1+\delta))} \text{Var}[\langle f, \mu_{\lambda\rho} \rangle] = V \int_{\partial B_d} f^2(s) \rho_0^{(d-1)/(d-1+2(1+\delta))}(s) ds. \quad (2.3)$$

Moreover, the finite-dimensional distributions $\lambda^{-(d-1)/2(d-1+2(1+\delta))}(\langle f_1, \bar{\mu}_{\lambda\rho} \rangle, \dots, \langle f_k, \bar{\mu}_{\lambda\rho} \rangle)$, $f_i \in \mathcal{C}_b(B_d)$, of $(\lambda^{-(d-1)/2(d-1+2(1+\delta))} \bar{\mu}_{\lambda\rho})$ converge as $\lambda \rightarrow \infty$ to those of a mean zero Gaussian field with covariance kernel

$$(f, g) \mapsto V \int_{\partial B_d} f(s)g(s) \rho_0^{(d-1)/(d-1+2(1+\delta))}(s) ds, \quad f, g \in \mathcal{C}_b(B_d).$$

Additionally, if $\delta = 0$, then for all $f \in \mathcal{C}_b(B_d)$

$$\sup_t \left| P \left[\frac{\langle f, \bar{\mu}_{\lambda\rho} \rangle}{\sqrt{\text{Var}\langle f, \bar{\mu}_{\lambda\rho} \rangle}} \leq t \right] - P[N(0, 1) \leq t] \right| = O \left(\lambda^{-(d-1)/2(d+1)} (\log \lambda)^{3+2(d-1)} \right). \quad (2.4)$$

Remarks. (i) Taking $f_1 \equiv 1$ (and all other $f_i \equiv 0$, $i = 2, \dots, k$) provides a central limit theorem for the cardinality of $\mathcal{V}(\mathcal{P}_{\lambda\rho})$.

(ii) Theorem 2.1 adds to the work of the following authors: (a) Groeneboom [19] and Cabo and Groeneboom [10], who prove a central limit theorem for the cardinality of $\mathcal{V}(\mathcal{P}_{\lambda\rho})$ when ρ is uniform and when $d = 2$, (b) Reitzner [35] who considers the one dimensional central limit theorem and who establishes a rate of convergence $O(\lambda^{-(d-1)/2(d+1)} (\log \lambda)^{2+2/(d+1)})$ to the normal for ρ uniform (whence $\delta = 0$ in our setting), without giving asymptotics for the limiting variance and covariance, and (c) Van Vu [44], who proves a central limit theorem for the cardinality of $\mathcal{V}(\{X_i\}_{i=1}^n)$, X_i i.i.d. uniform, but who also does not consider limiting covariances. Concerning rates, we believe that the power on the logarithm, namely $3 + 2(d - 1)$ can be reduced to $2(d - 1)$ but we have not tried for this sharper rate.

(iii) As shown by Reitzner (Lemma 7 of [35]), when $\delta = 0$ the right-hand side of (2.3) is strictly positive and finite whenever f is not identically zero.

2.2 Number of maximal points in an i.i.d. sample

For all $\bar{w} := (w, h_w)$, we define the downwards cone

$$K^\downarrow[\bar{w}] := \{(z, h_z) \in \mathbb{R}^{d-1} \times \mathbb{R}_+ : h_z \leq h_w - \psi(|z - w|)\}. \quad (2.5)$$

Consider $\psi(l) := l^\alpha$, $\alpha \in (0, 1]$, in subsection 1.1 so that $K[\mathbf{0}] := \{(w_1, \dots, w_d) : w_d \geq (w_1^2 + \dots + w_{d-1}^2)^{\alpha/2}\}$. Given a locally finite set $\mathcal{X} \subset \mathbb{R}^d$, a point $\bar{w} \in \mathcal{X}$ is called *K-maximal* iff \bar{w} does not belong to any $u \oplus K[\mathbf{0}]$ for $u \in \mathcal{X}$. When $\alpha \in (0, 1]$ we have the equivalence $\bar{y} \in K[\bar{x}]$ iff $K[\bar{y}] \subseteq K[\bar{x}]$ and $\bar{x} \in K^\downarrow[\bar{y}]$ iff $K^\downarrow[\bar{x}] \subseteq K^\downarrow[\bar{y}]$. It thus follows that for such ψ the present notion of maximality is just a rephrasing of the maximality notion as discussed in section one. Indeed, we see that \bar{w} is *K-maximal* or ψ -*extremal* in \mathcal{X} iff $\bar{w} \oplus K^\downarrow[\mathbf{0}]$ contains no other points in \mathcal{X} . This is not the case for $\alpha > 1$ where the equivalence $\bar{y} \in K[\bar{x}]$ iff $K[\bar{y}] \subseteq K[\bar{x}]$ does not hold.

Recalling that $\mathcal{M}_K(\mathcal{P}_{\lambda\rho})$ denotes the collection of *K-maximal* points in $\mathcal{P}_{\lambda\rho}$, and with ρ and A as in section 1.1, consider the induced *maximal point measure*

$$\mu_{\lambda\rho} := \sum_{x \in \mathcal{M}_K(\mathcal{P}_{\lambda\rho})} \delta_x.$$

Recalling the definitions of $I(f)$ and $J(f)$ at (1.7) and (1.8), respectively, we have:

Theorem 2.2 *With τ as given by (1.5) and $\alpha \in (0, 1]$, for all $f \in \mathcal{C}_b(A_+)$*

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\tau} \mathbb{E}[\langle f, \mu_{\lambda\rho} \rangle] = I(f) \quad (2.6)$$

and

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\tau} \text{Var}[\langle f, \bar{\mu}_{\lambda\rho} \rangle] = I(f^2) + J(f^2). \quad (2.7)$$

Moreover, the finite-dimensional distributions $(\langle f_1, \lambda^{-\tau/2} \bar{\mu}_{\lambda\rho} \rangle, \dots, \langle f_k, \lambda^{-\tau/2} \bar{\mu}_{\lambda\rho} \rangle)$, $f_1, \dots, f_k \in \mathcal{C}_b(A_+)$, of $\lambda^{-\tau/2} \bar{\mu}_{\lambda\rho}$ converge as $\lambda \rightarrow \infty$ to those of a mean zero Gaussian field with covariance kernel

$$(f, g) \mapsto I(fg) + J(fg), \quad f, g \in \mathcal{C}_b(A_+).$$

Additionally, if $\delta = 0$, then for all $f \in \mathcal{C}_b(A_+)$

$$\sup_t \left| P \left[\frac{\langle f, \bar{\mu}_{\lambda\rho} \rangle}{\sqrt{\text{Var} \langle f, \bar{\mu}_{\lambda\rho} \rangle}} \leq t \right] - P[N(0, 1) \leq t] \right| = O \left(\lambda^{-(d-1)/2d} (\log \lambda)^{3+2(d-1)} \right). \quad (2.8)$$

Theorem 2.2 admits de-Poissonization as follows. Let X_1, \dots, X_n be i.i.d. chosen in A_+ according to the density ρ , assumed to be integrable to 1, and consider the associated maximal point measure

$$\nu_n^\xi := \sum_{x \in \mathcal{M}_\kappa(\{X_i\}_{i=1}^n)} \delta_x.$$

We have then the following equivalent of Theorem 2.2 for binomial samples.

Theorem 2.3 *With τ as given by (1.5) and $\alpha \in (0, 1]$, for all $f \in \mathcal{C}_b(A_+)$*

$$\lim_{n \rightarrow \infty} n^{-\tau} \mathbb{E}[\langle f, \nu_n^\xi \rangle] = I(f) \quad (2.9)$$

and

$$\lim_{n \rightarrow \infty} n^{-\tau} \text{Var}[\langle f, \bar{\nu}_n^\xi \rangle] = I(f^2) + J(f^2). \quad (2.10)$$

Moreover, the finite-dimensional distributions $(\langle f_1, n^{-\tau/2} \bar{\nu}_n^\xi \rangle, \dots, \langle f_k, n^{-\tau/2} \bar{\nu}_n^\xi \rangle)$, $f_1, \dots, f_k \in \mathcal{C}_b(A_+)$, of $n^{-\tau/2} \bar{\nu}_n^\xi$ converge as $n \rightarrow \infty$ to those of a mean zero Gaussian field with covariance kernel

$$(f, g) \mapsto I(fg) + J(fg), \quad f, g \in \mathcal{C}_b(A_+).$$

Remark. Theorems 2.2 and 2.3 extend and generalize the work of (a) Barbour and Xia [4], who establish central limit theorems for the case of homogeneous spatial temporal input, with K the positive octant in \mathbb{R}^d , and who consider neither convergence of finite dimensional distributions nor convergence of variances, (b) Baryshnikov and Yukich [7], who establish convergence of finite dimensional distributions but who restrict to homogeneous temporal input ($\delta = 0$) as well as to the case $\psi(l) = l$ (i.e. $\alpha = 1$), and (c) Baryshnikov [6], who also restricts to homogeneous temporal input and does not consider convergence of finite dimensional distributions.

3 Proof of Main Results

In this section we prove Theorems 1.1 - 1.3. An essential component of the proofs involves introducing a notion of *localization*, which quantifies the decoupling property of the considered functional ξ over distant regions. It is straightforward to check that the proofs hold for ψ -growth without overlap when ψ is linear.

3.1 Stabilization for Ψ -growth functionals

With $B_{d-1}(y, r)$ standing as usual for the $(d-1)$ -dimensional ball centered at $y \in \mathbb{R}^{d-1}$ with radius $r \in (0, \infty)$, we denote by $C_{d-1}(y, r)$ the cylinder $B_{d-1}(y, r) \times \mathbb{R}_+$. Recalling $\bar{y} := (y, h_y)$, consider for all $r > 0$ the finite range version of $\xi(\bar{y}, \mathcal{X})$, namely

$$\xi_{[r]}(\bar{y}, \mathcal{X}) := \xi_{C_{d-1}(y, r)}(\bar{y}, \mathcal{X}),$$

that is $\xi_{[r]}(\bar{y}, \mathcal{X})$ depends only on the local behavior of \mathcal{X} with *spatial* coordinates restricted to the r -neighborhood of y . For a point process \mathcal{P} (usually chosen to be Poisson in the sequel) in $\mathbb{R}^{d-1} \times \mathbb{R}_+$, the *localization radius* of ξ at $\bar{y} \in \mathbb{R}^{d-1} \times \mathbb{R}_+$ is defined by

$$R^\xi := R^\xi[\bar{y}; \mathcal{P}] := \inf\{r \in \mathbb{R}_+ : \forall s \geq r \ \xi(\bar{y}, \mathcal{P}) = \xi_{[s]}(\bar{y}, \mathcal{P})\}. \quad (3.1)$$

In full analogy with $\xi^{(\lambda)}$ given by (1.17), we define for all $\lambda > 0$ the localization radius $R^{\xi^{(\lambda)}}[\cdot; \cdot]$ by

$$R^{\xi^{(\lambda)}} := R^{\xi^{(\lambda)}}[\bar{y}; \mathcal{P}'] := \inf\{r \in \mathbb{R}_+ : \forall s \geq r \ \xi^{(\lambda)}(\bar{y}', \mathcal{P}') = \xi_{[s]}^{(\lambda)}(\bar{y}', \mathcal{P}')\}.$$

Observe that the localization radius considered here formally differs from the stabilization radii considered in [7], [26] - [29], essentially defined for all $\bar{y} := (y, h)$ to be the smallest positive real r such that $\xi(\bar{y}, (\mathcal{P} \cap C_{d-1}(y, r)) \cup \mathcal{A}) = \xi(\bar{y}, (\mathcal{P} \cap C_{d-1}(y, r)))$ for all finite $\mathcal{A} \subset C_{d-1}^c(y, s)$.

However the ψ -extremal functional is in general extremely sensitive to the choice of the ‘outside’ configuration $\mathcal{A} \subset C_{d-1}^c(y, s)$, rendering the existence and use of standard stabilization radii a bit difficult. The benefit of the localization radius is that it considers only the outside configurations involving points from \mathcal{P} . However, since the localization radius shares many of the same properties as the stabilization radii in [7], [26] - [29], *we will abuse terminology and henceforth refer to the localization radius R^ξ as a stabilization radius.*

The following lemma shows that $\xi^{(\lambda)}$ given by (1.17) has a stabilization radius whose tail decays exponentially uniformly in large enough λ when \mathcal{P} is $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ given by (1.16) or when \mathcal{P} is given by $\mathcal{P}_{\lambda\rho}^{(\lambda)} \cup \{\bar{z}'_1, \dots, \bar{z}'_k\}$, $k \geq 1$, where \bar{z}'_i , $i = 1, \dots, k$ are certain deterministic points (fixed atoms). This result will prove useful later in showing exponential decay of correlation functions for ψ -growth processes.

Lemma 3.1 (i) *For A compact and convex there exists a constant C such that, uniformly in x and λ large enough, for all $\bar{y}' \in \lambda^\beta A \times \mathbb{R}_+$ and for all collections $\{\bar{z}'_1, \dots, \bar{z}'_k\} \subseteq \lambda^\beta A \times \mathbb{R}_+$ of deterministic points, $k \geq 0$, we have for all $L > 0$*

$$P \left[R^{\xi^{(\lambda)}}[\bar{y}'; \mathcal{P}_{\lambda\rho}^{*(\lambda)}] > L \right] \leq C \exp \left(-\frac{L^{\alpha+d-1}}{C} \right), \quad (3.2)$$

where $\mathcal{P}_{\lambda\rho}^{*(\lambda)} := \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup \{\bar{z}_1, \dots, \bar{z}_k\}$ so that in particular $\mathcal{P}_{\lambda\rho}^{*(\lambda)} = \mathcal{P}_{\lambda\rho}^{(\lambda)}$ for $k = 0$.

(ii) *An identical bound holds if instead $A := \mathbb{R}^{d-1}$ and $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ is replaced by a homogeneous Poisson point process on \mathbb{R}^{d-1} .*

Remark. In place of (3.2) we have uniformly in x and λ large enough, for all $\bar{y}' \in \lambda^\beta A \times \mathbb{R}_+$ and for all $L > 0$ the simpler bound

$$P \left[R^{\xi^{(\lambda)}}[\bar{y}'; \mathcal{P}_{\lambda\rho}^{*(\lambda)}] > L \right] \leq C \exp \left(-\frac{L}{C} \right). \quad (3.3)$$

Proof. We will only prove Lemma 3.1(i) as identical arguments handle Lemma 3.1(ii). Also, since the proof relies on probability bounds for certain regions being devoid of points of the underlying point process $\mathcal{P}_{\lambda\rho}^{*(\lambda)}$, as easily noted below, we can assume without loss of generality that $k = 0$ so that $\mathcal{P}_{\lambda\rho}^{*(\lambda)} = \mathcal{P}_{\lambda\rho}^{(\lambda)}$. Moreover, to simplify the argument below, we ignore the boundary effects arising when \bar{y}' is close to $\partial(\lambda^\beta A \times \mathbb{R}_+)$, noting that the absence of points of $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ in the vicinity of \bar{y}' can only decrease $R^{\xi^{(\lambda)}}[\bar{y}'; \mathcal{P}_{\lambda\rho}^{(\lambda)}]$. This allows us to avoid obvious but technical separate considerations for \bar{y}' close to $\partial(\lambda^\beta A \times \mathbb{R}_+)$. Also, we consider x fixed but arbitrary, keeping in mind that the required uniformity in x follows by the boundedness of ρ , both from above and away from 0.

Define for fixed $\bar{y}' := (y', h'_y)$ and all $\lambda \in [0, \infty]$ the *scaled upwards cone*

$$K^{(\lambda)}[\bar{y}'] := \{(v', h'_v) \in \mathbb{R}^{d-1} \times \mathbb{R}_+ : h'_v \geq h'_y + \psi^{(\lambda)}(|v' - y'|)\} \quad (3.4)$$

and the *scaled downwards cone*

$$K_{(\lambda)}^\downarrow[\bar{y}'] := \{(v', h'_v) \in \mathbb{R}^{d-1} \times \mathbb{R}_+ : h'_v \leq h'_y - \psi^{(\lambda)}(|v' - y'|)\}. \quad (3.5)$$

Note that $\bar{u}' \in K^{(\lambda)}[\bar{z}']$ iff $h'_u \geq h'_z + \psi(|u' - z'|)$ which is equivalent to $h'_z \leq h'_u - \psi(|z' - u'|)$, and thus the *duality* $\bar{u}' \in K^{(\lambda)}[\bar{z}']$ iff $\bar{z}' \in K_{(\lambda)}^\downarrow[\bar{u}']$.

To proceed, note that the event $\{R^{\xi^{(\lambda)}}[\bar{y}'; \mathcal{P}_{\lambda\rho}^{(\lambda)}] > L\}$ is equivalent to the event

$$E := \left\{ \exists r > L : \xi^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{(\lambda)}) \neq \xi_{[r]}^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{(\lambda)}) \right\},$$

and moreover $E \subset E_1 \cup E_2$, where E_1 and E_2 are defined below. Roughly speaking, the event E_1 ensures that \bar{y}' is extremal with respect to $\mathcal{P}_{\lambda\rho}^{(\lambda)} \cap C_{d-1}(y', r)$ for some $r > L$ but not necessarily with respect to $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ whereas E_2 is just the opposite.

Event E_1 : For some $r > L$ there exists a boundary point $\bar{u}' \in \partial(K^{(\lambda)}[\bar{y}']) \cap C_{d-1}(y', r)$, and such that $\bar{u}' \notin \bigcup_{\bar{z}' \in [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] \cap C_{d-1}(y', r)} K^{(\lambda)}[\bar{z}']$ but $\bar{u}' \in \bigcup_{\bar{z}' \in \mathcal{P}_{\lambda\rho}^{(\lambda)} \cap C_{d-1}(y', r)} K^{(\lambda)}[\bar{z}']$, i.e., $\xi_{[r]}^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{(\lambda)}) = 1$, but possibly $\xi^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{(\lambda)}) = 0$.

Event E_2 : For some $r > L$ there exists a boundary point $\bar{u}' \in \partial(K^{(\lambda)}[\bar{y}']) \cap C_{d-1}^c(y', r)$ such that $\bar{u}' \notin \bigcup_{\bar{z}' \in \mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}} K^{(\lambda)}[\bar{z}']$, but $K^{(\lambda)}[\bar{y}'] \cap C_{d-1}(y', r) \subset \bigcup_{\bar{z}' \in [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] \cap C_{d-1}(y', r)} K^{(\lambda)}[\bar{z}']$, i.e., $\xi^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{(\lambda)}) = 1$ but $\xi_{[r]}^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{(\lambda)}) = 0$.

On event E_1 writing $\bar{u}' := (u', h'_u)$, we easily check that

$$h'_u \geq \psi^{(\lambda)}\left(\frac{L}{2}\right). \quad (3.6)$$

Indeed, we have

- either $|u' - y'| \geq r/2$ or
- $d(u', \partial B_{d-1}(y', r)) \geq r/2$ and hence $d(u', z') \geq r/2$ for all $\bar{z}' \in \mathcal{P}_{\lambda\rho}^{(\lambda)} \cap C_{d-1}^c(y', r)$.

In both cases, on E_1 , (u', h'_u) falls into $K^{(\lambda)}[\bar{v}']$ for some \bar{v}' such that $|v' - u'| \geq r/2$, either with $\bar{v}' = \bar{y}'$ or $\bar{v}' \in \mathcal{P}_{\lambda\rho}^{(\lambda)} \cap C_{d-1}^c(y', r)$. Consequently, recalling that $r > L$ and using the definition of $K^{(\lambda)}[\cdot]$ we obtain (3.6) as required.

On E_1 we have $\bar{u}' \notin \bigcup_{\bar{z}' \in [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] \cap C_{d-1}(y', r)} K^{(\lambda)}[\bar{z}']$ implying that the downward cone $K_{(\lambda)}^\downarrow[\bar{u}']$ is devoid of points of $\mathcal{P}_{\lambda\rho}^{(\lambda)} \cap C_{d-1}(y', r)$. By the assumed properties of ψ and ρ , the integral of $\rho^{(\lambda)}$ over $K_{(\lambda)}^\downarrow[\bar{u}']$ is $\Omega(\text{Vol}(K_{(\lambda)}^\downarrow[\bar{u}']))$ which is

$$\Omega \left(\int_0^{h'_u} ([\psi^{(\lambda)}]^{-1}(h'_u - h'))^{d-1} dh' \right) = \Omega \left(\int_0^{h'_u} (h'_u - h')^{(d-1)/\alpha} dh' \right) = \Omega((h'_u)^{(\alpha+d-1)/\alpha}), \quad (3.7)$$

with the second equality following by the definition of $[\psi^{(\lambda)}]^{-1}$, and where we use $f(\lambda) = \Omega(g(\lambda))$ to signify that $f(\lambda)/g(\lambda)$ is asymptotically bounded away from zero. Clearly the integral of $\rho^{(\lambda)}$ over $K_{(\lambda)}^\downarrow[\bar{u}'] \cap C_{d-1}(y', r)$ for $\bar{u}' \in C_{d-1}(y', r)$ is of the same order.

Recalling from (1.18) that the intensity measure of the Poisson process $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ has its density given by $\rho^{(\lambda)}$, we thus conclude for fixed \bar{u}' that the probability of the considered event $\Xi[\bar{u}'] := \{K_{(\lambda)}^\downarrow[\bar{u}'] \cap [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] \cap C_{d-1}(y', r) = \emptyset\}$ satisfies

$$P[\Xi[\bar{u}']] \leq \exp(-\Omega((h'_u)^{(\alpha+d-1)/\alpha})). \quad (3.8)$$

To proceed, we recall that $r > L$ and we partition $\mathbb{R}^{d-1} \times \mathbb{R}_+$ into unit volume cubes and we let q_1, q_2, \dots be an enumeration of those cubes having non-empty intersection with $\partial(K^{(\lambda)}[\bar{y}'])$. Let

$$p_i := P \left[\exists \bar{u}' \in q_i : K_{(\lambda)}^\downarrow[\bar{u}'] \cap [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] \cap C_{d-1}(y', L) = \emptyset \right]$$

for all $i = 1, 2, \dots$ and note that by (3.8) we have

$$p_i \leq \exp(-\Omega((h'_q)^{(\alpha+d-1)/\alpha}))$$

where h'_q is the last coordinate of the center of the cube q_i .

We now have

$$P[E_1] \leq \sum_{i=1}^{\infty} p_i \leq C \int_{\psi^{(\lambda)}(L/2)}^{\infty} L^{d-2} \exp\left(-\frac{1}{C}(h'_u)^{(\alpha+d-1)/\alpha}\right) dh'_u$$

for some $0 < C < \infty$ in view of the discussion above. Here CL^{d-2} bounds the number of cubes in the set q_1, q_2, \dots of any fixed height $h'_u \geq \psi^{(\lambda)}(L/2)$.

Recalling that $\psi^{(\lambda)}(L/2) = (1+o(1))(L/2)^\alpha$, it follows (using a different choice of C if necessary) that

$$P[E_1] \leq C \exp\left(-\frac{1}{C}L^{\alpha+d-1}\right).$$

To estimate $P[E_2]$, note that for $\bar{u}' := (u', h'_u) \in \partial(K[\bar{y}'])$ lying in $C_{d-1}^c(y', r)$ we must have

$$h'_u \geq \psi^{(\lambda)}(r). \quad (3.9)$$

Further, since $\bar{u}' \notin \bigcup_{\bar{z}' \in \mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}} K^{(\lambda)}[\bar{z}']$ we have $K_{(\lambda)}^\downarrow[\bar{u}'] \cap [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] = \emptyset$. Denoting this event $\Xi^*[\bar{u}'] := \{K_{(\lambda)}^\downarrow[\bar{u}'] \cap [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] = \emptyset\}$, noting that as in (3.8) we have

$$P[\Xi^*[\bar{u}']] \leq \exp(-\Omega((h'_u)^{(\alpha+d-1)/\alpha})), \quad (3.10)$$

recalling that $r > L$ and proceeding in analogy with the case of event E_1 above, with (3.6) and (3.8) there replaced by (3.9) and (3.10) respectively and with $C_{d-1}^c(y', L)$ partitioned into unit volume cubes, we bound $P[E_2]$ by

$$P[E_2] \leq C \int_{s=L}^{\infty} s^{d-2} \int_{h'_y + \psi^{(\lambda)}(s)}^{\infty} \exp\left(-\frac{1}{C}(h'_u)^{(\alpha+d-1)/\alpha}\right) dh'_u ds$$

for some $0 < C < \infty$. It follows that $P[E_2] \leq C \exp(-L^{\alpha+d-1}/C)$. Since $P[R^{\xi^{(\lambda)}}[\bar{y}'; \mathcal{P}_{\lambda\rho}^{(\lambda)}] > L] = P[E] \leq P[E_1] + P[E_2]$, Lemma 3.1 follows. \square

Given $\bar{y} := (y', h'_y)$ we expect for large temporal h'_y that is ψ -extremal with small probability. Also, as previously noted in subsection 1.1, we expect for small α that \bar{y} is more likely to be ψ -extremal. The next lemma makes these probabilities a bit more precise and shows that the probability of having (y', h'_y) extreme in $\mathcal{P}_{\lambda\rho}^{*(\lambda)} := \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup \{\bar{z}'_1, \dots, \bar{z}'_k\}$, $k \geq 0$, with respect to $\psi^{(\lambda)}$ decays exponentially with h'_y uniformly in λ for λ large enough.

Lemma 3.2 *There exists a constant C such that, uniformly in λ large enough, for all $\bar{y}' \in \lambda^\beta A \times \mathbb{R}_+$ and $\{\bar{z}'_1, \dots, \bar{z}'_k\}$ we have*

$$P\left[\xi^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{*(\lambda)}) = 1\right] \leq C \exp\left(-\frac{1}{C}(h'_y)^{(\alpha+d-1)/\alpha}\right).$$

Proof. Clearly, since adding extra points to $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ decreases the probability of (y', h'_y) being extreme, we may without loss of generality choose $k = 0$ so that $\mathcal{P}_{\lambda\rho}^{*(\lambda)} = \mathcal{P}_{\lambda\rho}^{(\lambda)}$.

On the event $E := \{\xi^{(\lambda)}(\bar{y}', \mathcal{P}_{\lambda\rho}^{(\lambda)}) = 1\}$ there exists $\bar{u}' := (u', h'_u) \in \partial(K^{(\lambda)}[\bar{y}'])$ such that $\bar{u}' \notin \bigcup_{\bar{z}' \in \mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}} K^{(\lambda)}[\bar{z}']$, which is equivalent to $K_{(\lambda)}^\downarrow[\bar{u}'] \cap [\mathcal{P}_{\lambda\rho}^{(\lambda)} \setminus \{\bar{y}'\}] = \emptyset$. As in the proof of Lemma 3.1, for fixed \bar{u}' the probability of the last event does not exceed

$$\exp\left[-\int_{K_{(\lambda)}^\downarrow[\bar{u}']} \rho^{(\lambda)}(v' h'_v) dv' dh'_v\right] \leq C \exp\left(-\frac{1}{C}(h'_u)^{(\alpha+d-1)/\alpha}\right).$$

Recalling the relation $h'_u = h'_y + \psi^{(\lambda)}(|u' - y'|)$, putting $|u' - y'| = s$, and resorting again to a partition of $\mathbb{R}^{d-1} \times \mathbb{R}_+$ into unit volume cubes and summing up the respective probabilities as in the proof of Lemma 3.1, we obtain the required bound

$$P[E] \leq C \int_0^{\infty} s^{d-2} \int_{h'_y}^{\infty} \exp\left(-\frac{1}{C}(h'_u)^{(\alpha+d-1)/\alpha}\right) dh'_u ds \leq C \exp\left(-\frac{1}{C}(h'_y)^{(\alpha+d-1)/\alpha}\right).$$

\square

3.2 Proof of Theorem 1.1

Recall the definition of $\mathcal{P}_{\rho_x^{(\infty)}}$ from (1.21). One benefit of stabilization is that the one point correlation function $\mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right]$ is approximated for large r by the finite range version

$$\mathbb{E} \left[\xi_{[r]}^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right]$$

and similarly $\mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right]$ is approximated by its finite range version $\mathbb{E} \left[\xi_{[r]}^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right]$.

Using the large λ weak convergence of $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ to $\mathcal{P}_{\rho_x^{(\infty)}}$, one may approximate the first mentioned finite range version by the second and thus show that $\mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right]$ is asymptotically equal to $\mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right]$. This is spelled out in Lemma 3.3 below, which captures the essence of stabilization and which lies at the heart of the proof of Theorem 1.1. Note that when Lemma 3.3 is combined with Lemma 3.2, then it shows

$$\mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}^* \right) \right] \leq C \exp \left(-\frac{1}{C} (h')^{(\alpha+d-1)/\alpha} \right) \quad (3.11)$$

and therefore $I(f) < \infty$ for $f \in \mathcal{C}_b(A_+)$. Recall from (1.17) that $\xi^{(\lambda)}$ is the re-scaled version of ξ with dependency on x fixed.

Lemma 3.3 *For all $x \in A$ and $h' \in \mathbb{R}_+$ we have*

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] = \mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right].$$

Proof. Fix $x \in A$. Taking into account (1.19) and (1.21) and using the results of Section 3.5 in [38] (see Proposition 3.22 or Proposition 3.19 there combined with Proposition 3.6 (ii) ibidem) we observe that as $\lambda \rightarrow \infty$, $\mathcal{P}_{\lambda\rho}^{(\lambda)}$ converges weakly to $\mathcal{P}_{\rho_x^{(\infty)}}$ as a point process, see ibidem. Using Theorem 5.5 in [9] with $h_\lambda := \xi_{[r]}^{(\lambda)}((\mathbf{0}, h'), \cdot)$ and $h := \xi_{[r]}^{(\infty)}((\mathbf{0}, h'), \cdot)$ there we easily see that, by Lemma 3.1 and 3.2, under the law of the limit process $\mathcal{P}_{\rho_x^{(\infty)}}$ the discontinuity event E ibidem (an infinitesimal move of the point configuration alters the ξ -value for $(\mathbf{0}, h)$) is contained up to an event of probability 0 in the set of point configurations \mathcal{X} such that either the spatial coordinates of two points in \mathcal{X} coincide or such that there are at least two points $\bar{y}', \bar{y}'' \in \mathcal{X}$ such that the boundaries of the upwards cones $K^{(\infty)}[\bar{y}']$ and $K^{(\infty)}[\bar{y}'']$ (recall (3.4)), intersect in a point lying on the boundary of the upwards cone $K^{(\infty)}[(\mathbf{0}, h')]$, which clearly happens with probability 0 under the law of $\mathcal{P}_{\rho_x^{(\infty)}}$. Indeed, Lemma 3.2 states that no effects coming from $h \rightarrow \infty$ arise (no infinite range dependencies in h). A similar statement in space is provided by Lemma 3.1. Combining both these statements allows us to draw conclusions from the weak convergence of point processes

as we do in the above argument, see *ibidem* in [38]. Thus, Theorem 5.5 in [9] yields

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \left[\xi_{[r]}^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] = \mathbb{E} \left[\xi_{[r]}^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right]. \quad (3.12)$$

Let $R^\xi := R^{\xi^{(\lambda)}}[(\mathbf{0}, h'); \mathcal{P}_{\lambda\rho}^{(\lambda)}]$. We have for all $r > 0$ and all $\lambda > 0$

$$\begin{aligned} \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] &= \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \mathbf{1}_{R^\xi \leq r} \right] + \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \mathbf{1}_{R^\xi > r} \right] \\ &= \mathbb{E} \left[\xi_{[r]}^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \mathbf{1}_{R^\xi \leq r} \right] + \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \mathbf{1}_{R^\xi > r} \right]. \end{aligned}$$

By Lemma 3.1(i) (recall the bound (3.3)), Cauchy-Schwarz, and the boundedness of $\xi_{[r]}^{(\lambda)}$, uniformly in large λ and all $r > 0$

$$\mathbb{E} \left[\xi_{[r]}^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \mathbf{1}_{R^\xi > r} \right] \leq C \exp \left(-\frac{r}{C} \right)$$

for some C not depending on x . Likewise, uniformly in large λ , we have $\mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \mathbf{1}_{R^\xi > r} \right] \leq C \exp(-r/C)$. It follows that for large $\lambda > 0$ and all $r > 0$

$$\left| \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] - \mathbb{E} \left[\xi_{[r]}^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] \right| \leq 2C \exp \left(-\frac{r}{C} \right). \quad (3.13)$$

Similarly Lemma 3.1(ii) gives for all $r > 0$

$$\left| \mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right] - \mathbb{E} \left[\xi_{[r]}^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right] \right| \leq 2C \exp \left(-\frac{r}{C} \right).$$

Write

$$\begin{aligned} & \left| \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] - \mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right] \right| \\ & \leq \left| \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] - \mathbb{E} \left[\xi_{[r]}^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] \right| \\ & + \left| \mathbb{E} \left[\xi_{[r]}^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \right] - \mathbb{E} \left[\xi_{[r]}^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right] \right| \\ & + \left| \mathbb{E} \left[\xi_{[r]}^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right] - \mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \right] \right|. \end{aligned} \quad (3.14)$$

For fixed r , the second term on the right hand side of (3.14) goes to zero as $\lambda \rightarrow \infty$ by (3.12). The first and third terms are bounded above by $2C \exp(-r/C)$. Letting $r \rightarrow \infty$ completes the proof of Lemma 3.3. \square

Given Lemmas 3.2 and 3.3 we now prove Theorem 1.1 as follows. We have

$$\mathbb{E} [\langle f, \mu_{\lambda\rho}^\xi \rangle] = \int_A \int_0^\infty f(x, h_x) \mathbb{E} [\xi((x, h_x), \mathcal{P}_{\lambda\rho})] \lambda \rho(x, h_x) dh_x dx.$$

By (1.17) we have $\xi((x, h_x), \mathcal{P}_{\lambda\rho}) = \xi^{(\lambda)}((\mathbf{0}, h'_x), \mathcal{P}_{\lambda\rho}^{(\lambda)})$ and by (1.15) we have $\rho(x, h_x) = \lambda^{-\gamma\delta}\rho^{(\lambda)}(\mathbf{0}, h'_x)$. Thus putting $h'_x := \lambda^\gamma h_x$ and recalling $1 - \gamma(\delta + 1) = \tau$ (see (1.5) and (1.12)), we obtain

$$\mathbb{E}[\langle f, \mu_{\lambda\rho}^\xi \rangle] = \int_A \int_0^\infty f(x, h'_x \lambda^{-\gamma}) \mathbb{E}[\xi^{(\lambda)}((\mathbf{0}, h'_x), \mathcal{P}_{\lambda\rho}^{(\lambda)})] \lambda^\tau \rho^{(\lambda)}(\mathbf{0}, h'_x) dh'_x dx,$$

or simply

$$\lambda^{-\tau} \mathbb{E}[\langle f, \mu_{\lambda\rho}^\xi \rangle] = \int_A \int_0^\infty f(x, h'_x \lambda^{-\gamma}) \mathbb{E}[\xi^{(\lambda)}((\mathbf{0}, h'_x), \mathcal{P}_{\lambda\rho}^{(\lambda)})] \rho^{(\lambda)}(\mathbf{0}, h'_x) dh'_x dx.$$

We put

$$g_\lambda(x, h'_x) := \mathbb{E}[\xi^{(\lambda)}((\mathbf{0}, h'_x), \mathcal{P}_{\lambda\rho}^{(\lambda)})] \rho^{(\lambda)}(\mathbf{0}, h'_x).$$

For all $x \in A$ and $h'_x \in \mathbb{R}_+$, we have by Lemma 3.3 and (1.19)

$$\lim_{\lambda \rightarrow \infty} g_\lambda(x, h'_x) = \mathbb{E}[\xi^{(\infty)}((\mathbf{0}, h'_x), \mathcal{P}_{\rho_x^{(\infty)}})] \rho_0(x) h_x'^\delta$$

and moreover, by Lemma 3.2 for all $(x, h) \in A_+$, $g_\lambda(x, h'_x)$ is bounded uniformly in λ by the function $(x, h') \mapsto C'(h')^\delta \exp(-h'/C)$, which is integrable on A_+ . Consequently, the dominated convergence theorem yields

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\tau} \mathbb{E}[\langle f, \mu_{\lambda\rho}^\xi \rangle] = \int_A \int_0^\infty f(x, 0) \mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \cup \{(\mathbf{0}, h')\} \right) \right] \rho_0(x) (h')^\delta dh' dx. \quad (3.15)$$

Using the scaling relations (1.13, 1.14) and (1.6, 1.21) we see that

$$\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \cup \{(\mathbf{0}, h')\} \right) \stackrel{\mathcal{D}}{=} \xi^{(\infty)} \left((\mathbf{0}, [\rho_0(x)]^\gamma h'), \mathcal{P}_* \cup \{(\mathbf{0}, [\rho_0(x)]^\gamma h')\} \right), \quad (3.16)$$

with $\stackrel{\mathcal{D}}{=}$ standing for equality in law. Theorem 1.1 follows by using (3.16), changing variables $h'' := [\rho_0(x)]^\gamma h'$ in the integral in (3.15) and recalling that $\tau = 1 - \gamma(\delta + 1)$. \square

3.3 Proof of Theorem 1.2

Fix $x \in A$ and recall from (1.16) that $\mathcal{P}_{\lambda\rho}^{(\lambda)} := \mathcal{P}_{\lambda\rho}^{(\lambda)}[x]$. For all $\lambda > 0$, $h' \in \mathbb{R}_+$, and $(y', h'_y) \in \lambda^\beta A \times \mathbb{R}_+$, consider the pair correlation function for the re-scaled growth process:

$$\begin{aligned} c^{(\lambda)}((\mathbf{0}, h'), (y', h'_y)) &:= c_x^{(\lambda)}((\mathbf{0}, h'), (y', h'_y)) := & (3.17) \\ \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y) \right) \xi^{(\lambda)} \left((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h') \right) \right] &- \\ \mathbb{E} \xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right) \mathbb{E} \xi^{(\lambda)} \left((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \right). & \end{aligned}$$

Consider also the pair correlation function for the limit growth process $\xi^{(\infty)}$:

$$\begin{aligned} c_x^{(\infty)}((\mathbf{0}, h'), (y', h'_y)) &:= \\ \mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h), \mathcal{P}_{\rho_x^{(\infty)}} \cup (y', h'_y) \right) \xi^{(\infty)} \left((y', h'_y), \mathcal{P}_{\rho_x^{(\infty)}} \cup (\mathbf{0}, h') \right) \right] - \\ \mathbb{E} \xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \right) \mathbb{E} \xi^{(\infty)} \left((y', h'_y), \mathcal{P}_{\rho_x^{(\infty)}} \right). \end{aligned}$$

A second benefit of stabilization, as shown by the next lemma, is that it facilitates convergence of pair correlation functions and thus leads to variance asymptotics. The next lemma is the second order counterpart to Lemma 3.3.

Lemma 3.4 (convergence of two point correlation function) *For all $(x, h_x) := (x, h) \in A_+$, and $(y', h'_y) \in \lambda^\beta A \times \mathbb{R}_+$ we have*

$$\lim_{\lambda \rightarrow \infty} c_x^{(\lambda)}((\mathbf{0}, h'), (y', h'_y)) = c_x^{(\infty)}((\mathbf{0}, h'), (y', h'_y)).$$

Proof. In view of Lemma 3.3, it will suffice to show

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \mathbb{E} \left[\xi^{(\lambda)} \left((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y) \right) \xi^{(\lambda)} \left((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h') \right) \right] \\ = \mathbb{E} \left[\xi^{(\infty)} \left((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \cup (y', h'_y) \right) \xi^{(\infty)} \left((y', h'_y), \mathcal{P}_{\rho_x^{(\infty)}} \cup (\mathbf{0}, h') \right) \right]. \end{aligned} \quad (3.18)$$

Let $R^\xi := R^{\xi^{(\lambda)}}[(\mathbf{0}, h'); \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)]$ and let $R_{y'}^\xi := R^{\xi^{(\lambda)}}[(y', h'_y); \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')]$. For all $r > 0$ we let $E_r := \{R_{y'}^\xi \leq r, R^\xi \leq r\}$. We split the left hand side of (3.18) as

$$\begin{aligned} \mathbb{E} \left[\xi^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \mathbf{1}_{E_r} \right] + \\ \mathbb{E} \left[\xi^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \mathbf{1}_{E_r^c} \right]. \end{aligned}$$

The second expectation is bounded by $C \exp(-r/C)$ for some C not depending on x by Lemma 3.1(i) and by Cauchy-Schwarz. By the definition of the stabilization radius, the first is simply

$$\mathbb{E} \left[\xi_{[r]}^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi_{[r]}^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \mathbf{1}_{E_r} \right].$$

Again, by Lemma 3.1(i) and by Cauchy-Schwarz, for all $r > 0$, this is within $C \exp(-r/C)$ of

$$\mathbb{E} \left[\xi_{[r]}^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi_{[r]}^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \right],$$

hence

$$\begin{aligned} & \left| \mathbb{E} \left[\xi^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \right] - \right. \\ & \left. \mathbb{E} \left[\xi_{[r]}^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi_{[r]}^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \right] \right| \leq 2C \exp\left(\frac{-r}{C}\right) \end{aligned} \quad (3.19)$$

uniformly in x . Now, in analogy with (3.12) we have

$$\begin{aligned} & \lim_{\lambda \rightarrow \infty} \mathbb{E} \left[\xi^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi_{[r]}^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \right] = \\ & \mathbb{E} \left[\xi_{[r]}^{(\infty)}((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \cup (y', h'_y)) \xi_{[r]}^{(\infty)}((y', h'_y), \mathcal{P}_{\rho_x^{(\infty)}} \cup (\mathbf{0}, h')) \right]. \end{aligned} \quad (3.20)$$

By Lemma 3.1(ii) we have for all $r > 0$

$$\begin{aligned} & \left| \mathbb{E} \left[\xi^{(\infty)}((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \cup (y', h'_y)) \xi^{(\infty)}((y', h'_y), \mathcal{P}_{\rho_x^{(\infty)}} \cup (\mathbf{0}, h')) \right] - \right. \\ & \left. \mathbb{E} \left[\xi_{[r]}^{(\infty)}((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \cup (y', h'_y)) \xi_{[r]}^{(\infty)}((y', h'_y), \mathcal{P}_{\rho_x^{(\infty)}} \cup (\mathbf{0}, h')) \right] \right| \leq 2C \exp\left(-\frac{r}{C}\right) \end{aligned} \quad (3.21)$$

as in (3.19). Again, note that C does not depend on x since $\rho_0(x)$ is bounded away from zero.

Combining (3.19), (3.20) and (3.21) yields

$$\begin{aligned} & \limsup_{\lambda \rightarrow \infty} \left| \mathbb{E} \left[\xi^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \right] - \right. \\ & \left. \mathbb{E} \left[\xi^{(\infty)}((\mathbf{0}, h'), \mathcal{P}_{\rho_x^{(\infty)}} \cup (y', h'_y)) \xi^{(\infty)}((y', h'_y), \mathcal{P}_{\rho_x^{(\infty)}} \cup (\mathbf{0}, h')) \right] \right| \\ & \leq 4C \exp\left(-\frac{r}{C}\right) \end{aligned}$$

for all $r > 0$. We conclude the proof of Lemma 3.4 by letting $r \rightarrow \infty$. \square

Lemma 3.4 is not enough to establish second order asymptotics. We will also need that $c_x^{(\lambda)}$ is bounded by an integrable function on $A_+ \times \lambda^\beta A \times \mathbb{R}_+$, i.e., we will need to establish the exponential decay of the correlation function (3.17). This is done in the following lemma, which combined with Lemma 3.4, shows that

$$|c_x^{(\infty)}((\mathbf{0}, h'), (y', h'_y))| \leq C \exp\left(-\frac{1}{C} \max\left(\frac{|y'|}{2}, h'_y, h'\right)\right) \quad (3.22)$$

and therefore $J(f) < \infty$ for all $f \in \mathcal{C}_b(A_+)$.

Lemma 3.5 *There exists a constant C such that for all $\lambda > 0$, $(x, h_x) := (x, h) \in A_+$, and $(y', h'_y) \in \lambda^\beta A \times \mathbb{R}_+$, we have*

$$|c_x^{(\lambda)}((\mathbf{0}, h'), (y', h'_y))| \leq C \exp\left(-\frac{1}{C} \max\left(\frac{|y'|}{2}, h'_y, h'\right)\right).$$

Proof. Let $r \leq |y'|/2$ and note that by definition of $\xi_{[r]}^{(\lambda)}$, we have

$$\begin{aligned} \mathbb{E} \left[\xi_{[r]}^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (y', h'_y)) \xi_{[r]}^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)} \cup (\mathbf{0}, h')) \right] = \\ \mathbb{E} \left[\xi_{[r]}^{(\lambda)}((\mathbf{0}, h'), \mathcal{P}_{\lambda\rho}^{(\lambda)}) \right] \mathbb{E} \left[\xi_{[r]}^{(\lambda)}((y', h'_y), \mathcal{P}_{\lambda\rho}^{(\lambda)}) \right]. \end{aligned}$$

Recalling (3.13) and (3.19) we see that

$$|c_x^{(\lambda)}((\mathbf{0}, h'), (y', h'_y))| \leq 4C \exp\left(-\frac{r}{C}\right)$$

for all $r \leq |y'|/2$. In other words, putting $r = |y'|/2$ yields for all $(x, h) \in A_+$ and $(y', h'_y) \in \lambda^\beta A \times \mathbb{R}_+$

$$|c_x^{(\lambda)}((\mathbf{0}, h'), (y', h'_y))| \leq C \exp\left(-\frac{|y'|}{2C}\right).$$

Appealing to Lemma 3.2 shows

$$|c_x^{(\lambda)}((\mathbf{0}, h'), (y', h'_y))| \leq 2C \exp\left(-\frac{1}{C} \max(h'_y, h')\right).$$

Combining the previous two displays concludes the proof of Lemma 3.5. \square

Given Lemmas 3.4 and 3.5, we now prove Theorem 1.2 as follows. By Palm theory for Poisson processes (see e.g. Theorem 1.6 of [25]) we express $\text{Var}[\langle f, \mu_{\lambda\rho}^\xi \rangle]$ as

$$\begin{aligned} \lambda \int_{A_+} f^2(\bar{x}) \mathbb{E}[\xi(\bar{x}, \mathcal{P}_{\lambda\rho})] \rho(\bar{x}) d\bar{x} + \\ \lambda^2 \int_{A_+} \int_{A_+} f(\bar{x}) f(\bar{y}) c_x^{(1)}((\mathbf{0}, h_x), (y-x, h_y)) \rho(\bar{x}) \rho(\bar{y}) d\bar{x} d\bar{y} \end{aligned} \quad (3.23)$$

where $\bar{x} := (x, h_x)$ and $\bar{y} := (y, h_y)$.

Following the proof of Theorem 1.1 verbatim shows that after normalization by λ^τ , the first integral converges as $\lambda \rightarrow \infty$ to

$$\int_A \int_0^\infty f^2(x, 0) \mathbb{E}[\xi^{(\infty)}((\mathbf{0}, h'_x), \mathcal{P}_{\rho_x^{(\infty)}})] \rho_0(x) (h'_x)^\delta dh'_x dx,$$

which by the definition of $m^{(\infty)}$ and the scaling relation (3.16) equals

$$\int_A \int_0^\infty f^2(x, 0) m^{(\infty)}(\mathbf{0}, h'_x) \rho_0^\tau(x) (h'_x)^\delta dh'_x dx.$$

Making again the usual substitutions $y' = \lambda^\beta(y-x)$, $h'_x = \lambda^\gamma h_x$, and $h'_y = \lambda^\gamma h_y$ and recalling $\rho(x, h_x) = \lambda^{-\gamma\delta} \rho^{(\lambda)}(\mathbf{0}, h'_x)$, the second integral in (3.23) becomes

$$\lambda^{2-2\gamma-2\gamma\delta-\beta(d-1)} \int_A \int_{\lambda^\beta A} \int_0^\infty \int_0^\infty f(x, h'_x \lambda^{-\gamma}) f(\lambda^{-\beta} y' + x, h'_y \lambda^{-\gamma}) c_x^{(\lambda)}((\mathbf{0}, h'_x), (y', h'_y)) \cdot \rho^{(\lambda)}(\mathbf{0}, h'_x) \rho^{(\lambda)}(y', h'_y) dh'_x dh'_y dy' dx.$$

Recalling from (1.12) that $\beta(d-1) + \gamma(1+\delta) = 1$, we have by definition of τ (see (1.5)) that $2 - 2\gamma - 2\gamma\delta - \beta(d-1) = 1 - \gamma(1+\delta) = \tau$. After normalization by λ^τ the above integral equals

$$\int_A \int_{\lambda^\beta A} \int_0^\infty \int_0^\infty f(x, h'_x \lambda^{-\gamma}) f(\lambda^{-\beta} y' + x, h'_y \lambda^{-\gamma}) g_\lambda(x, h'_x, y', h'_y) dh'_x dh'_y dy' dx$$

where we put

$$g_\lambda(x, h'_x, y', h'_y) := c_x^{(\lambda)}((\mathbf{0}, h'_x), (y', h'_y)) \rho^{(\lambda)}(\mathbf{0}, h'_x) \rho^{(\lambda)}(y', h'_y).$$

Clearly $f(x, h'_x \lambda^{-\gamma}) f(\lambda^{-\beta} y' + x, h'_y \lambda^{-\gamma})$ converges to $f^2(x, 0)$ as $\lambda \rightarrow \infty$. Lemma 3.4 implies for all $(x, h'_x, y', h'_y) \in A_+ \times \lambda^\beta A \times \mathbb{R}_+$ that the product $g_\lambda(x, h'_x, y', h'_y) (h'_x)^{-\delta} (h'_y)^{-\delta}$ converges to

$$c_x^{(\infty)}((\mathbf{0}, h'_x), (y', h'_y)) \rho_0^2(x)$$

as $\lambda \rightarrow \infty$. Since, by Lemma 3.5 and **(R2)**, $g_\lambda(x, h'_x, y', h'_y) (h'_x)^\delta (h'_y)^\delta$ is dominated in absolute value by the integrable function

$$(x, h'_x, y', h'_y) \mapsto C' (h'_x)^\delta (h'_y)^\delta \exp\left(-\frac{1}{C} \max\left(\frac{|y'|}{2}, h'_x, h'_y\right)\right)$$

on $A_+ \times \mathbb{R}^{d-1} \times \mathbb{R}_+$ the dominated convergence theorem combined with relation (3.16) produces the desired limit (1.10). \square

3.4 Proof of Theorem 1.3

Given Theorems 1.1 and 1.2, one may prove Theorem 1.3 either by the method of cumulants [7] or by the Stein method [29]. The first approach shows that the Fourier transform of $\lambda^{-\tau/2} \langle f, \bar{\mu}_{\lambda\rho}^\xi \rangle$ namely

$$\mathbb{E} \exp\left[i\lambda^{-\tau/2} \langle f, \bar{\mu}_{\lambda\rho}^\xi \rangle\right],$$

converges as $\lambda \rightarrow \infty$ to the Fourier transform of a normal mean zero random variable with variance $\sigma_f^2 := I(f^2) + J(f^2)$. Even though we use a formally different version of stabilization, this is accomplished by following [7] nearly verbatim. Indeed, recall that Lemma 3.5 shows the exponential decay of the two point correlation function $c_x^{(\lambda)}((\mathbf{0}, h'), (y', h'_y))$. In a similar way we may establish the exponential decay of k -point correlation functions, and, more generally, that the k -point

correlation functions cluster exponentially, as shown in Lemma 5.2 of [7]. In this way we show (as in Lemma 5.3 of [7]) that for all $k = 3, 4, \dots$ and $f \in \mathcal{C}_b(A_+)$ that

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\tau k/2} \langle f^{\otimes k}, c_\lambda^k \rangle = 0, \quad (3.24)$$

where c_λ^k denotes the k th cumulant figuring in the logarithm of the Laplace transform (their existence follows by Lemma 3.2). This consequently shows that $\lambda^{-\tau/2} \langle f, \bar{\mu}_{\lambda\rho}^\xi \rangle$ converges to a mean zero normal random variable with variance σ_f^2 . The convergence of the finite dimensional distributions follows from the Cramer-Wold device and is standard (see e.g. p. 251 of [7] or [26]).

Alternatively we may also use the Stein method [29, 26]. This is a bit simpler and has the advantage of yielding rates of convergence when $\sigma_f^2 > 0$, as would be the case when $\delta = 0$ and $\alpha = 2$ (Lemma 7 of [35] combined with Section 4 below) or when $\alpha = 1$ (Theorem 2.2 of [8]). (When $\sigma_f^2 = 0$ then $\lambda^{\tau/2} \langle f, \bar{\mu}_{\lambda\rho}^\xi \rangle$ converges to a unit point mass.) Our proof is based closely on [29], which uses a formally different version of stabilization. For simplicity we assume $A = [0, 1]^{d-1}$.

Recalling that $\bar{x} := (x, h_x)$ we have

$$\langle f, \mu_{\lambda\rho}^\xi \rangle = \sum_{\bar{x} \in \mathcal{P}_{\lambda\rho}} \xi(\bar{x}, \mathcal{P}_{\lambda\rho}) f(\bar{x}) = \sum_{\bar{x} \in \mathcal{P}_{\lambda\rho}} \xi^{(\lambda)}((\mathbf{0}, h'_x), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x]) f((x, h'_x \lambda^{-\gamma})).$$

For all $L > 0$ let

$$\begin{aligned} T_\lambda := T_\lambda(L) &:= \sum_{\bar{x} \in \mathcal{P}_{\lambda\rho} \cap ([0, 1]^{d-1} \times [0, L\lambda^{-\gamma} \log \lambda])} \xi^{(\lambda)}((\mathbf{0}, h'_x), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x]) f((x, h'_x \lambda^{-\gamma})) \\ &= \sum_{\bar{x} \in \mathcal{P}_{\lambda\rho}: h'_x \leq L\lambda^{-\gamma} \log \lambda} \xi^{(\lambda)}((\mathbf{0}, h'_x), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x]) f((x, h'_x \lambda^{-\gamma})). \end{aligned}$$

By Lemma 3.2, given arbitrarily large $\kappa > 0$, if L is large enough then $\langle f, \mu_{\lambda\rho}^\xi \rangle$ and T_λ coincide except on a set with probability $O(\lambda^{-\kappa})$ in λ . Thus T_λ has the same asymptotic distribution as $\langle f, \mu_{\lambda\rho}^\xi \rangle$ and it suffices to find a rate of convergence to the standard normal for $(T_\lambda - \mathbb{E} T_\lambda) / \sqrt{\text{Var} T_\lambda}$.

Subdivide $[0, 1]^{d-1}$ into $V(\lambda) := \lambda^{\beta(d-1)} (\rho_\lambda)^{-(d-1)}$ sub-cubes C_i^λ of edge length $\lambda^{-\beta} \rho_\lambda$ and of volume $\lambda^{-\beta(d-1)} (\rho_\lambda)^{d-1}$, where $\rho_\lambda := M \log \lambda$ for some large M , exactly as in section 4 of [29].

Enumerate $\mathcal{P}_{\lambda\rho} \cap (C_i^\lambda \times L\lambda^{-\gamma} \log \lambda)$ by $\{\bar{X}_{i,j}\}_{j=1}^{N_i}$ where $\bar{X}_{i,j} := (x_{ij}, h_{ij})$. Re-write T_λ as

$$T_\lambda = \sum_{i=1}^{V(\lambda)} \sum_{j=1}^{N_i} \xi^{(\lambda)}((\mathbf{0}, h'_{ij}), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x_{ij}]) f((x_{ij}, h'_{ij} \lambda^{-\gamma})).$$

This is the analog of T_λ in [29].

For any random variable X and any $p > 0$, let $\|X\|_p := (\mathbb{E}[|X|^p])^{1/p}$. For all $1 \leq i \leq V(\lambda)$ we have $\sum_{j=1}^{N_i} \xi^{(\lambda)}((\mathbf{0}, h'_{ij}), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x_{ij}]) \leq N_i$, where N_i is Poisson with mean

$$\lambda \int_{C_i^\lambda \times [0, L\lambda^{-\gamma} \log \lambda]} \rho(u) du = O([\log \lambda]^{1+\delta}).$$

It follows by the boundedness of f that

$$\left\| \sum_{j=1}^{N_i} \xi^{(\lambda)}((\mathbf{0}, h'_{ij}), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x_{ij}]) f((x_{ij}, h'_{ij}\lambda^{-\gamma})) \right\|_3 \leq C \|f\| L^{1+\delta} (\log \lambda)^{1+\delta} (\rho_\lambda)^{d-1},$$

where $\|f\|$ denotes the essential supremum of f . This is the analog of Lemma 4.3 in [29] (putting $q = 3$ there) with an extra logarithmic factor.

For all $1 \leq i \leq V(\lambda)$ and $j = 1, 2, \dots$ let $R_{i,j}$ denote the radius of stabilization for $\xi^{(\lambda)}$ at $X_{i,j}$ (for $\mathcal{P}_{\lambda\rho}^{(\lambda)}$) if $1 \leq j \leq N_i$ and let $R_{i,j}$ be zero otherwise.

As in [29], put $E_i := \bigcap_{j=1}^{\infty} \{R_{i,j} \leq \rho_\lambda\}$ and let $E_\lambda := \bigcap_{i=1}^{V(\lambda)} E_i$. Then by Lemma 3.1 (i) we have $P[E_\lambda^c] \leq \lambda^{-\kappa}$ for κ arbitrarily large if M is large enough. This is the analog of (4.11) of [29].

Next, recalling $\rho_\lambda = M \log \lambda$, we define the analog of T'_λ in [29]:

$$T'_\lambda := \sum_{i=1}^{V(\lambda)} \sum_{j=1}^{N_i} \xi_{[\rho_\lambda]}^{(\lambda)}((\mathbf{0}, h'_{ij}), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x_{ij}]) f((x_{ij}, h'_{ij}\lambda^{-\gamma})).$$

Then we define for all $1 \leq i \leq V(\lambda)$

$$S_i := S_{Q_i} := (\text{Var} T'_\lambda)^{-1/2} \sum_{j=1}^{N_i} \xi_{[\rho_\lambda]}^{(\lambda)}((\mathbf{0}, h'_{ij}), \mathcal{P}_{\lambda\rho}^{(\lambda)}[x_{ij}]) f((x_{ij}, h'_{ij}\lambda^{-\gamma})).$$

We define $S_\lambda := \sum_{i=1}^{V(\lambda)} (S_i - \mathbb{E} S_i)$, noting that it is the analog of S in [29].

Notice that T'_λ is a close approximation to T_λ and that by definition of E_i , $1 \leq i \leq V(\lambda)$, it has a high amount of independence between summands. In fact by the independence property of Poisson point processes, it follows that S_i and S_k are independent whenever $d(C_i^\lambda, C_k^\lambda) > 2\lambda^{-\beta} \rho_\lambda$.

Next we define a graph $G_\lambda := (\mathcal{V}_\lambda, \mathcal{E}_\lambda)$ as follows. The set \mathcal{V}_λ consists of the sub-cubes $C_1^\lambda, \dots, C_{V(\lambda)}^\lambda$ and the edges $(C_i^\lambda, C_j^\lambda)$ belong to \mathcal{E}_λ if $d(C_i^\lambda, C_j^\lambda) \leq 2\lambda^{-\beta} \rho_\lambda$. Since S_i and S_k are independent whenever $d(C_i^\lambda, C_k^\lambda) > 2\lambda^{-\beta} \rho_\lambda$ it follows that G_λ is a dependency graph for $\{S_i\}_{i=1}^{V(\lambda)}$.

Now proceed exactly as in [29], noting that:

- (i) $V(\lambda) = \lambda^{\beta(d-1)} (\rho_\lambda)^{-(d-1)}$,
- (ii) the maximum degree of G_λ is bounded by 5^d ,
- (iii) for all $1 \leq i \leq V(\lambda)$ we have $\|S_i\|_3 \leq K (\text{Var}(T'_\lambda))^{-1/2} (\log \lambda)^{1+\delta} (\rho_\lambda)^{d-1} =: \theta[\lambda]$.

These bounds correspond to the analogous bounds (i), (ii), and (iii) on pages 54-55 of [29]. Moreover, provided $\sigma_f^2 > 0$, then the counterpart of (v) of [29] holds, namely,

$$\text{Var}[T'_\lambda] = \Theta(\text{Var}[T_\lambda]) = \Theta(\lambda^\tau).$$

Putting $q = 3$ in (4.1) and (4.18) of [29] gives a rate of convergence for both S_λ and $(T_\lambda - \mathbb{E}T_\lambda)/\sqrt{\text{Var}T_\lambda}$ to the standard normal. This rate is

$$O(V(\lambda)\theta[\lambda]^3) = O(\lambda^{\beta(d-1)}(\rho_\lambda)^{-(d-1)}(\lambda^\tau)^{-3/2}(\log \lambda)^{3(1+\delta)}\rho_\lambda^{3(d-1)}).$$

Recalling that $\tau = \beta(d-1)$ we rewrite this as

$$O(\lambda^{-\tau/2} \log \lambda^{3(1+\delta)+2(d-1)}). \quad (3.25)$$

This completes the proof of Theorem 1.3. \square

4 Proofs of Applications

The purpose of the present section is to derive Theorems 2.1 and 2.2 from our general theorems of section one.

Proof of Theorem 2.1. To derive Theorem 2.1 from our general theory, we translate the convex hull problem into the language of ψ -growth processes with overlap. To this end, recall first that for a compact convex body $C \subseteq \mathbb{R}^d$ we define its support function $h_C : S_{d-1} \rightarrow \mathbb{R}$ by

$$h_C(u) := \sup_{\bar{x} \in C} \langle \bar{x}, u \rangle, \quad u \in S_{d-1},$$

with now $\langle \cdot, \cdot \rangle$ standing for the usual scalar product in \mathbb{R}^d , see Section 1.7 in [40]. An easily verified and yet crucial feature of the support functional $h(\cdot)$ is that

$$h_{\text{conv}\{\bar{x}_1, \dots, \bar{x}_k\}}(u) = \max_{1 \leq i \leq k} h_{\{\bar{x}_i\}}(u), \quad u \in S_{d-1}, \quad (4.1)$$

for each collection $\{\bar{x}_1, \dots, \bar{x}_k\}$ of points in \mathbb{R}^d . Moreover, by definition it is clear that for all $u \in S_{d-1}$ we have $h_{\{\bar{x}\}}(u) = \langle \bar{x}, u \rangle$, $u \in S_{d-1}$.

This leads to the following way of describing $\mathcal{V}(\mathcal{P}_{\lambda\rho})$ considered in Theorem 2.1. For a particular realization $\{\bar{x}_1, \dots, \bar{x}_k\}$ of $\mathcal{P}_{\lambda\rho}$ in B_d we consider the collection $H[\bar{x}_1], \dots, H[\bar{x}_k]$ of *support hypergraphs* given by

$$H[\bar{x}] := \{(y, h_y) \in S_{d-1} \times \mathbb{R}_+ : h_y \geq 1 - h_{\{\bar{x}\}}(y)\}, \quad (4.2)$$

where h_y stands for the distance between \bar{y} and the boundary $S_{d-1} = \partial B_d$. A compact convex body is uniquely determined by its support functional (cf. Section 1.7 in [40]), and in view of (4.1), the set $\text{conv}(\{\bar{x}_1, \dots, \bar{x}_k\})$ is in one-to-one correspondence with the union $\bigcup_{i=1}^k H[\bar{x}_i]$. Further, the number of vertices in the convex hull is easily seen to coincide with the number of those \bar{x}_i , $i = 1, \dots, k$, for which $H[\bar{x}_i]$ is not completely contained in the union $\bigcup_{j \neq i} H[\bar{x}_j]$.

Next we shall also write $r_y := 1 - h_y$ for the distance between \bar{y} and the origin of \mathbb{R}^d . Note now that the intensity measure $\rho(\bar{x})d\bar{x}$, $\bar{x} \in B_d$, coincides with $\rho((x, r))r^{d-1}drdx = \rho((x, r))(1 - h)^{(d-1)}dhdx$ where $\bar{x} := (x, r)$, with $r \in [0, 1]$ denoting the distance between \bar{x} and the origin of \mathbb{R}^d , with $h := 1 - r$ and with $x \in S_{d-1}$ being the radial projection of \bar{x} onto $\partial B_d = S_{d-1}$. Observe also that the support hypergraph $H[\bar{x}]$ as given in (4.2) can be represented by

$$H[(x, r)] = \{(y, h_y) \in S_{d-1} \times \mathbb{R}_+ : h_y \geq 1 - r \cos(\text{dist}_{S_{d-1}}(x, y))\}$$

with $\text{dist}_{S_{d-1}}(x, y) := \cos^{-1}\langle x, y \rangle$ denoting the geodesic distance in S_{d-1} between x and y . Now put

$$\psi(l) := 1 - \cos(l).$$

Writing the inequality $h_y \geq 1 - r \cos(\text{dist}_{S_{d-1}}(x, y))$ as $h_y \geq 1 - r + r\psi(\text{dist}_{S_{d-1}}(x, y))$ we have

$$H[(x, r)] = \{(y, h_y) \in S_{d-1} \times \mathbb{R}_+ : h_y \geq h + r\psi(\text{dist}_{S_{d-1}}(x, y))\}, \quad (4.3)$$

in other words the support hypergraphs are remarkably similar to the upwards cones (1.1) described at the outset.

The above observations naturally suggest *identifying the cardinality of the studied set* $\mathcal{V}(\mathcal{P}_{\lambda\rho})$ *with the number of extreme points in the $r\psi$ -growth process with overlap in the sense of section one with the underlying point density* $\rho((x, r))r^{d-1} = \rho((x, r))(1 - h)^{d-1}$. Likewise the vertex empirical measure $\mu_{\lambda\rho}$ in (2.1) corresponds to the empirical measure $\mu_{\lambda\rho}^\xi$, $\xi := \xi(\psi; \cdot)$, see (1.2).

This identification is valid modulo the following issues though:

(1) the ‘spatial’ coordinate x of a point $\bar{x} := (x, r) \in B_d$ falls into S_{d-1} rather than into a subset A of \mathbb{R}^{d-1} , as required in section one,

(2) ψ as given above is monotone only in a neighborhood of 0, and moreover we do not have $\lim_{l \rightarrow \infty} \psi(l) = \infty$, which violates **(Ψ1)**,

(3) the support hypergraph $H[(x, r)]$ coincides with the (x, h) -shifted ψ -hypergraph $K[x, h]$ given by (1.1) only when $r = 1$ and hence only when $h = 0$; in general for $0 \leq r \leq 1$ the set $H[(x, r)]$ is an (x, h) -shifted $r\psi$ -hypergraph.

We claim, however, that the above three restrictions can be neglected in the asymptotic regime $\lambda \rightarrow \infty$ thus rendering the theory of section one applicable. Indeed, first note that the sphere S_{d-1} , unlike the boundary of a general smooth convex body, has a spatially homogeneous structure and so the behavior of ψ is independent of x , exactly as in section one. Moreover, the sphere S_{d-1} , being a smooth manifold, has a local geometry coinciding with that of \mathbb{R}^{d-1} , which takes care of issue (1). Concerning issues (2) and (3), for each $r \in (0, 1)$ the convex hull $\text{conv}(\mathcal{P}_{\lambda\rho})$ coincides with $\text{conv}(\mathcal{P}_{\lambda\rho} \cap (B_d \setminus B_d(0, r)))$ with overwhelming probability, i.e., the probability of the complement event goes to zero exponentially fast in λ , see the discussion in [22] and the references therein. This allows us to focus on the geometry of $\text{conv}(\mathcal{P}_{\lambda\rho})$ in a thin shell $B_d \setminus B_d(0, r)$ within a distance $1 - r$ from the boundary S_{d-1} .

Consequently, only the behavior of ψ in a neighborhood of 0 matters. Recalling that the standard re-scaling of subsection 1.3 involves scaling in the spatial directions by λ^β , it follows that for a given $\bar{x} := (x, r)$ and support hypergraph $H[\bar{x}]$, the contribution of points distant from x by more than $O(\lambda^{-\beta})$ is negligible in view of the argument in Lemma 3.1(i) and no distortions from the local Euclidean geometry have to be taken into account in the limit under this re-scaling. Likewise, we only have to control the geometry of $H[\bar{x}]$, $\bar{x} := (x, r)$, for r arbitrarily close to 1. This allows us to rewrite the proofs of Theorems 1.1 - 1.3 for the thus modified r -dependent ψ . Indeed, the stabilization Lemma 3.1 as well as Lemma 3.2 do not require any modifications in their proof and nor does Lemma 3.4 or Lemma 3.5. Consequently, the arguments leading to the central limit theorem in subsection 3.4 do not require modification either. In this context we note that the proof of Lemma 3.1 would break down if the sphere S_{d-1} were replaced by a non-convex set allowing for long-range dependencies between extreme points.

It only remains to show the limit arguments in subsections 3.3 and 3.2 remain valid for the modified ψ . To see that this is indeed the case we note that the arguments rely on two main ingredients: on stabilization which holds with no changes as stated above, and on re-scaling relations discussed in subsection 1.3. However, it is easily seen that the re-scaling relations and their proofs can be readily rewritten for the modified ψ , the only essential modification being to add one extra argument ($h := 1 - r$) to the ψ -function, which anyway vanishes in the scaling limit of subsection 1.3 with $h = 1 - r$ tending to 0 as discussed above (whereas the contribution coming from smaller h is negligible in view of Lemma 3.2). This discussion takes care of issues (2) and (3) above.

Thus we can now conclude that the considered convex hull process falls into the range of applicability of the general theory of section one, with $\alpha = 2$ in **(Ψ2)** and δ in **(R2)** coinciding

with that in the statement of Theorem 2.1. Thus, we obtain the required Theorem 2.1 as a consequence of the general Theorems 1.1 - 1.3. The rate of convergence follows from (3.25) by putting $\delta = 0$ and $\alpha = 2$. \square

Proofs of Theorems 2.2 and 2.3. Theorem 2.2 follows directly by the general theory in section one (Theorems 1.1 - 1.3 with $\alpha \in (0, 1]$). The rate (2.8) follows from (3.25) by putting $\delta = 0$ and $\alpha = 1$. We thus focus attention on establishing Theorem 2.3. The first lemma yields (2.9).

Lemma 4.1 *For all $f \in \mathcal{C}_b(A_+)$ we have*

$$|\mathbb{E}[\langle f, \nu_n^\xi \rangle] - \mathbb{E}[\langle f, \mu_{n\rho}^\xi \rangle]| = O(n^{-\tau'}). \quad (4.4)$$

Proof. For all $\bar{w} \in A_+$, let $p(\bar{w}) := \int_{K^\downarrow[\bar{w}]} \rho(u) du$, where $K^\downarrow[\bar{w}]$ is as in (2.5) with $\psi(l) = l^\alpha$. Note that in our current setting for all $w \in A_+$ we have $p(w) \in [0, 1]$ since ρ is a probability density. Also, note that $\psi^{(n)} \equiv \psi$ and $K^{(n)} \equiv K$ with $K^{(n)} := \{(y^{(n)}, h_y^{(n)}) : (y, h_y) \in K\}$, i.e., the self similarity under the re-scaling is immediate rather than emerging as $n \rightarrow \infty$. For all $s \in [0, 1]$ and $f \in \mathcal{C}_b(A_+)$ let $B_f(s) := \int_{p(\bar{w}) \leq s} f(\bar{w}) \rho(\bar{w}) d\bar{w}$. Recalling that for $\alpha \in (0, 1]$ the ψ -extremality of a point w in a given sample is equivalent to having no other sample points in $K^\downarrow[w]$ (see the discussion at the beginning of subsection 2.2), we have

$$\begin{aligned} \mathbb{E}[\langle f, \nu_n^\xi \rangle] &= n \int_{A_+} (1 - p(w))^{n-1} f(w) \rho(w) dw \\ &= n \int_0^1 \int_{p(w)=s} (1 - s)^{n-1} f(w) \rho(w) dw ds = n \int_0^1 (1 - s)^{n-1} dB_f(s) \end{aligned}$$

by Fubini's theorem. Similarly

$$\mathbb{E}[\langle f, \mu_{n\rho}^\xi \rangle] = n \int_0^1 e^{-ns} dB_f(s) \sim C_f n^\tau, \quad (4.5)$$

where the asymptotics are given by Theorem 1.1. B_f is monotone, non-decreasing and Karamata's Tauberian theorem (e.g. Theorem 2.3 in [41]) gives $B_f(s) \sim C_f s^{\tau'}$ as $s \rightarrow 0^+$. Notice

$$\begin{aligned} |\mathbb{E}[\langle f, \mu_{n\rho}^\xi \rangle] - \mathbb{E}[\langle f, \nu_n^\xi \rangle]| &= n \int_0^1 (e^{-ns} - (1 - s)^{n-1}) dB_f(s) \\ &\leq n \int_0^1 (e^{-ns} - e^{n \ln(1-s)}) dB_f(s) \\ &\leq Cn^2 \int_0^1 e^{-ns} s^2 dB_f(s) \end{aligned}$$

$$= Cn^2 \int_0^{1/n} e^{-ns} s^2 dB_f(s) + Cn^2 \int_{1/n}^1 e^{-ns} s^2 dB_f(s).$$

The first integral behaves like $Cn^{-\tau'}$ since $B_f(s) \sim C_f s^{\tau'}$ whereas the second behaves like $\frac{C}{n} \int_1^n u^2 e^{-u} dB_f(u/n) \leq C/n$, since B_f is bounded by $B_f(1)$. This gives (4.4). \square

We now establish the remainder of Theorem 2.3. Recall $\bar{u}' := (u', h'_u)$. For all $\lambda > 0$ define

$$A'(\lambda) := \left\{ \bar{y}' \in \lambda^\beta A \times \mathbb{R}_+ : \int_{K^1[\bar{y}']} \rho^{(\lambda)}(u) du \leq C \log \lambda \right\}.$$

Let $A(\lambda) := \{\bar{y} \in A_+ : \bar{y}' \in A'(\lambda)\}$ and put $a_\lambda := \int_{A(\lambda)} \rho(w) dw$. Note that by Lemma 3.2 the probability that a sample point from $\bar{\mathcal{X}}_n := \{X_i\}_{i=1}^n$ in $A_+ \setminus A(\lambda)$ is ψ -extremal is at most

$$C \exp\left(-\frac{n \log \lambda}{C\lambda}\right) \quad (4.6)$$

and the same holds for $\bar{\mathcal{X}}_n$ replaced by Poisson sample with intensity $n\rho$. Indeed, although Lemma 3.2 was originally established for Poisson samples, it is easily seen that the same proof works also for binomial samples, as it essentially relies on exponentially decaying upper bounds for probabilities of certain sets in A_+ being devoid of points of the underlying point process. Thus the ψ -extremal points are predominantly concentrated in $A(\lambda)$, a fact which we will use to show (2.10). First we find growth bounds for a_λ .

Lemma 4.2 *We have $a_\lambda \leq C(\log \lambda)^{\frac{\alpha(1+\delta)}{\alpha+d-1}} \lambda^{\frac{-\alpha(1+\delta)}{d-1+\alpha(1+\delta)}}$.*

Proof. If $M(\lambda) := \sup\{h_y : h_y \in A(\lambda)\}$ then note that $\alpha(\lambda)$ grows like $\int_0^{M(\lambda)} h_y^\delta dh_y = C(M(\lambda))^{1+\delta}$. We now find $M(\lambda)$.

If $\bar{y}' := (y', h'_y) \in A'(\lambda)$ then by (3.7) we have $h'_y \frac{\alpha+d-1}{\alpha} \leq C \log \lambda$. Since $h'_y = \lambda^\gamma h_y$ and since $\gamma = \beta\alpha$, it follows that

$$h_y \frac{\alpha+d-1}{\alpha} \lambda^{\beta(\alpha+d-1)} \leq C \log \lambda.$$

Since $\gamma(d-1)/\alpha = \tau$ we have $h_y \frac{\alpha+d-1}{\alpha} \lambda^{\gamma+\tau} \leq C \log \lambda$ or $h_y \leq (\log \lambda)^{\frac{\alpha}{\alpha+d-1}} \lambda^{\frac{-\alpha(\gamma+\tau)}{\alpha+d-1}}$, that is

$$M(\lambda) \leq (\log \lambda)^{\frac{\alpha}{\alpha+d-1}} \lambda^{\frac{-\alpha(\gamma+\tau)}{\alpha+d-1}}.$$

Recall that $\gamma + \tau = (\alpha + d - 1)/(d - 1 + \alpha(1 + \delta))$ to get the result. \square

The next lemma yields (2.10). The proof borrows heavily from [8] and for the sake of completeness we provide the details.

Lemma 4.3 For all $f \in \mathcal{C}_b(A_+)$ we have $\lim_{n \rightarrow \infty} n^{-\tau} \text{Var}[\langle f, \nu_n^\xi \rangle] = \lim_{n \rightarrow \infty} n^{-\tau} \text{Var}[\langle f, \mu_{n\rho}^\xi \rangle]$.

Proof. Recall $\bar{\mathcal{X}}_n := \{X_i\}_{i=1}^n$. Let $N_n := \text{card}\{\bar{\mathcal{X}}_n \cap A(n)\}$ and $N'_n := \text{card}\{\mathcal{P}_{n\rho} \cap A(n)\}$. For all $r = 1, 2, \dots$ denote by $e(r) := e_f(r)$ the expected value of the functional $\langle f \cdot \mathbf{1}(A(n)), \mu_n^\xi \rangle$ conditioned on $\{N(n) = r\}$, and by $v(r)$ the variance of this functional conditioned on $\{N(n) = r\}$. Let $\nu_n^A := \nu_n^{\xi, A}$ denote the point measure induced by the ψ -extremal points in $\{\bar{\mathcal{X}}_n \cap A(n)\}$. Similarly, let $\mu_{n\rho}^A := \mu_{n\rho}^{\xi, A}$ denote the point measure induced by the ψ -extremal points in $\{\mathcal{P}_{n\rho} \cap A(n)\}$.

By the bound (4.6) on the probability of a given point outside $A(n)$ being extremal, ν_n^A coincides with ν_n and $\mu_{n\rho}^{\xi, A}$ coincides with $\mu_{n\rho}^\xi$ except on a set with probability at most $nC \exp(-C \log n) = Cn^{-C+1}$. Since C can be chosen arbitrarily large it suffices to show that

$$\lim_{n \rightarrow \infty} n^{-\tau} \text{Var}[\langle f, \nu_n^A \rangle] = \lim_{n \rightarrow \infty} n^{-\tau} \text{Var}[\langle f, \mu_{n\rho}^A \rangle]. \quad (4.7)$$

The conditional variance formula implies that

$$\text{Var}[\langle f, \nu_n^A \rangle] = \text{Var}[e(N_n)] + \mathbb{E}[v(N_n)] \quad \text{and} \quad \text{Var}[\langle f, \mu_{n\rho}^A \rangle] = \text{Var}[e(N'_n)] + \mathbb{E}[v(N'_n)].$$

We prove (4.7) by showing that:

(i) the terms $\mathbb{E}[v(N_n)]$ and $\mathbb{E}[v(N'_n)]$ are dominant and that their ratio tends to one as $n \rightarrow \infty$, and

(ii) $\text{Var}[e(N_n)]$ and $\text{Var}[e(N'_n)]$ are both $o(n^\tau)$.

We will first show (ii) as follows. For all $s > 0$, recall that $B_f(s) := \int_{p(\bar{w}) \leq s} f(\bar{w}) \rho(\bar{w}) d\bar{w}$. By Fubini's theorem, for all $r = 1, 2, \dots$ and with $a_n = \int_{A(n)} \rho(w) dw$, we obtain

$$e(r) = \frac{r}{a_n} \int_{A(n)} \left(1 - \frac{p(w)}{a_n}\right)^{r-1} f(w) \rho(w) dw = \frac{r}{a_n} \int_0^{a_n} \left(1 - \frac{s}{a_n}\right)^{r-1} dB_f(s).$$

Letting Δ_r denote the difference $e(r+1) - e(r)$, we obtain

$$\Delta_r = \frac{1}{a_n} \int_0^{a_n} \left(1 - \frac{s}{a_n}\right)^r - \frac{rs}{a_n} \left(1 - \frac{s}{a_n}\right)^{r-1} dB_f(s).$$

Setting $u = rs/a_n$ and applying $B_f(s) \sim C_f s^{\tau'}$, we see that ($\tau = 1 - \tau'$):

$$|\Delta_r| \leq \frac{C_f}{r} \int_0^r \left| \left(1 - \frac{u}{r}\right)^r - u \left(1 - \frac{u}{r}\right)^{r-1} \right| \left(\frac{ua_n}{r}\right)^{-\tau} du.$$

Since $\sup_{r>0} \int_0^r \left| \left(1 - \frac{u}{r}\right)^r - u \left(1 - \frac{u}{r}\right)^{r-1} \right| u^{-\tau} du \leq C$ it follows that $|\Delta_r| \leq C \left(\frac{a_n}{r}\right)^{-\tau}$.

When $r \in I_n := (na_n - C(\log n)(na_n)^{1/2}, na_n + C(\log n)(na_n)^{1/2})$, then, by Lemma 4.2, for n large

$$|\Delta_r| \leq C(na_n)^{-1} n^\tau = Ca_n^{-1} n^{-\tau'} = Cn^{\frac{\alpha(1+\delta)}{d-1+\alpha(1+\delta)}} n^{-\tau'} (\log n)^{\frac{-\alpha(1+\delta)}{\alpha+d-1}}.$$

Recalling that $\tau' = (1 + \delta)\alpha/(d - 1 + \alpha(1 + \delta))$ we see that for $r \in I_n$ we have

$$|\Delta_r| \leq \Delta(n) := C(\log n)^{\frac{-\alpha(1+\delta)}{\alpha+d-1}}.$$

Write $e(N_n) = e(1) + \sum_{j=2}^{N_n} (e(j) - e(j-1))$ and observe that $e(N_n)$ differs from the constant $e(1) + \sum_{j=2}^{\mathbb{E}[N_n]} (e(j) - e(j-1))$ by at most

$$\sum_{j \in J_n} (e(j) - e(j-1))$$

where $J_n := (\min(\mathbb{E}[N_n], N_n), \max(\mathbb{E}[N_n], N_n))$. Thus

$$\text{Var}[e(N_n)] \leq \mathbb{E} \left[\sum_{j \in J_n} (e(j) - e(j-1)) \right]^2 \leq \mathbb{E} \left[\sum_{j \in J_n} (e(j) - e(j-1)) \mathbf{1}_{N_n \in I_n} \right]^2 + o(1),$$

by Cauchy-Schwarz and since (by increasing C in the definition of I_n) standard concentration inequalities (see e.g. Proposition A.2.3(ii,iii) and Proposition A.2.5(ii,iii) in [3]) show that $P[N_n \in I_n^c]$ can be made smaller than any negative power of n .

For $j \in J_n$ and $N_n \in I_n$ we have $j \in I_n$ and so $(e(j) - e(j-1)) \leq \Delta(n)$. Since the length of J_n is bounded by $|N_n - \mathbb{E} N_n|$, it follows that $\text{Var}[e(N_n)] \leq \text{Var}[N_n](\Delta(n))^2 + o(1)$. Note that $\text{Var}[N_n] \leq Cn^\tau (\log n)^{\alpha(1+\delta)/(\alpha+d-1)}$. It follows that $\text{Var}[e(N_n)] \leq Cn^\tau (\log n)^{-\alpha(1+\delta)/(\alpha+d-1)} + o(1)$, that is $\text{Var}[e(N_n)] = o(n^\tau)$. Similarly $\text{Var}[e(N'_n)] = o(n^\tau)$ and so condition (ii) holds.

We now show condition (i) by showing that the ratio $\mathbb{E}[v(N_n)]/\mathbb{E}[v(N'_n)]$ is asymptotically one, as $n \rightarrow \infty$. Let $p_{n,r} := P[N_n = r]$ and $p'_{n,r} := P[N'_n = r]$. Stirling's formula implies that for $|r - a_n n| \leq n^\beta$, where $0 < \beta < 1/2$,

$$\lim_{n \rightarrow \infty} \frac{p_{n,r}}{p'_{n,r}} = 1 \tag{4.8}$$

uniformly. Now, for $|r - a_n n| > n^\beta$, where $\beta \in (0, 1/2)$ is chosen so that $n^{2\beta}/na_n$ grows faster than some (small) power of n , we have that both $p_{n,r}$ and $p'_{n,r}$ are bounded by $C \exp(-n^\delta/C)$ for some C , $\delta > 0$ (see e.g. Proposition A.2.3 (i) and Proposition A.2.5. (i) in [3]). Write

$$\mathbb{E}[v(N_n)] = \sum_{|r - a_n n| \leq n^\beta} v(r) p_{n,r} + \sum_{|r - a_n n| > n^\beta} v(r) p_{n,r}.$$

The second sum is negligible since $0 < v(r) < r^2$ and $p_{n,r}$ is exponentially small. Consider the terms in the first sum. By (4.8), we have $p_{n,r} = p'_{n,r}(1 + o(1))$ uniformly for all $|r - a_n n| \leq n^\beta$ and since the terms in the first sum are positive it follows that

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}[v(N_n)]}{\mathbb{E}[v(N'_n)]} = 1. \tag{4.9}$$

Now from before we know $\text{Var}[\langle f, \mu_{n\rho}^A \rangle]$ has asymptotic growth $Cn^\tau, C > 0$. It follows that $\mathbb{E}[v(N'_n)]$ has the same growth, since $\text{Var}[e(N'_n)] = o(n^\tau)$. Thus by (4.9) and the growth bounds $\text{Var}[e(N_n)] = o(n^\tau)$ and $\text{Var}[e(N'_n)] = o(n^\tau)$, the desired identity (4.7) follows, completing the proof of Lemma 4.3. \square

We conclude the proof of Theorem 2.3 by showing for all $f \in \mathcal{C}_b(A_+)$

$$\lim_{n \rightarrow \infty} d_{TV}(n^{-\tau/2} \langle f, \bar{\nu}_n^\xi \rangle, n^{-\tau/2} \langle f, \bar{\mu}_{n\rho}^\xi \rangle) = 0,$$

where the total variation distance between two measures m_1 and m_2 is $d_{TV}(m_1, m_2) := \sup_B |m_1(B) - m_2(B)|$, where the sup runs over all Borel subsets in \mathbb{R}^d . Since $n^{-\tau/2} |\mathbb{E}[\langle f, \nu_n^\xi \rangle] - \mathbb{E}[\langle f, \mu_{n\rho}^\xi \rangle]| \rightarrow 0$ by (4.4) and since $n^{-\tau/2} \langle f, \bar{\mu}_{n\rho}^\xi \rangle$ converges in law to an appropriate Gaussian distribution, recalling that $a_n = o(1)$ (see Lemma 4.2), Theorem 2.3 follows at once from

Lemma 4.4 *For all $f \in \mathcal{C}_b(A_+)$ we have*

$$d_{TV}(\langle f, \nu_n^\xi \rangle, \langle f, \mu_{n\rho}^\xi \rangle) = O(a_n). \quad (4.10)$$

Proof. We follow the proof of Lemma 7.1 in [8]. Recall that ν_n^A is the measure induced by the maximal points in $\{(X_i, h_i)\}_{i=1}^n \cap A(n)$ and similarly let $\mu_{n\rho}^{\xi, A}$ be the measure induced by the maximal points in $\mathcal{P}_{n\rho} \cap A(n)$. If C is large enough in the definition of $A(n)$, then the probability that points in $A_+ \setminus A(n)$ contribute to ν_n^ξ or $\mu_{n\rho}^\xi$ is $O(n^{-2})$. It follows that for all $f \in \mathcal{C}_b(A_+)$

$$d_{TV}(\langle f, \mu_{n\rho}^\xi \rangle, \langle f, \mu_{n\rho}^{\xi, A} \rangle) = O(n^{-2}) = o(a_n) \text{ and } d_{TV}(\langle f, \nu_n^\xi \rangle, \langle f, \mu_n^{\xi, A} \rangle) = O(n^{-2}) = o(a_n).$$

Thus we only need to show $d_{TV}(\langle f, \nu_n^A \rangle, \langle f, \mu_n^{\xi, A} \rangle) = O(a_n)$.

Recall that N_n is the number of points from $\bar{\mathcal{X}}_n$ belonging to $A(n)$. Conditional on $N = r$, $\langle f, \nu_n^A \rangle$ is distributed as $\langle f, \tilde{\nu}_r^A \rangle$ where $\tilde{\nu}_r^A$ is the point measure induced by considering the maximal points among r points placed randomly according to the restriction of ρ to $A(n)$. The same is true for $\langle f, \mu_{n\rho}^{\xi, A} \rangle$ conditional on the cardinality of $\{\mathcal{P}_{n\rho} \cap A(n)\}$ taking the value r .

Hence, with $Bi(n, p)$ standing for a binomial random variable with parameters n and p and $Po(\alpha)$ standing for a Poisson random variable with parameter α we have for all $f \in \mathcal{C}_b(A_+)$

$$d_{TV}(\langle f, \nu_n^A \rangle, \langle f, \mu_{n\rho}^{\xi, A} \rangle) \leq C d_{TV}(Bi(n, a_n), Po(na_n)) \leq C \frac{1}{na_n} \sum_{i=1}^n (a_n)^2 \leq Ca_n,$$

where the penultimate inequality follows by standard Poisson approximation bounds (see e.g. (1.23) of Barbour, Holst, and Janson [3]). This is the desired estimate (4.10). \square

Acknowledgements. The authors gratefully acknowledge helpful discussions with Yuliy Baryshnikov, who in particular encouraged developing theory under the general condition **(R2)** and who also contributed to the proofs of Lemmas 4.1 and 4.3. The authors also thank an anonymous referee for comments leading to an improved exposition.

References

- [1] Z.-D. Bai, C.-C. Chao, H.-K. Hwang, and W.-Q. Liang (1998), On the variance of the number of maxima in random vectors and applications. *Ann. Appl. Prob.*, **8**, 886-895.
- [2] Z.-D. Bai, H.-K. Hwang, W.-Q. Liang, and T.-H. Tsai (2001), Limit theorems for the number of maxima in random samples from planar regions. *Electronic Journal of Prob.* **6**, no. 3, 1 -41.
- [3] A. D. Barbour, L. Holst, and S. Janson (1992), Poisson Approximation. Oxford Univ. Press.
- [4] A. D. Barbour and A. Xia (2001), The number of two dimensional maxima. *Advances Appl. Prob.* **33**, 727-750.
- [5] O. Barndorff-Nielsen and M. Sobel (1966), On the distribution of the number of admissible points in a vector. *Theory of Probability and its Applications* **11**, 249-269.
- [6] Yu. Baryshnikov (2000), Supporting-points processes and some of their applications. *Prob. Theory and Related Fields* **117**, 163-182.
- [7] Yu. Baryshnikov and J. E. Yukich (2005), Gaussian limits for random measures in geometric probability. *Annals Appl. Prob.*, **15**, 1A, 213-253.
- [8] Yu. Baryshnikov and J. E. Yukich (2005), Gaussian limits and maximal points. Preprint, electronically available via <http://www.lehigh.edu/~jey0/publications.html>.
- [9] P. Billingsley (1968), *Convergence of Probability Measures*, John Wiley & Sons.
- [10] A. J. Cabo and P. Groeneboom (1994), Limit theorems for functionals of convex hulls. *Probab. Theory Related Fields*, **100**, 31-55.
- [11] W.-M. Chen, H.-K. Hwang and T.-H. Tsai (2003), Efficient maxima finding algorithms for random planar samples. *Discrete Mathematics and Theor. Comp. Sci.*, **6**, 107-122.

- [12] S. N. Chiu (1997), A central limit theorem for linear Kolmogorov's birth-growth models. *Stochastic Proc. and Applic.*, **66**, 97-106.
- [13] S. N. Chiu and M. P. Quine (1997), Central limit theory for the number of seeds in a growth model in \mathbb{R}^d with inhomogeneous Poisson arrivals. *Annals of Appl. Prob.*, **7**, 802-814.
- [14] S. N. Chiu and M. P. Quine (2001), Central limit theorem for germination-growth models in \mathbb{R}^d with non-Poisson locations. *Advances Appl. Prob.*, **33**, no. 4.
- [15] S. N. Chiu and H. Y. Lee (2002), A regularity condition and strong limit theorems for linear birth growth processes. *Math. Nachr.*, 1 - 7.
- [16] D. J. Daley and D. Vere-Jones (1988), An Introduction to the Theory of Point Processes. Springer-Verlag.
- [17] L. Devroye (1993), Records, the maximal layer, and uniform distributions in monotone sets. *Computers Math. Applic.* **25**, 5, 19-31.
- [18] M. Ehrgott (2000), Multicriteria Optimization. Springer, Berlin.
- [19] P. Groeneboom (1988), Limit theorems for convex hulls. *Probab. Theory Related Fields*, **79**, 327-368.
- [20] L. Holst, M. P. Quine and J. Robinson (1996), A general stochastic model for nucleation and linear growth. *Annals Appl. Prob.*, **6**, 903-921.
- [21] I. Hueter (1999), Limit theorems for the convex hull of random points in higher dimensions. *Trans. Amer. Math. Soc.*, **351**, 4337-4363.
- [22] K. H. Küfer (1994), On the approximation of a ball by random polytopes, *Adv. Appl. Probab.* **26**, 876-892.
- [23] A. N. Kolmogorov (1937), On statistical theory of metal crystallisation. *Izvestia Academy of Science*, USSR, ser. Math. **3**, 355-360 (in Russian).
- [24] J. Møller (1992), Random Johnson-Mehl tessellations. *Advances Appl. Prob.* **24**, 814-844.
- [25] M. D. Penrose (2003), Random Geometric Graphs, Clarendon Press, Oxford.
- [26] M. D. Penrose (2007), Gaussian limits for random geometric measures, Electronic Journal of Probability, to appear.

- [27] M. D. Penrose and J. E. Yukich (2002), Limit theory for random sequential packing and deposition. *Ann. Appl. Probab.* **12**, 272-301.
- [28] M. D. Penrose and J. E. Yukich (2004), Weak laws of large numbers in geometric probability. *Ann. Appl. Probab.* **13**, 277-303.
- [29] M. D. Penrose and J. E. Yukich (2005), Normal approximation in geometric probability. In Stein's Method and Applications, Lecture Note Series, Institute for Mathematical Sciences, National University of Singapore, **5**, A. D. Barbour and Louis H. Y. Chen, Eds., 37-58, Electronically available via <http://www.lehigh.edu/~jey0/publications.html>.
- [30] J.-C. Pomerol and S. Barba-Romero (2000), Multicriterion Decision in Management. Kluwer Academic Publishers, Boston.
- [31] Reiss, R.-D. (1993), A Course on Point Processes. Springer, Berlin.
- [32] M. Reitzner (2002), Random points on the boundary of smooth convex bodies. *Trans. Amer. Math. Soc.* **354**, 2243-2278.
- [33] M. Reitzner (2003), Random polytopes and the Efron-Stein jackknife inequality. *Ann. Probab.* **32**, 2136-2166.
- [34] M. Reitzner (2005), The combinatorial structure of random polytopes. *Adv. Math* **191**, 178-208.
- [35] M. Reitzner (2005), Central limit theorems for random polytopes. *Prob. Th. Related Fields* **133**, 483-507.
- [36] A. Rényi (1962), Théorie des éléments saillants d'une suite d'observations. *Colloq. on Combinatorial Methods in Prob. Theory*, 104-115, Math. Institut, Aarhus Universitet, Denmark.
- [37] A. Rényi and R. Sulanke (1963), Über die konvexe Hülle von n zufällig gewählten Punkten II. *Z. Wahrscheinlichkeitstheorie und verw. Gebiete*, **2**, 75-84.
- [38] S. I. Resnick (1987), Extreme Values, Regular Variation and Point Processes. Springer-Verlag.
- [39] R. Schneider (1988), Random approximation of convex sets. *J. Microscopy* **151**, 211-227.
- [40] R. Schneider (1993), Convex bodies: the Brunn-Minkowski theory. (Encyclopaedia Math. Appl. **44**), Cambridge University Press.

- [41] E. Seneta (1976), Regularly Varying Functions. Lect. Notes in Mathematics, **508**, Springer Verlag.
- [42] D. Stoyan, W. Kendall, and J. Mecke (1995), Stochastic Geometry and Its Applications. John Wiley and Sons, Second Ed.
- [43] T. Schreiber (2003), Asymptotic geometry of high density smooth-grained Boolean models in bounded domains. *Adv. Appl. Prob.* **35**, 913-936.
- [44] Van Vu (2005), Central limit theorems for random polytopes in a smooth convex set. Preprint.

Tomasz Schreiber, Faculty of Mathematics and Computer Science, Nicholas Copernicus University, Toruń, Poland: tomeks@mat.uni.torun.pl

J.E. Yukich, Department of Mathematics, Lehigh University, Bethlehem PA 18015:
joseph.yukich@lehigh.edu