

Advanced Operations Research Techniques

IE316

Lecture 9

Dr. Ted Ralphs

Reading for This Lecture

- Bertsimas 3.5-3.9

Finding an Initial Basic Feasible Solution

- Recall that we need an *initial BFS* to start the simplex methods.
- Ideally, the initial basis matrix would also be the identity matrix (easy to invert).
- In some cases, this is easy to achieve.
 - If the formulation contains only inequalities and the origin is feasible, then there is no problem.
 - The initial BFS is simply $x = 0, s = b$ and the initial basis matrix is the identity matrix (easy to invert).
- For problems with equalities or where the origin is not feasible, things are more difficult.
- To deal with this, we need the concept of an *artificial variable*.
- Artificial variables are used in the equality rows like slack variables, but then forced to zero.

Obtaining a Feasible Solution

- Suppose we are given an LP $\min\{c^T x \mid Ax = b, x \geq 0\}$ already in standard form.
- Assume without loss of generality that $b \geq 0$.
- Then we can obtain an initial BFS by solving the following auxiliary LP.

$$\begin{aligned} \min \quad & \sum_{i=1}^m y_i \\ \text{s.t.} \quad & Ax + y = b \\ & x \geq 0 \\ & y \geq 0 \end{aligned}$$

- If the optimal value for this problem is zero, then we obtain a **feasible solution**.
- Otherwise, the original problem was **infeasible**.
- This is usually called the **Phase I LP**.

Obtaining a Basis

- The only difficulty occurs when the solution to **Phase I** is degenerate and there are still artificial variables in the basis.
- In this case, we need to perform some *degenerate pivots* to rotate the artificial variables out of the basis.
- This is done in the normal way except that the pivot element doesn't have to be positive.
- If the pivot element is zero, then that row is linearly dependent on the others and we can eliminate it.
- After we rotate all the artificial variables out of the basis, we are left with a BFS and the corresponding basis.

Two-phase Simplex Method

1. By multiplying some of the constraints by -1, change the problem so that $b \geq 0$.
2. Introduce auxiliary variables y_1, \dots, y_m , if necessary, and solve the auxiliary problem (Phase I).
3. If the optimal cost for the auxiliary problem is zero, then a **feasible solution** has been found. Otherwise, the problem is **infeasible**.
4. Eliminate any zero rows.
5. Rotate any remaining basic artificial variables out of the basis.
6. Drop the (nonbasic) artificial variables from the problem and compute the reduced costs for the nonbasic variables with the original objective function.
7. Continue the simplex algorithm as usual (Phase II).

The Big M method

- In the above method, we first solve the **Phase I LP** to obtain a BFS.
- Using that BFS, we start **Phase II**, which is solving the original problem.
- Another approach is to combine **Phase I** and **Phase II**.
- In this approach, we add the artificial variables and then change the objective function to

$$\sum_{j=1}^n c_j x_j + M \sum_{i=1}^m y_i$$

- M has to be large enough to force the artificial variables to zero.
- If the optimal solution has any artificial variables at nonzero level, then the original problem was **infeasible**.
- Otherwise, we obtain an **optimal solution**.
- In practice, two-phase simplex is usually used.

Computational Efficiency of the Simplex Method

- The **efficiency** of the method depends on the number of iterations.
- The number of iterations depends on how many extreme points are visited.
- It is easy to construct an example where there are 2^n extreme points and all of them are visited.
- This means in the worst case, the simplex method requires an **exponential number of iterations**.
- Of course, this depends on the pivoting rule, etc.
- What is the **best case**?

The Diameter of a Polyhedron

- The *distance* between two vertices of a polyhedron is the minimum number of pivots to get from one to the other.
- The *diameter* of a polyhedron is the maximum of the distances between all pairs of vertices.
- The diameter tells us something about the number of pivots we might be forced to make *in the best case*.
- Unfortunately, we don't know much about the diameter of general polyhedra.
- *On average*, the simplex algorithm performs extremely well if implemented correctly.