


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:


<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Topology and its Applications

www.elsevier.com/locate/topol

Homotopy type and v_1 -periodic homotopy groups of p -compact groups

Donald M. Davis

Lehigh University, Bethlehem, PA 18015, USA

ARTICLE INFO

Article history:

Received 15 July 2008

Accepted 15 July 2008

MSC:

55Q52

57T20

55N15

Keywords:

 v_1 -periodic homotopy groups p -compact groups

Adams operations

 K -theory

Invariant theory

ABSTRACT

We determine the v_1 -periodic homotopy groups of all irreducible p -compact groups (BX, X) . In the most difficult, modular, cases, we follow a direct path from their associated invariant polynomials to these homotopy groups. We show that, if p is odd, every irreducible p -compact group has X of the homotopy type of a product of explicit spaces related to p -completed Lie groups.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

In [5] and [4], the classification of irreducible p -compact groups was completed. This family of spaces extends the family of (p -completions of) compact simple Lie groups. The v_1 -periodic homotopy groups of any space X , denoted $v_1^{-1}\pi_*(X)_{(p)}$, are a localization of the portion of the homotopy groups detected by K -theory; they were defined in [20]. In [17] and [16], the author completed the determination of the v_1 -periodic homotopy groups of all compact simple Lie groups. Here we do the same for all the remaining irreducible p -compact groups.¹

Recall that a p -compact group [22] is a pair (BX, X) such that BX is p -complete and $X = \Omega BX$ with $H^*(X; \mathbb{F}_p)$ finite. Thus BX determines X and contains more structure than does X . The homotopy type and homotopy groups of X do not take into account this extra structure nor the multiplication on X . We show that, if p is odd, every irreducible p -compact group has X of the homotopy type of a product of explicit spaces related to p -completed Lie groups.

According to [5, 1.1, 11.1] and [4], the irreducible p -compact groups correspond to compact simple Lie groups² and the p -adic reflection groups listed in [2, Table 1] for which the character field is strictly larger than \mathbb{Q} . See [13, pp. 430–431] and [27, p. 165] for other listings of reflection groups. We use the usual notation $((BX_n)_p, (X_n)_p)$, where n is the Shephard–Todd numbering ([36] or any of the previously-mentioned tables) and p is the prime associated to the completion.

We will divide our discussion into four families of cases:

- (1) The compact simple Lie groups—infinite family 1, part of infinite family 2, and cases 28, 35–37 in the Shephard–Todd list.
- (2) The rest of the infinite families numbered 2a, 2b, and 3.

E-mail address: dmd1@lehigh.edu.

¹ If the groups $v_1^{-1}\pi_i(X)$ are finite, then p -completion induces an isomorphism $v_1^{-1}\pi_*(X) \rightarrow v_1^{-1}\pi_*(X_p)$ [9, p. 1252].

² Cases in which distinct compact Lie groups give rise to equivalent p -compact groups are discussed in [5, 11.4].

- (3) The nonmodular special cases, in which p does not divide the order of the reflection group. This is cases 4–27 and 29–34.
- (4) The modular cases, in which p divides the order of the reflection group. These are cases $(X_{12})_3$, $(X_{24})_2$, $(X_{29})_5$, $(X_{31})_5$, and $(X_{34})_7$. (Actually, we include $(X_{12})_3$ in case (3) along with the nonmodular cases, and the Dwyer–Wilkerson space $(X_{24})_2$ was handled in [6].)

Here is a brief summary of what we accomplish in each case. The author feels that his contributions here are nil in case (1),³ minuscule in case (2), modest in case (3), and significant in case (4).

- (1) Spaces X_1 , X_{28} , X_{35} , X_{36} , and X_{37} are, respectively, $SU(n)$, F_4 , E_6 , E_7 , and E_8 . These are p -compact groups for all primes p , although for small primes they were excluded by Clark and Ewing [13] because $H^*(BX; \mathbb{F}_p)$ is not a polynomial algebra. The exceptional Lie group G_2 is the case $m = 6$ in infinite family 2b. The spaces $SO(n)$, $Spin(n)$, and $Sp(n)$ appear in the infinite family 2a with $m = 2$. Simplification of the homotopy types of many of these, when p is odd, to products of spheres and spherically-resolved spaces was obtained in [31, (8.1), 8.1]. The v_1 -periodic homotopy groups of these spaces were computed in [18,7,17,16,8], and other papers. We will say no more about these cases.
- (2) In Section 2, we use work of Castellana and Broto and Møller to show that the spaces in the infinite families can be decomposed, up to homotopy, as products of factors of p -completions of unitary groups, spheres, and sphere bundles over spheres. See Corollary 2.3, Remark 2.6, and Theorem 2.8 for the specific results.
- (3) In Table 3.2, we list the homotopy types of all cases $(X_n)_p$ which are not products of spheres. There are 31 such cases. In each case, we give the homotopy type as a product of spheres and spaces which are spherically resolved with α_1 attaching maps. In Remark 3.3, we discuss the easily-computed v_1 -periodic homotopy groups of these spaces.
- (4) The most novel part of the paper is the determination of the v_1 -periodic homotopy groups of $(X_{29})_5$, $(X_{31})_5$, and $(X_{34})_7$. We introduce a direct, but nontrivial, path from the invariant polynomials to the v_1 -periodic homotopy groups. En route, we determine the Adams operations in $K^*(BX; \hat{\mathbb{Z}}_p)$ and $K^*(X; \hat{\mathbb{Z}}_p)$. In the case of $(X_{34})_7$, we give new explicit formulas for the invariant polynomials. We prove in Theorem 4.15 (resp. Theorem 5.17) that the homotopy type of $(X_{29})_5$ (resp. $(X_{34})_7$) is directly related to $SU(20)$ (resp. $SU(42)$). We prove in Theorem 4.20 that $(X_{31})_5$ has the homotopy type of the 5-completion of a factor of E_8 . These latter homotopy-type results and their proofs were pointed out and explained to the author by John Harper.

Recall that the universal cover of a connected p -compact group splits as a product of a torus and a number of simply-connected irreducible p -compact groups. Thus our results yield the v_1 -periodic homotopy groups of any connected p -compact group.

2. Infinite families 2 and 3

Family 3 consists of p -completed⁴ spheres S^{2m-1} with $p \equiv 1 \pmod m$, which is a loop space due to work of Sullivan [37]. The groups $v_1^{-1}\pi_*(S^{2m-1})_{(p)}$, originally due to Mahowald ($p = 2$) and Thompson (p odd), are given in [19, 4.2].

Family 2 consists of spaces $X(m, r, n)$ where $m > 1$, $r|m$, and $n > 1$. The “degrees” of $X(m, r, n)$ are $m, 2m, \dots, (n-1)m, \frac{m}{r}n$. These are the degrees of invariant polynomials under a group action used in defining the space. The Clark–Ewing table doubles the degrees to form the “type”, as these doubled degrees are the degrees of generators of $H^*(BX; \mathbb{F}_p)$ in the cases which they consider. For most⁵ of the irreducible p -compact groups X , $H^*(X; \mathbb{F}_p)$ is an exterior algebra on classes of grading $2d - 1$, where d ranges over the degrees. Family 2b consists of spaces $X(m, r, n)$ in which $n = 2$ and $r = m$, while family 2a is all other cases. The reason that these are separated is that 2b has more applicable primes. Indeed, for family 2a, there are p -compact groups when $p \equiv 1 \pmod m$, while for family 2b, these exist when $p \equiv \pm 1 \pmod m$, and also $p = 2$ if $m = 4$ or 6 , and $p = 3$ if $m = 3$ or 6 . The case $m = 6$ in family 2b is the exceptional Lie group G_2 . Note that all primes work when $m = 6$. The case $(p = 2, m = 4)$ has $X = Sp(2)$ or $PSp(2)$, while $(p = 3, m = 3)$ has $X = SU(3)$ or $PSU(3)$, the projective unitary group. In these cases, there are two inequivalent p -compact groups corresponding to the same \hat{Q}_p -reflection group; however, since $Sp(2) \rightarrow PSp(2)$ and $SU(3) \rightarrow PSU(3)$ are covering spaces, they induce isomorphisms of v_1 -periodic homotopy groups.

The following results of Broto and Møller [11] and Castellana [12] will be useful. They deal with the homotopy fixed-point space X^{hG} when G acts on a space of the same homotopy type as a space X . Here and throughout, C_m denotes a cyclic group of order m , and $U(N)$ is the p -completion of a unitary group.

Theorem 2.1. (See [11, 5.2, A.9].) *Let $m|(p - 1)$, $0 \leq s$, $\max(s, 1) < m$, and $n > 0$. Then C_m acts on $U(mn + s)$ via unstable Adams operations, and*

$$U(mn + s)^{hC_m} \simeq X(m, 1, n)$$

and is a factor in a product decomposition of $U(mn + s)$.

³ But he accomplished much in these cases in earlier papers such as [16–18].

⁴ All of our spaces are completed at an appropriate prime p . This will not always be present in our notation. For example, we will often write $SU(n)$ when we really mean its p -completion.

⁵ According to [34], the only exclusions are certain compact Lie groups when p is very small.

Theorem 2.2. (See [11, 5.2, A.11].) Let $m|(p - 1)$, $m \geq 2$, $r > 1$, and $n \geq 2$. Then there is an exotic action of C_m on $X(m, r, n)$ defined just before [11, A.11], and

$$X(m, r, n)^{hC_m} \simeq X(m, 1, n - 1)$$

is a factor in a product decomposition of $X(m, r, n)$.

Corollary 2.3. If $m|(p - 1)$ and $r > 1$, then

$$X(m, r, n) \simeq X(m, 1, n - 1) \times S^{2n\frac{m}{r}-1}$$

and $X(m, 1, n - 1)$ is a factor in a product decomposition of $U(m(n - 1))$.

Here $X(m, 1, 1)$ is interpreted as S^{2m-1} .

Proof. We use Theorem 2.2 to get the first factor. By the Kunnetth theorem, the other factor must have the same \mathbb{F}_p -cohomology as $S^{2n\frac{m}{r}-1}$, and hence must have the same homotopy type as this sphere. Now we apply Theorem 2.1 to complete the proof. \square

Remark 2.4. Our Corollary 2.3 appears as [12, 1.4], except that she has an apparent typo regarding the dimension of the sphere. Also, neither she nor [11] have the restriction $r > 1$, but it seems that the result is false for $r = 1$, since by induction it would imply that $X(m, 1, n)$ is a product of spheres, which is not usually true.

Remark 2.5. The referee points out that Theorem 2.2 holds for the four cases $((m, r, n) = (2, 1, 2), (4, 2, 2), (3, 3, 3),$ or $(2, 2, 4))$ which were omitted from [11, A.11] because the only thing that matters is that $A(m, 1, n) \subseteq N_{GL(L)}(G(m, r, n))$.

Remark 2.6. Let p be odd. By [31], for any N , p -completed $SU(N)$ splits as a product of $(p - 1)$ spaces, each of which has $H^*(-; \mathbb{F}_p)$ an exterior algebra on odd dimensional classes of dimensions $b, b + q, \dots, b + tq$, for some integers b and t . Here and throughout $q = 2(p - 1)$. Our space $X(m, 1, n - 1)$ will be a product of $(p - 1)/m$ of these spaces for $SU(m(n - 1))$. The v_1 -periodic homotopy groups of these spaces can be read off from those of $SU(m(n - 1))$, since the $(p - 1)$ factors have v_1 -periodic homotopy groups in nonoverlapping dimensions. Thus, to the extent that [18] is viewed as being a satisfactory description of $v_1^{-1}\pi_*(SU(n))_{(p)}$,⁶ Corollary 2.3 gives $v_1^{-1}\pi_*(X(m, r, n))_{(p)}$ provided $m|(p - 1)$.

Example 2.7. Let $p = 7$. Then $X(2, 2, 6) \simeq X(2, 1, 5) \times S^{11}$. There is a product decomposition

$$(SU(10))_7 \simeq B(3, 15) \times B(5, 17) \times B(7, 19) \times S^9 \times S^{11} \times S^{13},$$

where $B(2n + 1, 2n + 13)$ denotes a 7-completed S^{2n+1} -bundle over S^{2n+13} with attaching map α_1 . Then

$$X(2, 1, 5) \simeq B(3, 15) \times B(7, 19) \times S^{11}.$$

What remains for family 2 is the cases 2b when $m|(p + 1)$. These are the spaces $X(m, m, 2)$ with $m|(p + 1)$. Let $B(3, 2p + 1)$ denote the p -completion of an S^3 -bundle over S^{2p+1} with attaching map α_1 .

Theorem 2.8. If $m|(p + 1)$, then

$$X(m, m, 2) \simeq \begin{cases} B(3, 2p + 1), & m = p + 1, \\ S^3 \times S^{2m-1}, & m < p + 1. \end{cases}$$

Proof. Let $X = X(m, m, 2)$ with $m|(p + 1)$. Then $H^*(X; \mathbb{F}_p) = \Lambda[x_3, x_{2m-1}]$. If $m < p + 1$, then by the unstable Adams spectral sequence [10], both classes x_3 and x_{2m-1} are spherical. Indeed, the E_2 -term begins with towers in dimensions 3 and $2m - 1$ emanating from filtration 0, and no possible differentials. See Diagram 3.4. Because X is an H -space, the maps $S^3 \rightarrow X$ and $S^{2m-1} \rightarrow X$ yield a map $S^3 \times S^{2m-1} \rightarrow X$, and it is a p -equivalence by Whitehead's theorem.

On the other hand, suppose $m = p + 1$. We will show that $\mathcal{P}^1(x_3) = x_{2p+1}$. It then follows from [31, 7.1] that there is a p -equivalence $B(3, 2p + 1) \rightarrow X$.

To see that $\mathcal{P}^1(x_3) = x_{2p+1}$, we use the classifying space BX , which satisfies $H^*(BX; \mathbb{F}_p) = \mathbb{F}_p[y_4, y_{2p+2}]$. We will prove that $\mathcal{P}^1(y_4) = y_{2p+2} + Ay_4^{(p+1)/2}$, for some generator y_{2p+2} and some $A \in \mathbb{F}_p$, from which the desired result about the x 's follows immediately from the map $\Sigma X \rightarrow BX$, which in $H^*(-; \mathbb{F}_p)$ sends y_{j+1} to x_j and sends products to 0.

⁶ [18, 1.4] states that $v_1^{-1}\pi_{2k}(SU(n))_{(p)}$ is a cyclic p -group with exponent $\min(v_p(j!S(k, j)): j \geq n)$, where $S(-, -)$ denotes the Stirling number of the second kind. In [21], more tractable formulas were obtained if $n \leq p^2 - p + 1$. Here and throughout, $v_p(-)$ is the exponent of p .

First note that

$$\begin{aligned} \mathcal{P}^1(y_4) &= Ay_4^{(p+1)/2} + By_{2p+2}, \\ \mathcal{P}^1(y_{2p+2}) &= Cy_4^p + Dy_4^{(p-1)/2}y_{2p+2}, \end{aligned}$$

for some A, B, C, D in \mathbb{F}_p . By the unstable property of the Steenrod algebra,

$$\mathcal{P}^{p+1}(y_{2p+2}) = y_{2p+2}^p. \tag{2.9}$$

We must have

$$\mathcal{P}^p(y_{2p+2}) = \sum_{j=0}^{p-1} c_j y_4^{1+j(p+1)/2} y_{2p+2}^{p-1-j},$$

for some $c_j \in \mathbb{F}_p$. Since $\mathcal{P}^{p+1} = \mathcal{P}^1 \mathcal{P}^p$ and

$$\mathcal{P}^1(y_4^i y_{2p+2}^j) = i \mathcal{P}^1(y_4) y_4^{i-1} y_{2p+2}^j + j y_4^i \mathcal{P}^1(y_{2p+2}) y_{2p+2}^{j-1}, \tag{2.10}$$

the only way to obtain (2.9) is if $c_0 B = 1$ in \mathbb{F}_p . Thus B must be a unit, and the generator y_{2p+2} can be chosen so that $B = 1$. \square

3. Nonmodular individual cases

In this section, we consider all cases 4–34, excluding case 28 (which is F_4), in the Shephard–Todd numbering in which p does not divide the order of the reflection group. We obtain a very attractive result. One modular case, $(X_{12}, p = 3)$ is also included here. There is some overlap of our methods and results here with those in [29].

Theorem 3.1. *Let $X = (X_n)_p$ with $4 \leq n \leq 34$ and $n \neq 28$, excluding the modular cases $(X_{29})_5$, $(X_{31})_5$, and $(X_{34})_7$, which will be considered in the next two sections, and the modular case $(X_{24})_2$, which was analyzed in [6]. Then $X \simeq \prod S^{2d-1}$, where $2d$ ranges over the integers listed as the “type” in [13], except for the 31 cases listed in Table 3.2. In these, each $B(-, \dots, -)$ is built by fibrations from spheres of the indicated dimensions, with α_1 as each attaching map, and occurs as a factor in a product decomposition of the p -completion of some $SU(N)$.*

We will call the integers d , which are $1/2$ times the “type” numbers of Clark–Ewing, the “degrees”.

Remark 3.3. The v_1 -periodic homotopy groups of $B(2n + 1, 2n + 2p - 1)$ were obtained in [8, 1.3]. Those of $B(11, 23, 35)_7$ and $B(23, 35, 47, 59)_7$ were obtained in [8, 1.4]. Using [21, 1.5, 1.9], we find that for $\epsilon = 0, 1$,

$$v_1^{-1} \pi_{2t-\epsilon}(B(11, 35, 59, 83))_{(13)} \approx \begin{cases} 0, & t \not\equiv 5 \pmod{12}, \\ \mathbb{Z}/13^{\max(f_5(t), f_{17}(t), f_{29}(t), f_{41}(t))}, & t \equiv 5 \pmod{12}, \end{cases}$$

where $f_\gamma(t) = \min(\gamma, 4 + v_{13}(t - \gamma))$, while

$$v_1^{-1} \pi_{2t-\epsilon}(B(11, 47, 83))_{(19)} \approx \begin{cases} 0, & t \not\equiv 5 \pmod{18}, \\ \mathbb{Z}/19^{\max(f'_5(t), f'_{23}(t), f_{41}(t))}, & t \equiv 5 \pmod{18}, \end{cases}$$

where $f'_\gamma(t) = \min(\gamma, 3 + v_{19}(t - \gamma))$.

Proof of Theorem 3.1. It is straightforward to check that the pairs (case, prime) listed in Table 3.2 are the only nonmodular cases in [2, Table 1] in which an admissible prime p satisfies that $(p - 1)$ divides the difference of distinct degrees. Indeed all other admissible primes have $(p - 1)$ greater than the maximum difference of degrees. For example, case 30 requires $p \equiv 1, 4 \pmod{5}$, and the degrees are 2, 12, 20, 30. The first few primes of the required congruence are 11, 19, and 29. Clearly 10, 18, and 28 divide differences of these degrees, but no larger $(p - 1)$ can. Thus the unstable Adams spectral sequence argument used in proving Theorem 2.8 works the same way here to show that X is a product of S^{2d-1} in all cases not appearing in Table 3.2. In the relevant range, the E_2 -term will consist only of infinite towers, one for each generator. The first deviation from that is a \mathbb{Z}/p in filtration 1 in homotopy dimension $(2d - 1) + (2p - 3)$, where d is the smallest degree. This will always be greater than the dimension of the largest S^{2d-1} . A different proof that X is a product of spheres if and only if $p \geq \max\{d\}$ is given in [3, 2.1].

The next step is to show that the Steenrod operation \mathcal{P}^1 in $H^*(X; \mathbb{F}_p)$ must connect all the classes listed as adjacent generators in one of the B -spaces in Table 3.2. This was achieved independently of, and slightly earlier than, the author in [33] and [25]. We include several of our proofs, omitting the most complicated cases, to illustrate our methods and for the benefit of the reader without access to [33] and [25]. We accomplish this by considering the A -module $H^*(BX; \mathbb{F}_p)$.

Table 3.2
Cases in Theorem 3.1 which are not products of spheres

Case	Prime	Space
5	7	$B(11, 23)$
8	5	$B(15, 23)$
9	17	$B(15, 47)$
10	13	$B(23, 47)$
12	3	$B(11, 15)$
14	19	$B(11, 47)$
16	11	$B(39, 59)$
17	41	$B(39, 119)$
18	31	$B(59, 119)$
20	19	$B(23, 59)$
24	11	$B(7, 27) \times S^{11}$
25	7	$B(11, 23) \times S^{17}$
26	7	$B(11, 23, 35)$
26	13	$B(11, 35) \times S^{23}$
27	19	$B(23, 59) \times S^{13}$
29	13	$B(15, 39) \times S^7 \times S^{23}$
29	17	$B(7, 39) \times S^{15} \times S^{23}$
30	11	$B(3, 23) \times B(39, 59)$
30	19	$B(3, 39) \times S^{27} \times S^{59}$
30	29	$B(3, 59) \times S^{23} \times S^{39}$
31	13	$B(15, 39) \times B(23, 47)$
31	17	$B(15, 47) \times S^{23} \times S^{39}$
32	7	$B(23, 35, 47, 59)$
32	13	$B(23, 47) \times B(35, 59)$
32	19	$B(23, 59) \times S^{35} \times S^{47}$
33	7	$B(7, 19) \times B(11, 23, 35)$
33	13	$B(11, 35) \times S^7 \times S^{19} \times S^{23}$
34	13	$B(11, 35, 59, 83) \times B(23, 47)$
34	19	$B(11, 47, 83) \times B(23, 59) \times S^{35}$
34	31	$B(23, 83) \times S^{11} \times S^{35} \times S^{47} \times S^{59}$
34	37	$B(11, 83) \times S^{23} \times S^{35} \times S^{47} \times S^{59}$

With one exception⁷, all cases involving factors of $B(2m - 1, 2m + 2p - 3)$ are implied by Lemma 3.9 by applying $H^*(BX) \rightarrow H^{*-1}(X)$, which sends products to 0. Similarly, Lemma 3.11 covers the two cases with a factor $B(11, 23, 35)$.

Now we must show that the spaces X have the homotopy type claimed. The first 10 cases are immediate from [31, 7.1], and the two other nonproduct cases, i.e. $(X_{26})_7$ and $(X_{32})_7$, follow from [31, 7.2, 7.6]. Note that these results of [31] did not deal with p -completed spaces, but the obstruction theory arguments used there apply in the p -complete context. There are 15 additional types which we claim to be quasi p -regular. As defined in [32], a space is quasi p -regular if it is p -equivalent to a product of spheres and spaces of the form $B(2n + 1, 2n + 2p - 1)$. In [32] (see especially [32, pp. 330–334]), many exceptional Lie groups are shown to be quasi p -regular (for appropriate p) using a skeletal approach. We could use that approach here, but we prefer to use the unstable Adams spectral sequence (UASS). The two methods are really equivalent.

Let $q = 2p - 2$. In Diagrams 3.4 and 3.5, we illustrate the UASS for S^{2n+1} in dimension less than $2n + pq - 1$ and for $B(2n + 1, 2n + q + 1)$ in dimension less than $2n + 3q - 3$. Diagram 3.4 gives a nice interpretation of the statement of the homotopy groups in [38, 13.4]. If $n \geq p$, the paired dots in Diagram 3.4 will not occur in the pictured range. The nice thing about these charts is that the \mathbb{F}_p -cohomology groups of our spaces X are known to agree with that of their putative product decomposition as unstable algebras over the Steenrod algebra, and are of the required universal form for the UASS to apply; hence their UASS has E_2 -term the sum of the relevant charts of spheres and B -spaces. In all cases, there will be no possible differentials.


One can check that in all 15 cases in which X is claimed to be quasi p -regular, the towers in $\text{UASS}(X)$ corresponding to the spheres and the bottom cell of each $B(2n + 1, 2n + q + 1)$ cannot support a differential, and hence yield maps from the sphere or S^{2n+1} into X . Next one checks that $\pi_{2n+q}(X) = 0$ and $\pi_{4n+q+1}(X) = 0$. As these are the groups in which the obstruction to extending the map $S^{2n+1} \rightarrow X$ over $B(2n + 1, 2n + q + 1)$ lie, we obtain the desired extension. Finally, we take the product of maps $B \rightarrow X$ and $S^{2d_i-1} \rightarrow X$, using the group structure of X , to obtain the desired p -equivalence from a product of spheres and B -spaces into X .

The remaining cases, $(X_{33})_7$, $(X_{34})_{13}$, and $(X_{34})_{19}$, are handled similarly. The E_2 -term of the UASS converging to $\pi_*(X)$ is isomorphic to that of its putative product decomposition. For example, $E_2(X_{34})_{13}$ is the sum of Diagram 3.5 with $n = 11$ and $q = 24$ plus Diagram 3.6. We can map $S^{23} \rightarrow X$ and $S^{11} \rightarrow X$ corresponding to generators of homotopy groups. Then we can extend the first map over the 47- and 70-cells because $\pi_{46}(X) = 0$ and $\pi_{69}(X) = 0$. This gives a map $B(23, 47) \rightarrow X$. Similarly we can extend the second map over cells of $B(11, 35, 59, 83)$ of dimension 46, 70, 94, 118, 142, 105, 129, 153,

⁷ The exception is $(X_{34})_{19}$, which is handled in [33]. We thank Nishinobu for pointing out a gap in the argument for this case which appeared in an earlier version of this paper.

and 188. Taking the product of these two maps, using the multiplication of X , yields the desired 13-equivalence $B(23, 47) \times B(11, 35, 59, 83) \rightarrow X$. The other two cases are handled similarly. \square

Diagram 3.4. $UASS(S^{2n+1})$ in $\dim < 2n + pq - 1$. Here $n < p$.



Where there is a pair of dots, the grading at the bottom refers to the one on right, and the other is in grading 1 less.

Diagram 3.5. $UASS(B(2n + 1, 2n + q + 1))$ in $\dim < 2n + 3q - 3$.





Diagram 3.6. $UASS(B(11, 35, 59, 83)_{13})$ in $\dim < 200$.



An alternate proof of Theorem 3.1 can be obtained using [15, 1.3, 1.4], which can be interpreted as the following theorem. In Section 6, we will provide a proof of a strengthened version of this result.

Theorem 3.7. (See [15, 1.3, 1.4].) *If X is an H -space of rank $r < p - 1$ with torsion-free homology, then there are H -spaces X_1, \dots, X_r with $X_1 = S^{n_1}$ and $X_r = X$, and there are fibrations $X_{i-1} \rightarrow X_i \rightarrow S^{n_i}$ for $2 \leq i \leq r$ as in the diagram*

$$\begin{array}{ccccccc}
 X_1 & \longrightarrow & X_2 & \longrightarrow & \cdots & \longrightarrow & X_r \\
 & & \downarrow & & & & \downarrow \\
 & & S^{n_2} & & & & S^{n_r}
 \end{array} \tag{3.8}$$

The homotopy type of the p -localization of X is determined by the elements of $\pi_{n_i-1}(X_{i-1})$ associated to these fibrations.

In order to apply this, one would still need to determine the \mathcal{P}^1 -action and to check that the relevant homotopy groups $\pi_{n_i-1}(X_{i-1})$ are cyclic.

In the following lemmas, which were used above, g_i denotes a generator in grading i .

Lemma 3.9.

- (a) *If $m \not\equiv 1 \pmod p$ and $\mathbb{F}_p[g_{2m}, g_{2m+2p-2}]$ is an unstable A -algebra, then $\mathcal{P}^1 g_{2m} \equiv u g_{2m+2p-2} \pmod{\text{decomposables}}$, with $u \neq 0$.*
- (b) *The same conclusion holds if the unstable A -algebra contains additional generators in dimensions $d \not\equiv 2m \pmod{(2p-2)}$, provided also that $d + 2p - 2 \not\equiv 0 \pmod{(2m + 2p - 2)}$.*

Proof. (a) For dimensional reasons, we must have $\mathcal{P}^1 g_{2m} = \alpha g_{2m+2p-2}$ plus possibly a power of g_{2m} , for some $\alpha \in \mathbb{F}_p$, and $\mathcal{P}^1 g_{2m+2p-2} = g_{2m} Y$, for some polynomial Y . The unstable condition requires that $\mathcal{P}^{m+p-1} g_{2m+2p-2} = g_{2m+2p-2}^p$, and, since $m \not\equiv 1 \pmod p$, this equals, up to a unit, $\mathcal{P}^1 \mathcal{P}^{m+p-2} g_{2m+2p-2}$. For dimensional reasons,

$$\mathcal{P}^{m+p-2} g_{2m+2p-2} = \beta g_{2m} g_{2m+2p-2}^{p-1} + g_{2m}^3 Z \tag{3.10}$$

for some $\beta \in \mathbb{F}_p$ and some polynomial Z . By the Cartan formula (similar to (2.10)), the only way that \mathcal{P}^1 applied to (3.10) can yield $g_{2m+2p-2}^p$ is if both α and β are units.

(b) There are two ways that the additional generators could affect the argument for part (a). One is if several of them (possibly the same one) were multiplied together to get into the congruence of part (a). By the Cartan formula, \mathcal{P}^1 of such a product will still involve some of these additional generators as factors, and so cannot yield the $g_{2m+2p-2}^p$ term on which the argument focuses. The other way, pointed out to the author by Nishinobu, would be if there were a generator g_d such that $\mathcal{P}^1 g_d = u g_{2m+2p-2}^i$ for some $u \neq 0$ and $i > 0$, an eventuality excluded by our second hypothesis. If there were such a g_d and also $\mathcal{P}^{m+p-2} g_{2m+2p-2}$ included a term $g_d g_{2m+2p-2}^{p-i}$, this would provide an alternative way to achieve $g_{2m+2p-2}^p$ in $\mathcal{P}^{m+p-1} g_{2m+2p-2}$. \square

Lemma 3.11. *If $\mathbb{F}_7[g_{12}, g_{24}, g_{36}]$ is an unstable A -algebra, then, mod decomposables, $\mathcal{P}^1 g_{12} = u_1 g_{24}$ and $\mathcal{P}^1 g_{24} = u_2 g_{36}$ with $u_i \neq 0$. The same conclusion holds for $\mathbb{F}_7[g_{12}, g_{24}, g_{36}, g_8, g_{20}]$.*

Proof. We work modulo the ideal (g_{12}) , resp. (g_{12}, g_8, g_{20}) . Then $\mathcal{P}^1 g_{12} \equiv \alpha g_{24}$, $\mathcal{P}^1 g_{24} \equiv \beta g_{36}$, and $\mathcal{P}^1 g_{36} \equiv \gamma g_{24}^2$, for some α, β, γ in \mathbb{F}_7 . This latter term complicates things somewhat. We also have $\mathcal{P}^1 g_8 \equiv 0$ and $\mathcal{P}^1 g_{20} \equiv 0$, in the second case.

The unstable condition implies $\mathcal{P}^1 \mathcal{P}^{17} g_{36} = u g_{36}^7$ with $u \neq 0$. We use the Cartan formula as in the previous proof. The only way to get to g_{36}^7 by \mathcal{P}^1 is if $\beta \neq 0$, implying the result for $\mathcal{P}^1 g_{24}$.

However, there are two ways that $\mathcal{P}^1 \mathcal{P}^{11} g_{24}$ might yield g_{24}^7 , one via $\mathcal{P}^1(g_{12} g_{24}^6)$ and the other via $\mathcal{P}^1(g_{24}^5 g_{36})$. Instead, we consider $(\mathcal{P}^1)^5 \mathcal{P}^7 g_{24}$. We must have

$$\mathcal{P}^7 g_{24} \equiv \delta_1 g_{36}^3 + \delta_2 g_{24}^3 g_{36}$$

for some $\delta_i \in \mathbb{F}_7$.

Assume $\alpha = 0$, so that the ideal (g_{12}) or (g_{12}, g_8, g_{20}) is closed under the action of \mathcal{P}^1 . Then we can compute

$$\begin{aligned}
 (\mathcal{P}^1)^5 g_{36}^3 &\equiv \beta \gamma^4 g_{24}^7 - \beta^3 \gamma^2 g_{24} g_{36}^4, \\
 (\mathcal{P}^1)^5 (g_{24}^3 g_{36}) &\equiv \beta^2 \gamma^3 g_{24}^7 + 5 \beta^3 \gamma^2 g_{24}^4 g_{36}^2,
 \end{aligned}$$

and then

$$\begin{aligned} ug_{24}^7 &\equiv (\mathcal{P}^1)^5 \mathcal{P}^7(g_{24}) \\ &\equiv (\mathcal{P}^1)^5 (\delta_1 g_{36}^3 + \delta_2 g_{24}^3 g_{36}) \\ &\equiv \delta_1 (\beta \gamma^4 g_{24}^7 - \beta^3 \gamma^2 g_{24}^4 g_{36}^4) + \delta_2 (\beta^2 \gamma^3 g_{24}^7 + 5\beta^3 \gamma^2 g_{24}^4 g_{36}^2). \end{aligned}$$

Coefficients of g_{24}^7 imply $\beta \neq 0$, $\gamma \neq 0$, and some $\delta_i \neq 0$, but this then gives a contradiction regarding $g_{24}^4 g_{36}^4$ or $g_{24}^4 g_{36}^2$. Thus the assumption that $\alpha = 0$ must have been false. \square

4. 5-primary modular cases

In this section, we determine the v_1 -periodic homotopy groups of the modular 5-compact groups X_{29} and X_{31} . We pass directly from invariant polynomials to Adams operations in $K^*(X)$ and thence to $v_1^{-1}\pi_*(X)$. In Theorems 4.15 and 4.20, we relate the homotopy type of $(X_{29})_5$ and $(X_{31})_5$ to that of $SU(20)$ and E_8 . Theorem 4.15 was conjectured by the author in an earlier version of this paper. It and Theorem 4.20 and their proofs were provided to the author by John Harper.

The input to determining the Adams module $K^*(X_{29}; \hat{\mathbb{Z}}_5)$ is the following result due to Aguadé [1] and Maschke [30]. Throughout the rest of the paper, we will denote by $m_{(e_1, \dots, e_k)}$ the smallest symmetric polynomial on variables x_1, \dots, x_ℓ (the value of ℓ will be implicit) containing the term $x_1^{e_1} \cdots x_k^{e_k}$.

Theorem 4.1. *There is a reflection group G_{29} acting on $(\hat{\mathbb{Z}}_5)^4$, and there is a space BX_{29} and map $BT \rightarrow BX_{29}$ with $BT = K((\hat{\mathbb{Z}}_5)^4, 2)$ such that*

$$H^*(BX_{29}; \hat{\mathbb{Z}}_5) \approx H^*(BT; \hat{\mathbb{Z}}_5)^{G_{29}},$$

the invariants under the natural action of G_{29} on $H^*(BT; \hat{\mathbb{Z}}_5) = \hat{\mathbb{Z}}_5[x_1, x_2, x_3, x_4]$ with $|x_i| = 2$. Moreover, $H^*(BT; \hat{\mathbb{Z}}_5)^{G_{29}}$ is a polynomial algebra on the following four invariant polynomials:

$$\begin{aligned} f_4 &= m_{(4)} - 12m_{(1,1,1,1)}, \\ f_8 &= m_{(8)} + 14m_{(4,4)} + 168m_{(2,2,2,2)}, \\ f_{12} &= m_{(12)} - 33m_{(8,4)} + 330m_{(4,4,4)} + 792m_{(6,2,2,2)}, \\ f_{20} &= m_{(20)} - 19m_{(16,4)} - 494m_{(12,8)} - 336m_{(14,2,2,2)} + 716m_{(12,4,4)} \\ &\quad + 1038m_{(8,8,4)} + 7632m_{(10,6,2,2)} + 129012m_{(8,4,4,4)} + 106848m_{(6,6,6,2)}. \end{aligned}$$

Proof. The group G_{29} is the subgroup of $GL(4, \mathbf{C})$ generated by the following four matrices. These can be seen explicitly in [1].

$$\frac{1}{2} \begin{pmatrix} 1 & -1 & -1 & -1 \\ -1 & 1 & -1 & -1 \\ -1 & -1 & 1 & -1 \\ -1 & -1 & -1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Since $i \in \hat{\mathbb{Z}}_5$, these act on $(\hat{\mathbb{Z}}_5)^4$, and this induces an action on

$$H^*(K((\hat{\mathbb{Z}}_5)^4, 2)) \approx \hat{\mathbb{Z}}_5[x_1, x_2, x_3, x_4].$$

The invariants of this action were determined by Maschke [30] to be the polynomials stated in the theorem. Although he did not state them all explicitly, they can be easily generated by: (a) define ϕ , ψ_i , and χ as on his p. 501, then (b) define Φ_1, \dots, Φ_6 as on his p. 504, and finally (c) let $f_4 = -\frac{1}{2}\Phi_6$ and $f_8 = F_8$, $f_{12} = F_{12}$, and $f_{20} = F_{20}$ as on his p. 505. See also [36, p. 287] for a reference to this work.

Actually, Maschke's work and that of [36] involved finding generators for the complex invariant ring. To see that these integral polynomials generate the invariant ring over $\hat{\mathbb{Z}}_5$, one must show that they cannot be decomposed over $\mathbb{Z}/5$. For example, one must verify that f_{20} cannot be decomposed mod 5 as a linear combination of $f_8 f_{12}$, $f_4^2 f_{12}$, $f_4 f_8^2$, $f_4^3 f_8$, and f_4^5 . The need to do this was pointed out to the author by Kasper Andersen in a dramatic way, as will be described prior to Theorem 5.6. The verification here was performed by Andersen using a Magma program.

Aguadé [1] constructed the 5-compact group (BX, X) corresponding to this modular reflection group. \square

The approach based on the following proposition benefits from a suggestion of Clarence Wilkerson. Here we employ notation, standard in the p -compact group literature, that $H^*(-; \hat{\mathbb{Q}}_p)$ is short for $H^*(-; \hat{\mathbb{Z}}_p) \otimes_{\hat{\mathbb{Z}}_p} \hat{\mathbb{Q}}_p$.

Proposition 4.2. Let (BX, X) be a p -compact group with Weyl group G acting on $BT = K((\hat{\mathbb{Z}}_p)^n, 2)$. Suppose $H^*(BT; \hat{\mathbb{Q}}_p)^G = \hat{\mathbb{Q}}_p[f_1, \dots, f_k]$, where f_i is a polynomial in y_1, \dots, y_n with $y_j \in H^2(BT; \hat{\mathbb{Q}}_p)$ corresponding to the j th factor. Let $K^*(BT; \hat{\mathbb{Q}}_p) = \hat{\mathbb{Q}}_p[[x_1, \dots, x_n]]$ with x_i the class of $H - 1$ in the i th factor, where H is the complex Hopf bundle. Let $\ell_0(x) = \log(1 + x)$. Then

$$K^*(BX; \hat{\mathbb{Z}}_p) \approx \hat{\mathbb{Q}}_p[[f_1(\ell_0(x_1), \dots, \ell_0(x_n)), \dots, f_k(\ell_0(x_1), \dots, \ell_0(x_n))]] \cap \hat{\mathbb{Z}}_p[[x_1, \dots, x_n]].$$

Proof. The Chern character $K^*(BT; \hat{\mathbb{Q}}_p) \xrightarrow{\text{ch}} H^*(BT; \hat{\mathbb{Q}}_p)$ satisfies $\text{ch}(\ell_0(x_i)) = y_i$ and hence, since ch is a ring homomorphism, $\text{ch}(f_j(\ell_0(x_1), \dots, \ell_0(x_n))) = f_j(y_1, \dots, y_n)$. It commutes with the action of G , and hence sends invariants to invariants. Indeed

$$K^*(BT; \hat{\mathbb{Q}}_p)^G = \hat{\mathbb{Q}}_p[[f_1(\ell_0(x_1), \dots, \ell_0(x_n)), \dots, f_k(\ell_0(x_1), \dots, \ell_0(x_n))]]. \tag{4.3}$$

The invariant ring in $K^*(BT; \hat{\mathbb{Z}}_p)$ is just the intersection of (4.3) with $\hat{\mathbb{Z}}_p[[x_1, \dots, x_n]]$. Finally we use a result of [26] that $K^*(BX; \hat{\mathbb{Z}}_p) \approx K^*(BT; \hat{\mathbb{Z}}_p)^G$.

As pointed out by the referee, a standard Galois theory argument implies that $k = n$ in this proposition. \square

Thus with f_4, f_8, f_{12}, f_{20} as in Theorem 4.1, we wish to find algebraic combinations of

$$f_4(\ell_0(x_1), \dots, \ell_0(x_4)), \dots, f_{20}(\ell_0(x_1), \dots, \ell_0(x_4))$$

which have coefficients in $\hat{\mathbb{Z}}_5$. A theorem of [26] which states that for a p -compact group BX there is an isomorphism $K^*(BX; \hat{\mathbb{Z}}_p) \approx \hat{\mathbb{Z}}_p[[g_1, \dots, g_k]]$, and the collapsing, for dimensional reasons, of the Atiyah–Hirzebruch spectral sequence

$$H^*(BX; K^*(\text{pt}; \hat{\mathbb{Z}}_p)) \Rightarrow K^*(BX; \hat{\mathbb{Z}}_p) \tag{4.4}$$

implies that the generators g_j can be chosen to be of the form $f_j(x_1, \dots, x_n) \bmod$ higher degree polynomials.

Finding these algebraic combinations can be facilitated by using the p -typical log series

$$\ell_p(x) = \sum_{n \geq 0} x^{p^n} / p^n.$$

By [24], there is a series $h(x) \in \mathbb{Z}_{(p)}[[x]]$ such that $\ell_0(h(x)) = \ell_p(x)$ and $h(x) \equiv x \pmod{x^2}$. Let $x'_i = h(x_i)$. For any $c_e \in \hat{\mathbb{Q}}_p$ with $e = (e_4, e_8, e_{12}, e_{20})$, we have

$$\sum c_e f_4(\ell_p(x_1), \dots, \ell_p(x_4))^{e_4} \cdots f_{20}(\ell_p(x_1), \dots, \ell_p(x_4))^{e_{20}} \tag{4.5}$$

$$= \sum c_e f_4(\ell_0(x'_1), \dots, \ell_0(x'_4))^{e_4} \cdots f_{20}(\ell_0(x'_1), \dots, \ell_0(x'_4))^{e_{20}}, \tag{4.6}$$

where the sums are taken over various e . We will find c_e so that (4.5) is in $\hat{\mathbb{Z}}_p[[x_1, \dots, x_4]]$. Thus so is (4.6), and hence also $\sum c_e f_4(\ell_0(x_1), \dots, \ell_0(x_4))^{e_4} \cdots f_{20}(\ell_0(x_1), \dots, \ell_0(x_4))^{e_{20}}$, since $h(x) \in \mathbb{Z}_{(p)}[[x]]$.

A Maple program, which implements an algorithm described in the proof, was used to prove the following result.

Theorem 4.7. Let f_4, f_8, f_{12}, f_{20} be as in Theorem 4.1, and let

$$F_j = F_j(x_1, \dots, x_4) = f_j(\ell_0(x_1), \dots, \ell_0(x_4)).$$

Then the following series are 5-integral through grading 20; i.e., their coefficients of all monomials $x_1^{e_1} \cdots x_4^{e_4}$ with $\sum e_i \leq 20$ are 5-integral.

$$\begin{aligned} &F_4 - \frac{1}{10}F_4^2 - \frac{1}{5}F_8 - \frac{16}{25}F_{12} - \frac{7}{25}F_4F_8 + \frac{4}{25}F_4^3 - \frac{13}{125}F_4F_{12} - \frac{57}{125}F_4^2F_8 \\ &\quad - \frac{102}{125}F_4^4 - \frac{62}{125}F_8^2 - \frac{64}{125}F_{20} - \frac{4}{625}F_4^5 - \frac{42}{125}F_4^2F_{12} - \frac{11}{25}F_4F_8^2 - \frac{72}{125}F_8F_{12}; \\ &F_8 - \frac{8}{5}F_{12} - \frac{7}{25}F_8^2 - \frac{4}{25}F_{20} - \frac{21}{125}F_8F_{12}; \\ &F_{12} - \frac{2}{5}F_8^2 - \frac{1}{5}F_{20} - \frac{4}{25}F_8F_{12}. \end{aligned}$$

Proof. As observed in the paragraph preceding the theorem, it suffices to show that the same is true for $\tilde{F}_j = f_j(x_1 + \frac{1}{5}x_1^5, \dots, x_4 + \frac{1}{5}x_4^5)$. The advantage of this is to decrease the number of terms which must be kept track of and looked at. We work one grading at a time, expanding relevant products of F_j 's as combinations of monomial symmetric polynomials in the fixed grading, and then solving a system of linear equations to find the combinations that work. We illustrate with the calculation for modifications of F_4 in gradings 8 and then 12.

In grading 8, we have

$$\begin{aligned} \tilde{F}_4 &= \frac{4}{5}m_{(8)} - \frac{12}{5}m_{(5,1,1,1)}, \\ \tilde{F}_8 &= m_{(8)} + 14m_{(4,4)} + 168m_{(2,2,2,2)}, \\ \tilde{F}_4^2 &= m_{(8)} - 24m_{(5,1,1,1)} + 2m_{(4,4)} + 144m_{(2,2,2,2)}. \end{aligned}$$

We wish to choose a and b so that, in grading 8, $\frac{a}{5}\tilde{F}_8 + \frac{b}{5}\tilde{F}_4^2 \equiv \tilde{F}_4 \pmod{\text{integers}}$. Thus we must solve a system of mod 5 equations for a and b with augmented matrix

$$\left(\begin{array}{cc|c} 1 & 1 & 4 \\ 0 & -24 & -12 \\ 14 & 2 & 0 \\ 168 & 144 & 0 \end{array} \right).$$

The solution is $a = 1$, $b = 1/2$. We could also have used $b = 3$ since we are working mod 5.

Let $\tilde{F}'_4 = \tilde{F}_4 - \frac{1}{10}\tilde{F}_4^2 - \frac{1}{5}\tilde{F}_8$. In grading 12, we have

$$\begin{aligned} \tilde{F}'_4 &= -\frac{6}{25}m_{(12)} - \frac{12}{5}m_{(8,4)} - \frac{96}{5}m_{(6,2,2,2)} + \frac{12}{5}m_{(9,1,1,1)} + \frac{12}{25}m_{(5,5,1,1)}, \\ \tilde{F}'_{12} &= m_{(12)} - 33m_{(8,4)} + 330m_{(4,4,4)} + 792m_{(6,2,2,2)}, \\ \tilde{F}_8\tilde{F}'_4 &= m_{(12)} + 15m_{(8,4)} + 42m_{(4,4,4)} + 168m_{(6,2,2,2)} - 12m_{(9,1,1,1)} - 168m_{(5,5,1,1)} - 2016m_{(3,3,3,3)}, \\ \tilde{F}_4^3 &= m_{(12)} + 3m_{(8,4)} + 6m_{(4,4,4)} + 432m_{(6,2,2,2)} - 36m_{(9,1,1,1)} - 72m_{(5,5,1,1)} - 1728m_{(3,3,3,3)}. \end{aligned}$$

We wish to choose a , b , and c so that, in grading 12, $\frac{a}{25}\tilde{F}'_{12} + \frac{b}{25}\tilde{F}_8\tilde{F}'_4 + \frac{c}{25}\tilde{F}_4^3 \equiv \tilde{F}'_4 \pmod{\text{integers}}$. Thus we must solve a system of equations mod 25 whose augmented matrix is

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & -6 \\ -33 & 15 & 3 & -60 \\ 330 & 42 & 6 & 0 \\ 792 & 168 & 432 & -480 \\ 0 & -12 & -36 & 60 \\ 0 & -168 & -72 & 12 \\ 0 & -2016 & -1728 & 0 \end{array} \right).$$

The solution is $a = 16$, $b = 7$, and $c = -4$.

We perform similar calculations for \tilde{F}_8 in grading 12, then for \tilde{F}'_4 , \tilde{F}'_8 , and \tilde{F}'_{12} in gradings 16 and then 20. \square

By the observation in the paragraph involving (4.4), the modified versions of F_4 , F_8 , and F_{12} given in Theorem 4.7, and also F_{20} , can be modified similarly in all subsequent gradings, yielding generators of the power series algebra $K^*(BX_{29}; \hat{\mathbb{Z}}_5)$ which we will call G_4 , G_8 , G_{12} , and G_{20} . By [26], $K^*(X_{29}; \hat{\mathbb{Z}}_5)$ is an exterior algebra on classes z_3 , z_7 , z_{11} , and z_{19} in $K^1(-)$ obtained using the map $e: \Sigma X = \Sigma \mathcal{O}BX \rightarrow BX$ and Bott periodicity $B: K^1(X) \rightarrow K^{-1}(X)$ by $z_i = B^{-1}e^*(G_{i+1})$. The following determination of the Adams operations is essential for our work on v_1 -periodic homotopy groups. Here and elsewhere $QK^1(-)$ denotes the indecomposable quotient.

Theorem 4.8. *The Adams operation ψ^k in $QK^1(X_{29}; \hat{\mathbb{Z}}_5)$ on the generators z_3 , z_7 , z_{11} , and z_{19} is given by the matrix*

$$\begin{pmatrix} k^3 & 0 & 0 & 0 \\ \frac{1}{5}k^3 - \frac{1}{5}k^7 & k^7 & 0 & 0 \\ \frac{24}{25}k^3 - \frac{8}{25}k^7 - \frac{16}{25}k^{11} & \frac{8}{5}k^7 - \frac{8}{5}k^{11} & k^{11} & 0 \\ \frac{92}{125}k^3 - \frac{12}{125}k^7 - \frac{16}{125}k^{11} - \frac{64}{125}k^{19} & \frac{12}{25}k^7 - \frac{8}{25}k^{11} - \frac{4}{25}k^{19} & \frac{1}{5}k^{11} - \frac{1}{5}k^{19} & k^{19} \end{pmatrix}.$$

Proof. We first note that

$$\psi^k(\ell_0(x)) = \ell_0(\psi^k x) = \ell_0((x+1)^k - 1) = \log((x+1)^k) = k \log(x+1) = k\ell_0(x).$$

Since F_{4j} is homogeneous of degree $4j$ in $\ell_0(x_i)$, $\psi^k(F_{4j}) = k^{4j}F_{4j}$. We can use this to determine ψ^k on the generators G_i which are defined as algebraic combinations of F_{4j} 's. We then apply e^* to this formula to obtain ψ^k in $K^{-1}(X_{29}; \hat{\mathbb{Z}}_5)$. Since e^* annihilates decomposables, we need consider only the linear terms in the expressions which express G_i in terms of F_{4j} 's. On the basis (over \mathbb{Q}_5) $(e^*(F_4), e^*(F_8), e^*(F_{12}), e^*(F_{20}))$, the matrix of ψ^k is $D = \text{diag}(k^4, k^8, k^{12}, k^{20})$. On the basis (over $\hat{\mathbb{Z}}_5$)

$$(e^*(G_4), e^*(G_8), e^*(G_{12}), e^*(G_{20})),$$

it is $P^{-1}DP$, where

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -\frac{1}{5} & 1 & 0 & 0 \\ -\frac{16}{25} & -\frac{8}{5} & 1 & 0 \\ -\frac{64}{125} & -\frac{4}{25} & -\frac{1}{5} & 1 \end{pmatrix}$$

is the change-of-basis matrix, obtained using the linear terms in Theorem 4.7. The matrix in the statement of the theorem is obtained by dividing $P^{-1}DP$ by k , since ψ^k in $K^1(-)$ corresponds to ψ^k/k in $K^{-1}(-)$. \square

We can use Theorem 4.8 to obtain the v_1 -periodic homotopy groups of $(X_{29})_5$ as follows.

Theorem 4.9. *The groups $v_1^{-1}\pi_*(X_{29})_{(5)}$ are given by*

$$v_1^{-1}\pi_{2t-1}(X_{29}) \approx v_1^{-1}\pi_{2t}(X_{29}) \approx \begin{cases} 0, & t \not\equiv 3 \pmod{4}, \\ \mathbb{Z}/5^3, & t \equiv 3, 15 \pmod{20}, \\ \mathbb{Z}/5^{\min(8, 3+v_5(t-7-4\cdot 5^4))}, & t \equiv 7 \pmod{20}, \\ \mathbb{Z}/5^{\min(12, 3+v_5(t-11-4\cdot 5^8))}, & t \equiv 11 \pmod{20}, \\ \mathbb{Z}/5^{\min(20, 3+v_5(t-19-12\cdot 5^{16}))}, & t \equiv 19 \pmod{20}. \end{cases}$$

Proof. We use the result of [9] that $v_1^{-1}\pi_{2t}(X)_{(5)}$ is presented by the matrix $\begin{pmatrix} (\psi^5)^T \\ (\psi^2)^T - 2^t I \end{pmatrix}$. We form this matrix by letting $k = 5$ and 2 in the matrix of Theorem 4.8 and letting $x = 2^t$, obtaining

$$\begin{pmatrix} 125 & -15\,600 & -31\,274\,880 & -9\,765\,631\,257\,408 \\ 0 & 78\,125 & -78\,000\,000 & -3\,051\,773\,400\,000 \\ 0 & 0 & 48\,828\,125 & -3\,814\,687\,500\,000 \\ 0 & 0 & 0 & 19\,073\,486\,328\,125 \\ 8-x & -24 & -1344 & -268\,704 \\ 0 & 128-x & -3072 & -84\,480 \\ 0 & 0 & 2048-x & -104\,448 \\ 0 & 0 & 0 & 524\,288-x \end{pmatrix}. \tag{4.10}$$

If $x \not\equiv 3 \pmod{5}$, which is equivalent to $t \not\equiv 3 \pmod{4}$, then the units along the diagonal of the bottom half of (4.10) imply that the group presented is 0. Henceforth, we assume $x \equiv 3 \pmod{5}$.

Pivoting on the units (over $\mathbb{Z}_{(5)}$) in positions (5, 2) and (7, 4) and removing their rows and columns does not change the group presented. We now have a 6-by-2 matrix, whose nonzero entries are polynomials in x of degree 1 or 2. The polynomial in new position (5, 2) is nonzero mod 5 for such x , and so we pivot on it, and remove its row and column. The five remaining entries are ratios of polynomials with denominator nonzero mod 5. Let p_1, \dots, p_5 denote the polynomials in the numerators. The group $v_1^{-1}\pi_{2t}(X_{29})_{(5)}$ is $\mathbb{Z}/5^e$, where $e = \min(v(p_1(x)), \dots, v(p_5(x)))$, where $x = 2^t$. We abbreviate $v_5(-)$ to $v(-)$ throughout the remainder of this section. We have

$$\begin{aligned} p_1 &= -71\,122\,941\,747\,658\,752 + 9\,480\,741\,773\,824\,512x - 74\,067\,383\,851\,199x^2 + 33\,908\,441\,866x^3, \\ p_2 &= -66\,750\,692\,556\,800\,000 + 8\,897\,903\,174\,800\,000x - 69\,512\,640\,100\,000x^2 + 31\,789\,306\,250x^3, \\ p_3 &= -8\,327\,872 \cdot 10^{10} + 11\,101\,145 \cdot 10^9x - 86\,731\,015\,625\,000x^2 + 39\,736\,328\,125x^3, \\ p_4 &= 4 \cdot 10^{19} - 533\,203\,125 \cdot 10^{10}x + 41\,656\,494\,140\,625\,000x^2 - 19\,073\,486\,328\,125x^3, \\ p_5 &= 1\,099\,511\,627\,776 - 146\,567\,856\,128x + 1\,145\,324\,544x^2 - 526\,472x^3 + x^4. \end{aligned}$$

For values of m listed in the table, we compute and present in Table 4.12 the tuples (e_0, e_1, e_2, e_3) so that, up to units,

$$p_i(2^m + y) = 5^{e_0} + 5^{e_1}y + 5^{e_2}y^2 + 5^{e_3}y^3 \tag{4.11}$$

(plus y^4 if $i = 5$). Considerable preliminary calculation underlies the choice of these values of m .

Table 4.12
Exponents of polynomials

m	i				
	1	2	3	4	5
3	3, 2, 1, 0	∞ , 7, 6, 5	∞ , 12, 12, 10	∞ , 21, 20, 19	∞ , 3, 2, 1
15	3, 2, 1, 0	8, 7, 6, 5	13, 12, 11, 10	22, 21, 20, 19	4, 4, 3, 2
$7 + 4 \cdot 5^4$	8, 2, 1, 0	8, 7, 6, 5	17, 12, 11, 10	26, 21, 21, 19	8, 3, 2, 1
$11 + 4 \cdot 5^8$	12, 2, 1, 0	12, 7, 7, 5	13, 12, 11, 10	30, 21, 20, 19	12, 3, 2, 1
$19 + 12 \cdot 5^{16}$	23, 2, 2, 0	20, 7, 6, 5	21, 12, 11, 10	22, 21, 20, 19	20, 3, 2, 1

Recall that $v(2^{4 \cdot 5^i} - 1) = i + 1$, as is easily proved by induction. Thus

$$p(2^{m+20j}) = p(2^m + 2^m(2^{20j} - 1)) = p(2^m + 25j \cdot u), \tag{4.13}$$

with u a unit. Hence

$$\min\{v(p_i(2^{3+20j})): 1 \leq i \leq 5\} = 3$$

since

$$p_1(2^{3+20j}) = p_1(2^3 + 5^2 ju) = 5^3 + 5^2 \cdot 5^2 ju + 5(5^2 ju)^2 + (5^2 ju)^3,$$

omitting some unit coefficients. Here we have set $y = 5^2 ju$ in (4.11). Replacing 3 by 15 yields an identical argument. This yields the second line of Theorem 4.9.

We use Table 4.12 to show

$$\min\{v(p_i(2^{19+12 \cdot 5^{16}+20j})): 1 \leq i \leq 5\} = \min(20, 4 + v(j)) = \min(20, 3 + v(20j)). \tag{4.14}$$

Indeed, for $v(j) \leq 16$, the minimum is achieved when $i = 1$, with the 4 coming as $2 + 2$ with one 2 being from the 25 in (4.13) and the other 2 being the first 2 in the last row of Table 4.12. If $v(j) > 16$, the minimum is achieved when $i = 2$, using the first 20 in the last row of Table 4.12. The last case of Theorem 4.9 follows easily from (4.14), and the other two parts of Theorem 4.9 are obtained similarly.

To see that $v_1^{-1}\pi_{2t-1}(X_{29}) \approx v_1^{-1}\pi_{2t}(X_{29})$, we argue in three steps. First, the two groups have the same order using [9, 8.5] and the fact that the kernel and cokernel of an endomorphism of a finite group have equal orders. Second, by [16, 4.4], a presentation of $v_1^{-1}\pi_{2t-1}(X_{29})$ is given by $(\begin{smallmatrix} \psi^5 \\ \psi^2 - 2^t \end{smallmatrix})$, i.e. like that for $v_1^{-1}\pi_{2t}(X_{29})$ except that the two submatrices are not transposed. Third, we pivot on this matrix, which is (4.10) with the top and bottom transposed, and find that we can pivot on units three times, so that the group presented is cyclic. \square

One of the factors in the product decomposition of $SU(20)_5$ given in [31] is an H -space $B_3^5(5)$ whose \mathbb{F}_5 -cohomology is an exterior algebra on classes of grading 7, 15, 23, 31, and 39, and which is built from spheres of these dimensions by fibrations. By [40], there is a product decomposition

$$(SU(20)/SU(15))_5 \simeq S^{31} \times S^{33} \times S^{35} \times S^{37} \times S^{39}.$$

Let $B(7, 15, 23, 39)$ denote the fiber of the composite

$$B_3^5(5) \rightarrow SU(20)_5 \rightarrow (SU(20)/SU(15))_5 \xrightarrow{\rho} (S^{31})_5.$$

Theorem 4.15 (Harper). *There is a homotopy equivalence*

$$(X_{29})_5 \simeq B(7, 15, 23, 39).$$

Note that this result requires more than Theorem 3.7 because the ranks of these H -spaces are not less than $p - 1$. We will provide Harper's proof of this result in Section 6. Here we just remark that our work above is required in the proof, for the entry in position (4, 3) of the matrix of Theorem 4.8 implies that the 39-cell of $(X_{29})_5$ is attached to the 23-cell by α_2 , which is not detected by primary Steenrod operations. This information is required in order to compare the two spaces in Theorem 4.15.

We can determine the Adams operations and v_1 -periodic homotopy groups of $(X_{31})_5$ by an argument very similar to that used above for $(X_{29})_5$. We shall merely sketch. The analogue of Theorem 4.1 is

Theorem 4.16. *There is an isomorphism $H^*(BX_{31}; \hat{\mathbb{Z}}_5) \approx H^*(BT; \hat{\mathbb{Z}}_5)^{G_{31}}$, where G_{31} has the four generators given for G_{29} in the proof of Theorem 4.1 and also $\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$. Then $H^*(BT; \hat{\mathbb{Z}}_5)^{G_{31}}$ is a polynomial ring on the generators f_8, f_{12} , and f_{20} given in Theorem 4.1 together with*

$$\begin{aligned} f_{24} = & m_{(24)} - 66m_{(20,4)} + 1023m_{(16,8)} + 2180m_{(12,12)} + 1\,293\,156m_{(8,8,4,4)} \\ & + 267\,096m_{(12,4,4,4)} + 2\,121\,984m_{(6,6,6,6)} + 620\,352m_{(10,6,6,2)} - 4032m_{(14,6,2,2)} \\ & - 190\,080m_{(10,10,2,2)} - 11\,892m_{(12,8,4)} - 4938m_{(16,4,4)} - 24\,534m_{(8,8,8)} - 2304m_{(18,2,2,2)}. \end{aligned}$$

The analogue of Theorem 4.7 is

Theorem 4.17. Let $f_8, f_{12}, f_{20}, f_{24}$ be as in Theorem 4.16, and let

$$F_j = f_j(\ell_0(x_1), \dots, \ell_0(x_4)).$$

Then the following series are 5-integral through grading 24.

$$\begin{aligned} F_8 - \frac{8}{5}F_{12} - \frac{7}{25}F_8^2 - \frac{4}{25}F_{20} - \frac{21}{125}F_8F_{12} - \frac{99}{125}F_{24} - \frac{597}{625}F_8^3 - \frac{558}{625}F_{12}^2, \\ F_{12} - \frac{2}{5}F_8^2 - \frac{1}{5}F_{20} - \frac{4}{25}F_8F_{12} - \frac{18}{25}F_{24} - \frac{74}{125}F_8^3 - \frac{11}{125}F_{12}^2, \\ F_{20} - \frac{3}{5}F_{24} - \frac{2}{5}F_8^3. \end{aligned}$$

The analogue of Theorem 4.8 is

Theorem 4.18. The Adams operation ψ^k in $K^1(X_{31}; \hat{\mathbb{Z}}_5)$ on the generators z_7, z_{11}, z_{19} , and z_{23} is given by the matrix

$$\begin{pmatrix} k^7 & 0 & 0 & 0 \\ \frac{8}{5}k^7 - \frac{8}{5}k^{11} & k^{11} & 0 & 0 \\ \frac{12}{25}k^7 - \frac{8}{25}k^{11} - \frac{4}{25}k^{19} & \frac{1}{5}k^{11} - \frac{1}{5}k^{19} & k^{19} & 0 \\ \frac{279}{125}k^7 - \frac{168}{125}k^{11} - \frac{12}{125}k^{19} - \frac{99}{125}k^{23} & \frac{21}{25}k^{11} - \frac{3}{25}k^{19} - \frac{18}{25}k^{23} & \frac{3}{5}k^{19} - \frac{3}{5}k^{23} & k^{23} \end{pmatrix}.$$

The analogue of Theorem 4.9 is

Theorem 4.19. The groups $v_1^{-1}\pi_*(X_{31})_{(5)}$ are given by

$$v_1^{-1}\pi_{2t-1}(X_{31}) \approx v_1^{-1}\pi_{2t}(X_{31}) \approx \begin{cases} 0, & t \not\equiv 3 \pmod{4}, \\ \mathbb{Z}/5^4, & t \equiv 15 \pmod{20}, \\ \mathbb{Z}/5^{\min(7, 4+\nu_5(t-7))}, & t \equiv 7 \pmod{20}, \\ \mathbb{Z}/5^{\min(11, 4+\nu_5(t-11))}, & t \equiv 11 \pmod{20}, \\ \mathbb{Z}/5^{\min(19, 4+\nu_5(t-19))}, & t \equiv 19 \pmod{20}, \\ \mathbb{Z}/5^{\min(23, 4+\nu_5(t-23))}, & t \equiv 23 \pmod{20}. \end{cases}$$

The author thanks the referee for astutely noticing a mistake in an earlier version of Theorem 4.19.

In [39, Proposition 2.3], it is proved that the exceptional Lie group E_8 , localized at 5, admits a product decomposition as $X_0(E_8) \times X_2(E_8)$ with $H^*(X_0(E_8); \mathbb{F}_5) \approx \Lambda(x_{15}, x_{23}, x_{39}, x_{47})$. In Section 6, we will prove the following result, which was pointed out by John Harper.

Theorem 4.20 (Harper). There is a homotopy equivalence

$$(X_{31})_5 \simeq X_0(E_8).$$

Note that Theorem 4.19 agrees with the calculation of $v_1^{-1}\pi_*(E_8)_{(5)}$ in [17, 1.1].

5. The 7-primary modular case

In this section, we first give in Theorem 5.1 new explicit formulas for the six polynomials which generate as a polynomial algebra the invariant ring of the complex reflection group G_{34} of [36], called the Mitchell group in [14]. Over $\hat{\mathbb{Z}}_7$, the invariant ring of G_{34} is also a polynomial algebra, but the generators must be altered slightly from the complex case, as we show prior to Theorem 5.6. Next we use this information to find explicit generators for $K^*(BX_{34}; \hat{\mathbb{Z}}_7)$ in Theorem 5.6, and from this the Adams operations in $QK^1(X_{34}; \hat{\mathbb{Z}}_7)$ in Theorem 5.15. These in turn enable us to compute the v_1 -periodic homotopy groups $v_1^{-1}\pi_*(X_{34})_{(7)}$. Finally, we prove in Theorem 5.17 that $(X_{34})_7$ has the homotopy type of a space formed from $SU(42)$. This result was conjectured by the author and proved by John Harper.

Theorem 5.1. The complex invariants of the reflection group G_{34} (defined in the proof) form a polynomial algebra

$$\mathbb{C}[x_1, \dots, x_6]^{G_{34}} \approx \mathbb{C}[f_6, f_{12}, f_{18}, f_{24}, f_{30}, f_{42}]$$

with generators given by

$$f_{6k} = (1 + (-1)^k 27^{k-1} \cdot 5)m_{(6k)} + \sum_{s=1}^k \binom{6k}{3s} (1 + (-1)^{k+s} 27^{k-1})m_{(6k-3s, 3s)} + \sum_{\mathbf{e}} (\mathbf{e})m_{\mathbf{e}},$$

where \mathbf{e} ranges over all partitions $\mathbf{e} = (e_1, \dots, e_r)$ of $6k$ with $3 \leq r \leq 6$ satisfying $e_i \equiv e_j \pmod{3}$ for all i, j , and $e_i \equiv 0 \pmod{3}$ if $r < 6$. Here also (\mathbf{e}) denotes the multinomial coefficient $(e_1 + \dots + e_r)! / (e_1! \dots e_r!)$, and $m_{\mathbf{e}}$ the monomial symmetric polynomial, which is the shortest symmetric polynomial in x_1, \dots, x_6 containing $x_1^{e_1} \dots x_r^{e_r}$.

For example, we have

- $f_6 = -4m_{(6)} + 40m_{(3,3)} + 720m_{(1,1,1,1,1,1)}$;
- $f_{12} = 136m_{(12)} - 26\binom{12}{3}m_{(9,3)} + 28\binom{12}{6}m_{(6,6)} + \sum(\mathbf{e})m_{\mathbf{e}}$, where \mathbf{e} ranges over $\{(6, 3, 3), (3, 3, 3, 3), (2, 2, 2, 2, 2), (7, 1, 1, 1, 1, 1), (4, 4, 1, 1, 1, 1)\}$;
- $f_{18} = (1 - 5 \cdot 27^2)m_{(18)} + \binom{18}{3}(1 + 27^2)m_{(15,3)} + \binom{18}{6}(1 - 27^2)m_{(12,6)} + \binom{18}{9}(1 + 27^2)m_{(9,9)} + \sum(\mathbf{e})m_{\mathbf{e}}$, where \mathbf{e} ranges over $\{(12, 3, 3), (9, 6, 3), (9, 3, 3, 3), (6, 6, 6), (6, 6, 3, 3), (6, 3, 3, 3, 3), (3, 3, 3, 3, 3, 3), (13, 1, 1, 1, 1, 1), (10, 4, 1, 1, 1, 1), (7, 4, 4, 1, 1, 1), (4, 4, 4, 4, 1, 1), (7, 7, 1, 1, 1, 1), (8, 2, 2, 2, 2, 2), (5, 5, 2, 2, 2, 2)\}$.

Proof of Theorem 5.1. As described in [36], the reflection group G_{34} is generated by reflections across the following hyperplanes in \mathbb{C}^6 : $x_i - x_j = 0$, $x_1 - \omega x_2 = 0$, and $x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 0$. Here $\omega = e^{2\pi i/3}$. It follows easily that G_{34} is generated by all permutation matrices together with the following two:

$$\begin{pmatrix} 0 & \omega^2 & 0 & 0 & 0 & 0 \\ \omega & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad I - \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}. \tag{5.2}$$

In [14], Conway and Sloane consider G_{34} instead as the automorphisms of a certain $\mathbb{Z}[\omega]$ -lattice in \mathbb{C}^6 . The lattice has 756 vectors of norm 2. There are none of smaller positive norm. 270 of these vectors are those with ω^a in one position, $-\omega^b$ in another, and 0 in the rest. Here, of course, a and b can be 0, 1, or 2. The other 486 are those of the form $\pm \frac{1}{\sqrt{-3}}(\omega^{a_1}, \dots, \omega^{a_6})$ such that $\sum a_i \equiv 0 \pmod{3}$.

To verify that this lattice approach to G_{34} is consistent with the reflection approach, one can check that the reflection matrices permute these 756 vectors. It is obvious that permutation matrices do, and easily verified for the first matrix of (5.2). The second matrix of (5.2), which has order 2, sends

- $(\omega, -\omega^2, 0, 0, 0, 0)$ to $\frac{1}{\sqrt{-3}}(\omega^2, \omega, 1, 1, 1, 1)$;
- $\frac{1}{\sqrt{-3}}(1, 1, 1, 1, 1, 1)$ to $-\frac{1}{\sqrt{-3}}(1, 1, 1, 1, 1, 1)$;
- $\frac{1}{\sqrt{-3}}(1, 1, 1, \omega, \omega, \omega)$ to $-\frac{1}{\sqrt{-3}}(\omega, \omega, \omega, 1, 1, 1)$;
- $\frac{1}{\sqrt{-3}}(1, 1, \omega, \omega, \omega^2, \omega^2)$ to itself;
- $(1, -1, 0, 0, 0, 0)$ to itself.

After permutation, negation, and multiplication by ω , this takes care of all cases.

Let

$$p_m(x_1, \dots, x_6) = \sum_{(v_1, \dots, v_6)} (v_1 x_1 + \dots + v_6 x_6)^m, \tag{5.3}$$

where the sum is taken over the 756 vectors described above. Then p_m is invariant under G_{34} for every positive integer m . It is proved in [14, Theorem 10] that the ring of complex invariant polynomials is given by

$$\mathbb{C}[x_1, \dots, x_6]^{G_{34}} = \mathbb{C}[p_6, p_{12}, p_{18}, p_{24}, p_{30}, p_{42}]. \tag{5.4}$$

In [14], several other lattices isomorphic to the above one are described, any of which can be used to give a different set of vectors v and invariant polynomials p_m , still satisfying (5.4). The one that we have selected seems to give the simplest polynomials; in particular, the only ones with integer coefficients.

We have $p_{6k} = S_1 + S_2$, where $S_1 = \sum_{i \neq j} \sum_{a,b=0}^2 (\omega^a x_i - \omega^b x_j)^{6k}$, with $1 \leq i, j \leq 6$, and

$$S_2 = \frac{2}{(-3)^{3k}} \sum_{a_i=0}^2 (\omega^{a_1} x_1 + \dots + \omega^{a_5} x_5 + \omega^{-a_1 - \dots - a_5} x_6)^{6k}.$$

The coefficient of 2 on S_2 is due to the ± 1 . Note that the sum for S_1 has $6 \cdot 5 \cdot 3^2$ terms, while that for S_2 has 3^5 terms. Next note that if a term T^{6k} occurs in either sum, then so does $(\omega T)^{6k}$ and $(\omega^2 T)^{6k}$, and all are equal. Thus we obtain $S_1 = 3 \sum_{i \neq j} \sum_{b=0}^2 (x_i - \omega^b x_j)^{6k}$ and

$$S_2 = 3 \frac{2}{(-3)^{3k}} \sum_{a_2, \dots, a_5=0}^2 (x_1 + \omega^{a_2} x_2 + \dots + \omega^{a_5} x_5 + \omega^{-a_2 - \dots - a_5} x_6)^{6k}.$$

We simplify S_1 further as

$$\begin{aligned} S_1 &= 3 \sum_{\ell=0}^{6k} (-1)^\ell \binom{6k}{\ell} \sum_{i \neq j} x_i^\ell x_j^{6k-\ell} \sum_{b=0}^2 \omega^{b\ell} \\ &= 9 \sum_{s=0}^{2k} (-1)^s \binom{6k}{3s} \sum_{i \neq j} x_i^{3s} x_j^{6k-3s} \\ &= 18 \left(5m_{(6k)} + \sum_{s=1}^k (-1)^s \binom{6k}{3s} m_{(6k-3s, 3s)} \right). \end{aligned}$$

At the first step, we have used that $\sum_{b=0}^2 \omega^{b\ell}$ equals 0 if $\ell \not\equiv 0 \pmod 3$, and equals 3 if $\ell \equiv 0 \pmod 3$. At the second step, we have noted that $\sum_{i \neq j} x_i^{3s} x_j^{6k-3s}$ equals $m_{(6k-3s, 3s)}$ if $s \notin \{0, k, 2k\}$, it equals $2m_{(3k, 3k)}$ if $s = k$, and equals $5m_{(6k)}$ if $s = 0$ or $2k$.

The sum S_2 becomes

$$\begin{aligned} S_2 &= \frac{6}{(-3)^{3k}} \sum_{\mathbf{e}} (\mathbf{e}) \sum_{a_2=0}^2 (\omega^{e_2-e_6})^{a_2} \dots \sum_{a_5=0}^2 (\omega^{e_5-e_6})^{a_5} x_1^{e_1} \dots x_6^{e_6} \\ &= \frac{6}{(-27)^k} \sum_{e_1 \equiv \dots \equiv e_6 \pmod 3} (\mathbf{e}) 3^4 x_1^{e_1} \dots x_6^{e_6}. \end{aligned}$$

Then $(-27)^k(S_1 + S_2)/486$ equals the expression which we have listed for f_{6k} in the statement of the theorem. We have chosen to work with this rather than p_{6k} itself for numerical simplicity. It is important that the omitted coefficient is not a multiple of 7.

For good measure, we show that (5.3) is 0 if $m \not\equiv 0 \pmod 6$. If $m \not\equiv 0 \pmod 3$, then replacing terms T^m by $(\omega T)^m$ leaves the sums like S_1 and S_2 for (5.4) unchanged while, from a different perspective, it multiplies them by ω^m . Thus the sums are 0. If $m \equiv 3 \pmod 6$, the term in S_1 corresponding to $\sum x_i^{3s} x_j^{m-3s}$ occurs with opposite sign to that corresponding to $\sum x_i^{m-3s} x_j^{3s}$, and so $S_1 = 0$. For S_2 , the $(\pm 1)^m$ will cause pairs of terms to cancel. \square

Remark 5.5. The only other place known to the author where explicit formulas for invariant polynomials of G_{34} exist is [28], where they occupy 190 pages of dense text when printed.

As pointed out by Kasper Andersen, $f_{42} - (f_6)^7$ is divisible by 7. This is easily seen by expanding $(f_6)^7 = (\sum (v_1 x_1 + \dots + v_6 x_6)^6)^7$ by the multinomial theorem. The need for this became apparent to Andersen, as the author had thought that the invariant ring of G_{34} over $\hat{\mathbb{Z}}_7$ was $\hat{\mathbb{Z}}_7[f_6, \dots, f_{42}]$, and this would have led to an impossible conclusion for the Adams operations in $QK^1(X_{34}; \hat{\mathbb{Z}}_7)$.

Let $h_{42} = \frac{1}{7}(f_{42} - (f_6)^7)$. Then we have the following result, for which we are grateful to Andersen.

Theorem 5.6. *The invariant ring of G_{34} over $\hat{\mathbb{Z}}_7$ is given by*

$$\hat{\mathbb{Z}}_7[x_1, \dots, x_6]^{G_{34}} = \hat{\mathbb{Z}}_7[f_6, f_{12}, f_{18}, f_{24}, f_{30}, h_{42}].$$

Proof. A Magma program written and run by Andersen showed that each of these asserted generators is indecomposable over $\mathbb{Z}/7$. (This is what failed when f_{42} was used; it equals $(f_6)^7$ over $\mathbb{Z}/7$.) Thus the result follows from (5.4). \square

Since f_{36} is invariant under G_{34} , it follows from (5.4) that it can be decomposed over \mathbb{C} in terms of $f_6, f_{12}, f_{18}, f_{24}$, and f_{30} . The nature of the coefficients in this decomposition was not so clear. It turned out that all coefficients were rational numbers which are 7-adic units. We make this precise in

Theorem 5.7. *f_{36} can be decomposed as*

$$q_1 f_6 f_{30} + q_2 f_{12} f_{24} + q_3 f_{18}^2 + q_4 f_6^2 f_{24} + q_5 f_6 f_{12} f_{18} + q_6 f_{12}^3 + q_7 f_6^3 f_{18} + q_8 f_6^2 f_{12}^2 + q_9 f_6^4 f_{12} + q_{10} f_6^6$$

with

$$q_1 = 944\,610\,925\,401/15\,161\,583\,716,$$

$$q_2 = 733\,671\,261/19\,519\,520,$$

$$q_3 = 243\,068\,633/9\,781\,739,$$

$$\begin{aligned}
 q_4 &= -133\,840\,666\,859\,131\,062\,549/73\,986\,709\,144\,034\,080, \\
 q_5 &= -1\,758\,887\,990\,521\,258\,018\,071\,215\,403/629\,320\,589\,839\,873\,719\,708\,800, \\
 q_6 &= -1\,602\,221\,942\,044\,323/4\,879\,880\,000\,000, \\
 q_7 &= 4\,011\,206\,338\,081\,535\,787\,030\,788\,541/114\,421\,925\,425\,431\,585\,401\,600, \\
 q_8 &= 701\,461\,342\,458\,322\,269\,763\,709\,951\,654\,931/15\,733\,014\,745\,996\,842\,992\,720\,000\,000, \\
 q_9 &= -11\,844\,219\,519\,446\,025\,955\,021\,712\,628\,669/22\,348\,032\,309\,654\,606\,523\,750\,000, \\
 q_{10} &= 26\,589\,469\,730\,264\,682\,368\,719\,198\,549\,833/22\,348\,032\,309\,654\,606\,523\,750\,000.
 \end{aligned}$$

Each of these coefficients q_i is a 7-adic unit; i.e. no numerator or denominator is divisible by 7.

Proof. The ten products, $f_6 f_{30}, \dots, f_6^6$, listed above are the only ones possible. We express each of these products as a combination of monomial symmetric polynomials m_e . We use Magma to do this. The length of $m_{(e_1, \dots, e_r)}$ is defined to be r . We only kept track of components of the products of length ≤ 4 . This meant that we only had to include components of length ≤ 4 of the various f_{6k} being multiplied.

There were 34 m_e 's of length ≤ 4 . These correspond to the partitions of 36 into multiples of 3. (Note that monomials with subscripts $\equiv 1$ or $2 \pmod 3$ only occur for us if the length is 6. Not having to deal with them simplifies our work considerably.) Indeed, there was one of length 1, six of length 2, twelve of length 3, and fifteen of length 4. Magma expressed each monomial such as $f_6 f_{30}$ or f_6^6 as an integer combination of these, plus monomials of greater length. We just ignored in the output all those of greater length. The coefficients in these expressions were typically 12 to 15 digits. We also wrote f_{36} as a combination of monomial symmetric polynomials of length ≤ 4 , ignoring the longer ones. This did not require any fancy software, just the multinomial coefficients from Theorem 5.1.

Now we have a linear system of 34 linear equations with integer coefficients in 10 unknowns. The unknowns are the coefficients q_i in the equation at the beginning of Theorem 5.7, and the equations are the component monomials of length ≤ 4 . Miraculously, there was a unique rational solution, as given in the statement of this theorem.

If it were not for the fact that the Conway–Sloane Theorem 5.4 guarantees that there must be a solution when all monomial components (of length ≤ 6) are considered, then we would have to consider them all, but the fact that we got a unique solution looking at only the monomial components of length ≤ 4 implies that this solution will continue to hold in the other unexamined components. \square

Next we wish to modify the generators in Theorem 5.6 to obtain generators of $QK^1(X_{34}; \hat{\mathbb{Z}}_7)$. Similarly to Theorem 4.7, we let $\ell_0(x) = \ln(1+x)$, and

$$F_i = F_i(x_1, \dots, x_6) = f_i(\ell_0(x_1), \dots, \ell_0(x_6)). \tag{5.8}$$

A major calculation is required to modify the classes F_i so that their coefficients are in $\hat{\mathbb{Z}}_7$; i.e. they do not have 7's in the denominators. As observed after (4.4), it will be enough to accomplish this through grading 42 (with grading of x_i considered to be 1).

Theorem 5.9. The following expressions are 7-integral through grading 42:

- $F_{30} + \frac{5}{7}F_{36} + \frac{22}{7^2}F_{42}$;
- $F_{24} + \frac{4}{7}F_{30} + \frac{45}{7^2}F_{36} + \frac{104}{7^3}F_{42}$;
- $F_{18} + \frac{3}{7}F_{24} + \frac{20}{7^2}F_{30} + \frac{157}{7^3}F_{36} + \frac{526}{7^4}F_{42}$;
- $F_{12} + \frac{2}{7}F_{18} + \frac{45}{7^2}F_{24} + \frac{109}{7^3}F_{30} + \frac{1391}{7^4}F_{36} + \frac{6201}{7^5}F_{42}$;
- $F_6 + \frac{1}{7}F_{12} + \frac{22}{7^2}F_{18} + \frac{204}{7^3}F_{24} + \frac{1107}{7^4}F_{30} + \frac{9682}{7^5}F_{36} + \frac{100682}{7^6}F_{42}$.

It was very surprising that just linear terms were needed here. Decomposable terms were certainly expected. The analogue for G_{29} in Theorem 4.7 involved many decomposables. It would be interesting to know why Theorem 5.9 works with just linear terms; presumably this pattern will continue into higher gradings.

Proof of Theorem 5.9. Similarly to the proof of Theorem 4.7, we define

$$\tilde{F}_i = \tilde{F}_i(x_1, \dots, x_6) = f_i(\ell_p(x_1), \dots, \ell_p(x_6)),$$

and observe that a polynomial in the \tilde{F}_i 's is 7-integral if and only if the same polynomial in the F_i 's is.

Next note that in the range of concern for Theorem 5.9 $\ell_7(x) = x + x^7/7$. If we define

$$h_i = h_i(x_1, \dots, x_6) = f_i(x_1 + x_1^7, \dots, x_6 + x_6^7),$$

then Theorem 5.9 is clearly equivalent to

Statement 5.10. For $t \geq 1$ and grading ≤ 42 ,

- $h_{30} + 5h_{36} + 22h_{42} \equiv 0 \pmod{7^t}$ in grading $30 + 6t$;
- $h_{24} + 4h_{30} + 45h_{36} + 104h_{42} \equiv 0 \pmod{7^t}$ in grading $24 + 6t$;
- $h_{18} + 3h_{24} + 20h_{30} + 157h_{36} + 526h_{42} \equiv 0 \pmod{7^t}$ in grading $18 + 6t$;
- $h_{12} + 2h_{18} + 45h_{24} + 109h_{30} + 1391h_{36} + 6201h_{42} \equiv 0 \pmod{7^t}$ in grading $12 + 6t$;
- $h_6 + h_{12} + 22h_{18} + 204h_{24} + 1107h_{30} + 9682h_{36} + 100682h_{42} \equiv 0 \pmod{7^t}$ in grading $6 + 6t$.

We use Maple to verify Statement 5.10. Our f_i 's are given in Theorem 5.1 in terms of m_e 's. To evaluate $m_e(x_1 + x_1^7, \dots, x_6 + x_6^7)$, the following result keeps the calculation manageable (e.g. it does not involve a sum over all permutations). Partitions can be written either in increasing order or decreasing order; we use increasing. If (a_1, \dots, a_r) is an r -tuple of positive integers, let $s(a_1, \dots, a_r)$ denote the sorted form of the tuple; i.e. the rearranged version of the tuple so as to be in increasing order. For example, $s(4, 2, 3, 2) = (2, 2, 3, 4)$.

Proposition 5.11. The component of $m_{(e_1, \dots, e_r)}(x_1 + x_1^7, \dots, x_6 + x_6^7)$ in grading $\sum e_i + 6t$ is

$$\sum_{\mathbf{j}} \frac{P(e_1 + 6j_1, \dots, e_r + 6j_r)}{P(e_1, \dots, e_r)} \binom{e_1}{j_1} \cdots \binom{e_r}{j_r} m_{s(e_1+6j_1, \dots, e_r+6j_r)},$$

where $\mathbf{j} = (j_1, \dots, j_r)$ ranges over all r -tuples of nonnegative integers summing to t , and $P(a_1, \dots, a_r)$ is the product of the factorials of repetend sizes.

For example, $P(4, 2, 3, 3) = 2!$ because there are two 3's, $P(3, 1, 3, 3, 1, 2) = 3!2!$, and $P(3, 4, 2, 1) = 1$.

Example 5.12. We consider as a typical example, the component of

$$m_{(3,3,9,15)}(x_1 + x_1^7, \dots, x_6 + x_6^7)$$

in grading 42. Table 5.13 lists the possible values of \mathbf{j} and the contribution to the sum. The final answer is the sum of everything in the right-hand column.

Proof of Proposition 5.11. $m_{(e_1, \dots, e_r)}(x_1 + x_1^7, \dots, x_6 + x_6^7)$ is related to

$$\sum_{\sigma} \left(x_{\sigma(1)}^{e_1} + \binom{e_1}{1} x_{\sigma(1)}^{e_1+6} + \binom{e_1}{2} x_{\sigma(1)}^{e_1+12} + \cdots \right) \cdots \left(x_{\sigma(r)}^{e_r} + \binom{e_r}{1} x_{\sigma(r)}^{e_r+6} + \cdots \right) \tag{5.14}$$

summed over all permutations σ in Σ_r . If t values of e_i are equal, then (5.14) will give $t!$ times the correct answer. That is the reason that we divide by $P(\mathbf{e})$. If $(e_1 + 6j_1, \dots, e_r + 6j_r)$ contains s equal numbers, then the associated m will be obtained from each of $s!$ permutations, which is the reason that $P(e_1 + 6j_1, \dots, e_r + 6j_r)$ appears in the numerator. \square

We continue now with the proof of Theorem 5.9. At first, mimicking Theorem 4.7, we were allowing for products of h 's in addition to the linear terms which appear in Statement 5.10, but it was turning out that what was needed to satisfy the congruences was just the linear term. If just a linear term was going to work, the coefficients could be obtained by just looking at monomials of length 1. They were computed by Maple, using that, by Theorem 5.1 and Proposition 5.11, the coefficient of $m_{(6k+6t)}$ in h_{6k} is $(1 + (-1)^k 27^{k-1} \cdot 5) \binom{6k}{t}$. Write the k th expression from the bottom of Statement 5.10 as $\sum_{j \geq 0} a_{j,k} h_{6k+6j}$. We require that the coefficient of $m_{(6k+6t)}$ in $\sum_{j=0}^t a_{j,k} h_{6k+6j}$ is $0 \pmod{7^t}$. But this coefficient equals

$$\sum_{j=0}^t a_{j,k} \binom{6k+6j}{t-j} (1 + (-1)^{k+j} 27^{k+j-1} \cdot 5).$$

Table 5.13
Terms for Example 5.12

\mathbf{j}	Term
(2, 0, 0, 0)	$\binom{3}{2} m_{3,9,15,15}$
(0, 2, 0, 0)	$\binom{3}{2} m_{3,9,15,15}$
(0, 0, 2, 0)	$\binom{9}{2} m_{3,3,15,21}$
(0, 0, 0, 2)	$\binom{15}{2} m_{3,3,9,27}$
(1, 1, 0, 0)	$3 \cdot 3 \cdot 3 m_{9,9,9,15}$
(1, 0, 1, 0)	$3 \cdot 9 m_{3,9,15,15}$
(1, 0, 0, 1)	$3 \cdot 15 m_{3,9,9,21}$
(0, 1, 1, 0)	$3 \cdot 9 m_{3,9,15,15}$
(0, 1, 0, 1)	$3 \cdot 15 m_{3,9,9,21}$
(0, 0, 1, 1)	$9 \cdot 15 m_{3,3,15,21}$

We solve iteratively for $a_{j,k}$, starting with $a_{0,k} = 1$, and obtain the values in Statement 5.10. Note that it first gives $a_{1,1} \equiv 1 \pmod{7}$. If we had chosen a value such as 8 or -6 instead of 1, then the value of $a_{2,1}$ would be different than 22. So these numbers $a_{j,k}$ are not uniquely determined. These different choices just amount to choosing a different basis for $QK^1(X_{34}; \hat{\mathbb{Z}}_7)$.

Verifying Statement 5.10 required running many Maple programs. For each line of Statement 5.10, a verification had to be made for each relevant t -value, from two t -values for the first line down to six t -values for the last line. Moreover, for each of these pairs (line number, t -value), it was convenient to use a separate program for monomials of each length 2, 3, 4, and 5, and then, for monomials of length 6, it was done separately for those with subscripts congruent to 0, 1, or 2 mod 3. Thus altogether $(2 + 3 + 4 + 5 + 6)(4 + 3) = 140$ Maple programs were run. The programs had enough similarity that one could be morphed into another quite easily, and a more skillful programmer could incorporate them all into the same program.

Note that expanding from f_j to h_j does not change the number of components in monomials, nor does it change the mod 3 value of the sum of the subscripts (i.e. exponents) in the monomials. This is simpler than the situation in the proof of Theorem 5.7. The algorithm is quite easy. For each combination of h 's in Statement 5.10, replace each h_{6j} by the combination of m_e 's in f_{6j} in Theorem 5.1, but expanded using Proposition 5.11. \square

To obtain the Adams operations in $QK^1(X_{34}; \hat{\mathbb{Z}}_7)$, we argue similarly to the paragraph which precedes Theorem 4.8. First note that F_{36} decomposes in terms of F_{6i} 's exactly as does f_{36} in terms of f_{6i} 's in Theorem 5.7. We can⁸ modify by decomposables in dimensions greater than 42 to obtain 7-integral classes $G_6, G_{12}, G_{18}, G_{24}$, and G_{30} which agree with the classes of Theorem 5.9 (with F_{36} decomposed) through dimension 42. There is also a 7-integral class G_{42} which agrees with $\frac{1}{7}(F_{42} - (F_6)^7)$ in dimension 42. These generate $K^*(BX_{34}; \hat{\mathbb{Z}}_7)$ as a power series algebra. As in the preamble to Theorem 4.8, then $z_i := B^{-1}e^*(G_{i+1})$ for $i = 5, 11, 17, 23, 29$, and 41 form a basis for $QK^1(X_{34}; \hat{\mathbb{Z}}_7)$, and e^* annihilates decomposables.

Similarly to the situation for $(X_{29})_5$ in the proof of Theorem 4.8, if we let

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{7} & 1 & 0 & 0 & 0 & 0 \\ \frac{22}{49} & \frac{2}{7} & 1 & 0 & 0 & 0 \\ \frac{204}{343} & \frac{45}{49} & \frac{3}{7} & 1 & 0 & 0 \\ \frac{1107}{2401} & \frac{109}{343} & \frac{20}{49} & \frac{4}{7} & 1 & 0 \\ \frac{16647}{16807} & \frac{1399}{2401} & \frac{183}{343} & \frac{6}{49} & \frac{1}{7} & 1 \end{pmatrix},$$

then the matrix of ψ^k on the basis $\{z_5, z_{11}, z_{17}, z_{23}, z_{29}, z_{41}\}$ is

$$P^{-1} \text{diag}(k^5, k^{11}, k^{17}, k^{23}, k^{29}, k^{41}) P.$$

The entries in the last row of P are 7 times the coefficients of F_{42} in Theorem 5.9 reduced mod 1. Those coefficients were multiplied by 7 because z_{41} is related to $\frac{1}{7}F_{42}$ rather than to F_{42} .

Using this, we compute the ν_1 -periodic homotopy groups, similarly to Theorem 4.9. Note the remarkable similarity with that result. Here, of course, $\nu(-)$ denotes the exponent of 7 in an integer.

Theorem 5.15. *The groups $\nu_1^{-1}\pi_*(X_{34})_{(7)}$ are given by*

$$\nu_1^{-1}\pi_{2t-1}(X_{34}) \approx \nu_1^{-1}\pi_{2t}(X_{34}) \approx \begin{cases} 0, & t \not\equiv 5 \pmod{6}, \\ \mathbb{Z}/7^5, & t \equiv 5, 35 \pmod{42}, \\ \mathbb{Z}/7^{\min(12, 5+\nu(t-11-12\cdot 7^6))}, & t \equiv 11 \pmod{42}, \\ \mathbb{Z}/7^{\min(18, 5+\nu(t-17-18\cdot 7^{12}))}, & t \equiv 17 \pmod{42}, \\ \mathbb{Z}/7^{\min(24, 5+\nu(t-23-18\cdot 7^{18}))}, & t \equiv 23 \pmod{42}, \\ \mathbb{Z}/7^{\min(30, 5+\nu(t-29-12\cdot 7^{24}))}, & t \equiv 29 \pmod{42}, \\ \mathbb{Z}/7^{\min(42, 5+\nu(t-41-24\cdot 7^{36}))}, & t \equiv 41 \pmod{42}. \end{cases}$$

Proof. The group $\nu_1^{-1}\pi_{2t}(X)_{(7)}$ is presented by $\left(\begin{smallmatrix} (\psi^7)^T \\ (\psi^3)^T - 3^t I \end{smallmatrix}\right)$, since 3 generates $\mathbb{Z}/49^\times$. We let $x = 3^t$ and form this matrix analogously to (4.10). Five times we can pivot on units, removing their rows and columns, leaving a column matrix with 7 polynomials in x . The 7-exponent of $\nu_1^{-1}\pi_{2t}(X)_{(7)}$ is the smallest of that of these polynomials (with $x = 3^t$). This will be 0 unless $x \equiv 5 \pmod{7}$, which is equivalent to $t \equiv 5 \pmod{6}$. We find that two of these polynomials will always yield, between them, the smallest exponent. Similarly to (4.11) and Table 4.12, we write these polynomials as $p_i(3^m + y)$ for carefully-chosen values of m . Much preliminary work is required to discover these values of m . Ignoring unit coefficients and ignoring higher-power terms whose coefficients will be sufficiently divisible that they will not affect the divisibility, these polynomials will be as in Table 5.16.

⁸ But we need not bother to do so explicitly.

Table 5.16
Certain $p_i(3^m + y)$ (linear part only)

m	p_1	p_2
5, 35	$7^5 + 7^4 y$	$7^{\geq 12} + 7^{11} y$
$11 + 12 \cdot 7^6$	$7^{12} + 7^4 y$	$7^{12} + 7^{11} y$
$17 + 18 \cdot 7^{12}$	$7^{18} + 7^4 y$	$7^{18} + 7^{11} y$
$23 + 18 \cdot 7^{18}$	$7^{25} + 7^4 y$	$7^{24} + 7^{11} y$
$29 + 12 \cdot 7^{24}$	$7^{31} + 7^4 y$	$7^{30} + 7^{11} y$
$41 + 24 \cdot 7^{36}$	$7^{43} + 7^4 y$	$7^{42} + 7^{11} y$

The claim of the theorem follows from Table 5.16 by the same argument as was used in the proof of Theorem 4.9. For t in the specified congruence, if $3^t = 3^m + y$, then $v(y) = v(t - m) + 1 \geq 2$, similarly to (4.13). For example, if $t \equiv 11 \pmod{42}$, and $3^t = 3^{11+12 \cdot 7^6} + y$, then $v(y) = v(t - 11 - 12 \cdot 7^6) + 1$. Thus $\min(v(p_1(3^t)), v(p_2(3^t)))$ will be determined by the 7^5 in p_1 if $t \equiv 5, 35 \pmod{42}$, while in the other cases, it is determined by the $7^4 y$ in p_1 or the constant term in p_2 .

The groups $v_1^{-1}\pi_{2t-1}(X_{34})$ are cyclic by an argument similar to the one described at the end of the proof of Theorem 4.9, and have the same order as $v_1^{-1}\pi_{2t}(X_{34})$ for the standard reason described there. \square

Similarly to the discussion preceding Theorem 4.15, one of the factors in the product decomposition of $SU(42)_7$ given in [31] is an H -space $B_5^7(7)$ whose \mathbb{F}_7 -cohomology is an exterior algebra on classes of grading 11, 23, 35, 47, 59, 71, and 83, and which is built from spheres of these dimensions by fibrations. Using [40], we can obtain a degree-1 map $B_5^7(7) \rightarrow S^{71}$. Let $B_7 := B(11, 23, 35, 47, 59, 83)$ denote its fiber. The following result was conjectured by the author and proved by John Harper. Its proof will be described in the last line of the paper.

Theorem 5.17 (Harper). *There is a homotopy equivalence $(X_{34})_7 \simeq B_7$.*

6. Proofs provided by John Harper

In this section, we provide proofs of Theorems 4.15, 4.20, and 5.17, which were explained to the author by John Harper. We begin with the following strengthening of Theorem 3.7.

Theorem 6.1. *Theorem 3.7 is true for $r \leq p - 1$.*

Proof. Although the proof presented in [15] works for $r = p - 1$ just as it does for $r < p - 1$, we take this opportunity to explain some aspects of it more thoroughly. For $\lambda \neq 0 \in \mathbb{Z}/p$, a π_λ -space (resp. Q_λ -space) is one which admits a self-map inducing multiplication by λ in $\pi_*(-) \otimes \mathbb{Z}/p$ (resp. $QH^*(-; \mathbb{Z}/p)$). A π_λ -map (resp. Q_λ -map) is a map between spaces of the indicated type which commutes up to homotopy with the self-maps.

The first part of the proof, extending [15, Theorem 1.3], involves showing that if X is an H -space of rank $r \leq p - 1$ and with torsion-free homology, then there is a π_λ -map $X \rightarrow S^n$ whose homotopy fiber Y is an H -space of rank $r - 1$ with torsion-free homology. The proof of this is exactly as in [15, p. 358], noting that [41, 4.2.2] is valid (and stated) for $r = p - 1$ as well as for $r < p - 1$. That Y is an H -space follows from [15, Theorem 1.1] since its rank is less than $p - 1$. This construction can be iterated to yield (3.8).

We use Lemma 6.2 to deduce the part of Theorem 6.1 which says that the homotopy type of X is determined by certain homotopy classes. Suppose we have (3.8) and a primed version. Suppose we have shown that X_i and X'_i are p -equivalent H -spaces and that the elements $\alpha \in \pi_{n_{i+1}-1}(X_i)$ and $\alpha' \in \pi_{n_{i+1}-1}(X'_i)$ correspond under this equivalence. Then, in the notation of Lemma 6.2, there are equivalences as π_λ -spaces

$$X_{i+1} \simeq (X_i)^\alpha \simeq (X'_i)^{\alpha'} \simeq X'_{i+1}.$$

If $i + 1 < p - 1$, then by [15, Theorem 1.1], there are p -equivalent H -space structures on X_{i+1} and X'_{i+1} , extending the induction. Note that for X_{p-1} and X'_{p-1} we do not assert an equivalence as H -spaces, only as π_λ -spaces. \square

Lemma 6.2. *Suppose $Y \rightarrow X \xrightarrow{g} S^n$ is a fibration, with X a π_λ -space and Y an H -space with $\text{rank}(Y) < p - 1$. Let $\alpha \in \pi_{n-1}(Y)$ denote $\partial(\iota_n)$ in the homotopy sequence of the fibration. As in [15, p. 351], let $Y^\alpha = D_+^n \times Y \cup_c D_-^n \times Y$, where $c(x, y) = (x, \alpha(x)y)$, using the H -space multiplication of Y . Then there is a homotopy equivalence $X \simeq Y^\alpha$ which is a π_λ -map.*

Proof. We expand on some aspects of the proof given in [15, p. 358] and correct several confusing typos. As explained there, Y^α is a π_λ -space and admits an inclusion $Y \cup_\alpha e^n \hookrightarrow Y^\alpha$.

On the other hand, the given fibration is fiber homotopy equivalent to $Y \rightarrow X' \rightarrow S^n$, with $X' = D_+^n \times Y \cup_\gamma D_-^n \times Y$, where $\gamma: S^{n-1} \times Y \rightarrow S^{n-1} \times Y$ is the clutching function defined by the homotopy equivalences $D_\pm^n \times Y \rightarrow g^{-1}(D_\pm^n)$. We will work with X' , but in the end may replace it with X . Note that X' inherits the π_λ -structure of X .

Since $\gamma|S^{n-1} \times \{*\}$ represents α , the relative n -skeleta $(Y^\alpha, Y)^n$ and $(X', Y)^n$ both equal $Y \cup_\alpha e^n$. Thus the n th stages of the Postnikov systems of Y^α and X' are built from the same Eilenberg–MacLane spaces and same k -invariants, and so there exists a π_λ -map $(Y^\alpha)_n \rightarrow (X')_n$ of these n th stages, inducing a cohomology isomorphism in dimension $\leq n$. We wish to show that this extends to a π_λ -map $Y^\alpha \rightarrow X'$, which will then be a homotopy equivalence by the Five Lemma applied to the homotopy sequences of the fibrations $Y \rightarrow Y^\alpha \rightarrow S^n$ and $Y \rightarrow X' \rightarrow S^n$, and Whitehead's theorem.

This requires that we consider the primitive Postnikov systems (PPS) of the map $X' \rightarrow (X')_n$. The *primitive Postnikov system*, as described, for example, in [27, p. 426ff], applies to a rational equivalence, and gives a Postnikov tower in which all fibers are mod p Eilenberg–MacLane spaces. Our map $X' \rightarrow (X')_n$ is a rational equivalence because the only infinite homotopy groups of the spheres that build X' are present in $(X')_n$. By [15, Lemma 2.8], the maps in the PPS are π_λ -maps. We will also use that, by [15, Lemma 2.4], all our π_λ -maps are also Q_λ -maps.

We will show that the composite $Y^\alpha \rightarrow (Y^\alpha)_n \rightarrow (X')_n$ lifts through the PPS of $X' \rightarrow (X')_n$. Assume there is a lifting to a π_λ -map $Y^\alpha \rightarrow E_s$, where E_s is some stage in the PPS. Since the k -invariants in the PPS are λ -eigenvectors by [15, Lemma 2.8], so are their images in $H^*(Y^\alpha; \mathbb{F}_p)$. A basis for $H^*(Y^\alpha; \mathbb{F}_p)$ consists of products of no more than $p - 1$ generators, each of which is a λ -eigenvector for the Q_λ -map of Y^α . An element in this basis which is not one of the generators is a λ^i -eigenvector for some $2 \leq i \leq p - 1$. Since, for such i , $\lambda^i \not\equiv \lambda \pmod p$, we deduce that a nonzero image of a k -invariant can only equal one of the generators⁹ of $H^*(Y^\alpha; \mathbb{F}_p)$. However, the k -invariants are in dimension greater than that of the generators of $H^*(Y^\alpha; \mathbb{F}_p)$. We conclude that the image of the k -invariants must equal 0, and so the map lifts to $Y^\alpha \rightarrow E_{s+1}$. By [15, 2.7], this lifting may be chosen to be a π_λ -map. Since the PPS has only finitely many stages through $\dim(Y^\alpha)$, we obtain the desired lifting $Y^\alpha \rightarrow X'$. \square

We now apply these general results to our specific situations.

Proof of Theorem 4.20. Both spaces $(X_{31})_5$ and $X_0(E_8)$ are H -spaces with mod-5 cohomology an exterior algebra on classes of dimension 15, 23, 39, and 47. By Theorem 6.1, there exist diagrams of fibrations

$$\begin{array}{ccccccc} S^{15} & \longrightarrow & X_2 & \longrightarrow & X_3 & \longrightarrow & (X_{31})_5 \\ & & f_2 \downarrow & & f_3 \downarrow & & f_4 \downarrow \\ & & S^{23} & & S^{39} & & S^{47} \end{array}$$

and

$$\begin{array}{ccccccc} S^{15} & \longrightarrow & X'_2 & \longrightarrow & X'_3 & \longrightarrow & X_0(E_8) \\ & & f'_2 \downarrow & & f'_3 \downarrow & & f'_4 \downarrow \\ & & S^{23} & & S^{39} & & S^{47} \end{array}$$

and the homotopy types of the spaces are determined by relevant elements of homotopy groups.

Always localized at 5, we have $\pi_{22}(S^{15}) \approx \mathbb{Z}/5$ generated by α_1 . The elements in $\pi_{22}(S^{15})$ which determine both X_2 and X'_2 are nonzero multiples of α_1 , and so by Theorem 3.7 we obtain an equivalence $X'_2 \simeq X_2$. The claim about these homotopy classes is proven for $(X_{31})_5$ from the entry in position (2, 1) in the Adams operation matrix in Theorem 4.18; Adams' e -invariant says that α_1 attaching maps are present iff the relevant Adams operation involves $u(k^n - k^{n+p-1})/p$ with u a unit. Similarly for $X_0(E_8)$, the presence of α_1 can be deduced from the Adams operations in E_8 as given in [17, Proposition 3.5], although here the well-known Steenrod algebra action in $H^*(E_8)$ also implies the attaching map. We now identify X'_2 with X_2 in our notation.

The homomorphism $\pi_{38}(X_2) \rightarrow \pi_{38}(S^{23})$ is a surjection $\mathbb{Z}/25 \rightarrow \mathbb{Z}/5$. See, e.g., Diagram 3.5. The $\frac{1}{5}(k^{11} - k^{19})$ in position (3, 2) of Theorem 4.18 and in the formula for $\psi^k(\chi_{11})$ in [17, Proposition 3.5] tell that in both of our sequences, the determining element of $\pi_{38}(X_2)$ is a generator, i.e., it maps to $\alpha_2 \in \pi_{38}(S^{23})$, and so, by Theorem 3.7, there is a homotopy equivalence $X_3 \simeq X'_3$. This is probably the only place that any of our calculations with $(X_{31})_5$ are required in proving Theorem 4.20. The α_1 attaching maps can be seen by Steenrod operations, but α_2 requires secondary operations, which were used in [23] to see the α_2 attaching map in $X_0(E_8)$, or Adams operations in K -theory, as we have done.

We identify X_3 and X'_3 . The homomorphism $\pi_{46}(X_3) \rightarrow \pi_{46}(S^{39})$ is a surjection $\mathbb{Z}/5^3 \rightarrow \mathbb{Z}/5$. (The cyclicity of $\pi_{46}(X_3)$ can be proved easily using [17, Proposition 5.5].) As in the previous two paragraphs, Adams operations imply that our determining element in both sequences is a generator, i.e., it maps to $\alpha_1 \in \pi_{46}(S^{39})$, and so we obtain the asserted equivalence from Theorem 6.1. \square

Proof of Theorem 4.15. We first show that the map $B_3^5 \rightarrow S^{31}$ described prior to Theorem 4.15 can be chosen to be a π_λ -map. By [15, Lemma 2.8], the generating map $S^{31} \rightarrow Z(\mathbb{Z}, 31)$ has a PPT in which all maps are π_λ -maps. The map

⁹ Or a linear combination of generators in the same dimension.

$B_3^5 \xrightarrow{g} K(\mathbb{Z}, 31)$ is a π_λ -map, since $\pi_{31}(B_3^5)$ is cyclic. By [40], g lifts to a map $B_3^5 \rightarrow S^{31}$. By the Lifting Theorem [15, 2.7a], this lifting, through each stage of the PPT, is a π_λ -map.

Thus, by [15, Lemma 2.3], the space which we call $B(7, 15, 23, 39)$ in Theorem 4.15 is a π_λ -space. Let f denote the composite $B(7, 15, 23, 39) \rightarrow B_3^5 \rightarrow SU(20) \rightarrow S^{39}$. Its fiber, $B_3^3 = B(7, 15, 23)$, is one of the four factors in $SU(12)_5$, and hence is an H -space.

On the other hand, Theorem 6.1 yields fibrations of H -spaces, localized at 5,

$$\begin{array}{ccccccc}
 S^7 & \longrightarrow & X_2 & \longrightarrow & X_3 & \longrightarrow & (X_{29})_5 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & S^{15} & & S^{23} & & S^{39}.
 \end{array} \tag{6.3}$$

We can almost apply Theorem 6.1 to obtain the desired equivalence of $(X_{29})_5$ and $B(7, 15, 23, 39)$. The only thing missing is that we do not know that $B(7, 15, 23, 39)$ is an H -space.

Instead, we first use Theorem 3.7 to show that X_3 (of (6.3)) and B_3^3 are homotopy equivalent H -spaces. Note that in the proof of Theorem 6.1 we obtained equivalent H -structures as long as the rank was less than $p - 1$. In this case, the homotopy classes that must correspond are, in both cases, $\alpha_1 \in \pi_{14}(S^7)_{(5)} \approx \mathbb{Z}/5$ and a generator of $\pi_{22}(B(7, 15)) \approx \mathbb{Z}/25$, mapping to $\alpha_1 \in \pi_{22}(S^{15})$. The α_1 's in B_3^3 are well known by the action of \mathcal{P}^1 , while in X_3 they may be deduced from entries in the matrix of Theorem 4.8 in positions (2, 1) and (3, 2).

For the fibrations $X_3 \rightarrow (X_{29})_5 \rightarrow S^{39}$ and $B_3^3 \rightarrow B(7, 15, 23, 39) \rightarrow S^{39}$, the relevant homotopy groups, always localized at 5, $\pi_{38}(X_3)$ and $\pi_{38}(B_3^3)$, are both $\mathbb{Z}/5^3$, and the relevant classes are generators, which we will call α_2 since they each map to $\alpha_2 \in \pi_{38}(S^{23})$. This is seen for X_3 from the entry in position (4, 3) of the matrix of Theorem 4.8, while for B_3^3 it can be obtained from the Adams operation calculation in [35, pp. 667–668]. Thus, by Lemma 6.2, we have homotopy equivalences

$$(X_{29})_5 \simeq (X_3)^{\alpha_2} \simeq (B_3^3)^{\alpha_2} \simeq B(7, 15, 23, 39),$$

as desired. \square

The proof of Theorem 5.17 is entirely analogous, and is omitted.

Acknowledgements

The author is grateful to Kasper Andersen, John Harper, Hirokazu Nishinobu, Clarence Wilkerson, and the referee for valuable suggestions related to this project, each of which is specifically described in the paper.

References

- [1] J. Aguadé, Constructing modular classifying spaces, *Israel J. Math.* 66 (1989) 23–40.
- [2] K.K.S. Andersen, The normalizer splitting conjecture for p -compact groups, *Fund. Math.* 161 (1999) 1–16.
- [3] K.K.S. Andersen, T. Bauer, J. Grodal, E.K. Pedersen, A finite loop space not rationally equivalent to a compact Lie group, *Invent. Math.* 157 (2004) 1–10.
- [4] K.K.S. Andersen, J. Grodal, The classification of 2-compact groups, Preprint, math.AT/0611437.
- [5] K.K.S. Andersen, J. Grodal, J.M. Møller, A. Viruel, The classification of p -compact groups for p odd, *Ann. of Math.* (2) 167 (2008) 95–210.
- [6] M. Bendersky, D.M. Davis, v_1 -periodic homotopy groups of the Dwyer–Wilkerson space, Preprint, arXiv:0706.0993.
- [7] M. Bendersky, D.M. Davis, 2-primary v_1 -periodic homotopy groups of $SU(n)$, *Amer. J. Math.* 114 (1991) 465–494.
- [8] M. Bendersky, D.M. Davis, M. Mimura, v_1 -periodic homotopy groups of exceptional Lie groups—torsion-free cases, *Trans. Amer. Math. Soc.* 333 (1992) 115–136.
- [9] A.K. Bousfield, The K -theory localizations and v_1 -periodic homotopy groups of H -spaces, *Topology* 38 (1999) 1239–1264.
- [10] A.K. Bousfield, E. Curtis, D. Kan, D. Quillen, D. Rector, J. Schlesinger, The mod p lower central series and the Adams spectral sequence, *Topology* 5 (1966) 331–342.
- [11] C. Broto, J.M. Møller, Chevalley p -local finite groups, *Algebr. Geom. Topol.* 7 (2007) 1809–1919.
- [12] N. Castellana, Quillen Grassmannians as non-modular homotopy fixed points, *Math. Z.* 248 (2004) 477–493.
- [13] A. Clark, J. Ewing, The realization of polynomial algebras as cohomology rings, *Pacific J. Math.* 50 (1974) 425–434.
- [14] J.H. Conway, N.J.A. Sloane, The Coxeter–Todd lattice, the Mitchell group, and related sphere packings, *Math. Proc. Cambridge Philos. Soc.* 93 (1983) 421–440.
- [15] G. Cooke, J.R. Harper, A. Zabrodsky, Torsion-free mod p H -spaces of low rank, *Topology* 18 (1979) 349–359.
- [16] D.M. Davis, Representation types and 2-primary homotopy groups of certain compact Lie groups, *Homology, Homotopy Appl.* 5 (2003) 297–324.
- [17] D.M. Davis, From representation theory to homotopy groups, *Mem. Amer. Math. Soc.* 759 (2002).
- [18] D.M. Davis, v_1 -periodic homotopy groups of $SU(n)$ at odd primes, *Proc. London Math. Soc.* (3) 43 (1991) 529–544.
- [19] D.M. Davis, Computing v_1 -periodic homotopy groups of spheres and certain Lie groups, in: *Handbook of Algebraic Topology*, Elsevier, 1995, pp. 993–1049.
- [20] D.M. Davis, M. Mahowald, Some remarks of v_1 -periodic homotopy groups, *London Math. Soc. Lecture Note Ser.* 176 (1992) 55–72.
- [21] D.M. Davis, H. Yang, Tractable formulas for v_1 -periodic homotopy groups of $SU(n)$ when $n \leq p^2 - p + 1$, *Forum Math.* 8 (1996) 585–620.
- [22] W.G. Dwyer, C.W. Wilkerson, Homotopy fixed-point methods for Lie groups and finite loop spaces, *Ann. of Math.* (2) 139 (1994) 395–442.
- [23] D.L. Gonçalves, The mod-5 splitting of the compact exceptional Lie group E_8 , *Publ. Res. Inst. Math. Sci.* 19 (1983) 1–6.
- [24] H. Hasse, Die Gruppe der p^n -primären Zahlen für ein Primteiler von p , *J. Reine Angew. Math.* 176 (1936) 174–183.
- [25] Y. Hemmi, H. Nishinobu, K. Ogi, The action of the Steenrod algebra on the cohomology of p -compact groups, Preprint.

- [26] A. Jeanneret, A. Osse, The K -theory of p -compact groups, *Comment. Math. Helv.* 72 (1997) 556–581.
- [27] R.M. Kane, *The Homology of Hopf Spaces*, North-Holland, 1988.
- [28] G. Kemper, et al., A database of invariant rings, www.maths.abdn.ac.uk/~bensond/html/archive/kemper-koerding-malle-matzat-vogel-wiese.html.
- [29] R. Levi, On finite groups and homotopy theory, *Mem. Amer. Math. Soc.* 567 (1995).
- [30] H. Maschke, Ueber die quarternäre, endliche, lineare Substitutionsgruppe der Borchardt'schen Moduln, *Math. Ann.* 30 (1887) 496–515.
- [31] M. Mimura, G. Nishida, H. Toda, Mod p decomposition of compact Lie groups, *Publ. Res. Inst. Math. Sci.* 13 (1977) 627–680.
- [32] M. Mimura, H. Toda, Cohomology operations and the homotopy of compact Lie groups, I, *Topology* 9 (1970) 317–336.
- [33] H. Nishinobu, On the action of the Steenrod algebra on the cohomology of a certain p -compact group, *Kochi J. Math.* 2 (2007) 125–153.
- [34] D. Notbohm, Spaces with polynomial mod- p cohomology, *Math. Proc. Cambridge Philos. Soc.* 126 (1999) 277–292.
- [35] K. Potocka, The number of summands in v_1 -periodic homotopy groups of $SU(n)$, *Topology Appl.* 153 (2005) 647–677.
- [36] G.C. Shephard, J.A. Todd, Finite unitary reflection groups, *Canad. J. Math.* 6 (1954) 274–304.
- [37] D.P. Sullivan, *Geometric Topology, the 1970 MIT notes*, Springer-Verlag, 2003.
- [38] H. Toda, *Composition Methods in the Homotopy Groups of Spheres*, *Ann. of Math. Stud.*, vol. 49, Princeton University Press, Princeton, 1962.
- [39] C. Wilkerson, Self-maps of classifying spaces, in: *Lecture Notes in Math.*, vol. 418, Springer-Verlag, 1974, pp. 150–157.
- [40] A. Yamaguchi, On the p -regularity of Stiefel manifolds, *Publ. Res. Inst. Math. Sci.* 25 (1989) 355–380.
- [41] A. Zabrodsky, Endomorphisms in the homotopy category, in: *Contemp. Math.*, vol. 44, Amer. Math. Soc., 1985, pp. 227–277.