

Holding Your Own: Establishing Sister Chromatid Cohesion

Robert V. Skibbens

Dept. of Biological Sciences, Lehigh University, Bethlehem, Pennsylvania 18015, USA

Chromosome Segregation and Genome Integrity

At cell division, both daughter cells receive only one copy of the parental genome. Not surprisingly, chromatin plays a role at numerous stages to ensure that genome segregation occurs with high fidelity. First, each chromosome is duplicated to produce two identical chromosomes called sister chromatids. Second, sister chromatids become paired, or glued together, along the entire chromosome length (Sumner 1991; Selig et al. 1992; Guacci et al. 1994). This pairing, or cohesion, provides a mechanism by which the identity of sister chromatids can be maintained over time. Third, sister pairs associate with the mitotic spindle apparatus via kinetochores, specialized protein complexes that assemble on centromeres. This association produces a highly defined geometry: The kinetochore of one chromatid associates with microtubules from the spindle pole opposite that of her sister. Cohesion between sister chromatids facilitates this orientation, most probably by physically constraining sister kinetochores to face in opposite directions (Rieder 1982; Waters et al. 1996; Michaelis et al. 1997; Tanaka et al. 2000a). Fourth, kinetochores associated with spindle microtubules generate poleward-pulling forces that act to separate sister chromatids (McNeill and Berns 1981; Rieder et al. 1986; Skibbens et al. 1995). Cohesion opposes these kinetochore pulling forces, stretching the intervening chromatin and generating tension across the chromatid pair. This tension is critical for coordinating sister chromatid movement during mitosis (Skibbens et al. 1995), and for cell cycle progression. In support of the latter, kinetochores capture microtubules by chance, such that initial attachments are not always correct (Nicklas 1997). Fortunately, cells have evolved a mitotic checkpoint mechanism that halts cell cycle progression in response to inappropriate kinetochore-microtubule interactions or in the absence of tension (Hoyt et al. 1991; Li and Murray 1991; Nicklas et al. 1995; Li and Nicklas 1997; Waters et al. 1998, 1999; Skibbens et al. 1999). Only after all of the chromatid pairs are properly oriented to the mitotic spindle apparatus does dissolution of cohesion occur, allowing sister chromatids to segregate away from each

other and move into the newly forming daughter cells (Fig. 1).

Numerous advances have facilitated the molecular dissection of *cis* and *trans* factors required for establishing sister chromatid cohesion. A seminal finding that propelled forward our understanding of cohesion and cell cycle regulation was that anaphase onset (sister separation) and exit from mitosis (M-phase to interphase transition) are separable events: The dissolution of sister chromatid cohesion could now be studied in an isolated context (Holloway et al. 1993; Surana et al. 1993; Irniger et al. 1995). Other advancements, including the adaptations of fluorescent in situ hybridization (FISH) and GFP-tagged repressor/operator methodologies, allowed for the visualization of individual sister chromatid loci in yeast cells, an organism in which mutants are readily generated and the genome sequence is known (Guacci et al. 1994; Straight et al. 1996; Michaelis et al. 1997; Megee and Koshland 1999). Finally, analyses of DNA from coimmunoprecipitates, in combination with functional assays of cohesion in minichromosomes, have provided a vehicle to identify DNA sequences that participate in cohesion (Dedon et al. 1991; Meluh and Koshland 1997; Blat and Kleckner 1999; Megee and Koshland 1999; Megee et al. 1999; Tanaka et al. 1999).

In this review, we take a *cerevisiae*-centric view of the recent advances that have brought insights, as well as intrigue, to the mechanism by which *cis* and *trans* factors participate in cohesion establishment.

Structural Cohesins

In budding yeast, several structural cohesion proteins (or cohesins) have been identified: Smc1p, Smc3p, Mcd1p/Scs1p, and Irr1p/Scs3p (Strunnikov et al. 1993; Kurlandzka et al. 1995; Guacci et al. 1997; Michaelis et al. 1997; Toth et al. 1999). These cohesins are conserved across evolution and their characterizations have been summarized in numerous reviews (Hirano 1999; Nasmyth 1999; Strunnikov and Jessberger 1999). Briefly, cohesin proteins are thought to function as structural components in sister chromatid cohesion based on three criteria. First, each protein is required for cohesion such that loss of function in any one results in sister chromatid separation before anaphase onset (Guacci et al. 1997; Michaelis et al. 1997; Toth et al. 1999). All loci separate in cohesin mutant cells, in-

E-MAIL rvs3@lehigh.edu; **FAX (610) 758-4004.**

Article and publication are at www.genome.org/cgi/doi/10.1101/gr.153600.




Figure 1 Cell cycle in terms of chromosome dynamics.

dicating that the mechanism of cohesion is conserved along the chromosome length. Note that factors unrelated to cohesins have been identified from other organisms that localize to, or function in, cohesion only at centromeres (Saitoh et al. 1997; Moore and Orr-Weaver 1998; Tang et al. 1998). Second, cohesins are nuclear proteins that associate with chromatin from S-phase through the bulk of mitosis: That portion of the cell cycle when sister chromatid cohesion must be maintained (Guacci et al. 1997; Michaelis et al. 1997; Toth et al. 1999; Ciosk et al. 2000). Data from yeast and *Xenopus* support a model in which all four cohesins function as a complex (Guacci et al. 1997; Michaelis et al. 1997; Losada et al. 1998; Toth et al. 1999). Third, cohesins dissociate from chromatin at anaphase onset when sister chromatids separate (Guacci et al. 1997; Michaelis et al. 1997; Toth et al. 1999; Uhlmann et al. 1999), although the kinetics of Smc1p and Smc3p dissociation are delayed relative to those of Mcd1p/Scs1p and Irr1p/Scs3p (Tanaka et al. 1999; Ciosk et al. 2000). In summary, these results are consistent with a model in which a complex required for cohesion associates with chromatin during the period of cohesion maintenance.

Cohesin dissociation from chromatin has been interpreted as a biochemical marker for anaphase onset and sister separation. For Mcd1p/Scs1p, dissociation is ultimately regulated by the anaphase-promoting complex (APC). APC, in association with the APC cofactor Cdc20p, targets Pds1p (an inhibitor of anaphase that exists in a complex with Esp1p) for degradation (Cohen-Fix et al. 1996; Schwab et al. 1997; Visintin et al. 1997; Ciosk et al. 1998). Pds1p degradation frees Esp1p to function as a positive regulator of anaphase onset.

In yeast, Esp1p facilitates the two-site cleavage and dissociation of the cohesin Mcd1p/Scs1p from chromatids, possibly via an endopeptidase activity (Ciosk et al. 1998; Uhlmann et al. 1999; Nasmyth et al. 2000). In contrast, little is known concerning the mechanism and extent of Smc1p and Smc3p dissociation. Several reviews provide various details concerning anaphase onset and the dissolution of cohesion (Farr and Cohen-Fix 1999; Nasmyth 1999; Peters 1999).

Sites of Cohesion

With cohesion proteins in hand, researchers focused on identifying the DNA sequences that participate in cohesion. Such an analysis is daunting, not only because of the genome size of most organisms, but because it was unknown whether cohesins associated with (1) all DNA sequences, (2) specific sequences repeated throughout the genome, or (3) specific sites but in a sequence-independent manner. Numerous approaches were used to address this issue. Chromatin immunoprecipitation (CHIP) tests for cohesin deposition at both endogenous chromosomes and at ectopic DNA sequences inserted within yeast chromosomes or minichromosomes (Blat and Kleckner 1999; Megee et al. 1999; Tanaka et al. 1999). While CHIP is a powerful approach and has been used to identify cohesion sequences at relatively high resolution (200–300 bp), this method relies on signals obtained from a population of cells. In addition, cohesin subunits participate in different complexes to perform a variety of tasks (e.g., DNA recombination/repair, chromosome condensation, and gene expression), requiring the performance of other assays showing that cohesin deposition participates functionally in cohesion (Jessberger et al. 1996; Guacci et al. 1997; Donze et al. 1999). To produce an overview of cohesion deposition along an entire chromosome, the CHIP method was modified such that radio-labeled DNA obtained from chromatin coimmunoprecipitations was used to probe membrane filters that contained DNA fragments spanning chromosome III (Blat and Kleckner 1999). Thus, the sites and spacing of cohesion deposition were obtained for a whole chromosome, albeit with reduced resolution. In addition to CHIP-based methodologies, Megee and Koshland (1999) exploited the small size of yeast plasmids to minimize cohesion binding-site redundancy and induce recombination events to regulate *in vivo* excision of highly defined DNA sequences from plasmids. Sister chromatid dissociation of the resulting recombination products could then be directly monitored using FISH and GFP-tagged minichromosomes, thus providing a

functional assay for cohesion on a per cell basis (Megee and Koshland 1999; Megee et al. 1999). In close agreement, the centromere was identified as a predominant cohesin-binding site (Blat and Kleckner 1999; Megee et al. 1999; Tanaka et al. 1999), consistent with cytological studies from higher eukaryotes in which sister chromatid centromeres are tightly paired and often not individually resolvable (Sumner 1991).

What portion of the centromere is required for cohesion? The yeast centromere is composed of three centromere DNA elements (CDE I, CDE II, and CDE III) that together comprise approximately 125 bp (Hyman and Sorger 1995). While the full-length centromere functions in cohesion and is a substrate for Mcd1p/Sccl1p deposition, a single base pair substitution in CDE III significantly decreased Mcd1p/Sccl1p deposition and cohesion between. CDE III allowed for only very limited cohesion binding and did not function as a cohesion site until a portion of CDE II was included. These results reveal that cohesion at the centromere is directed by as little as 100 bp that predominantly requires CDE III (a portion of CDE II followed by all of CDE III) (Megee and Koshland 1999; Megee et al. 1999; Tanaka et al. 1999). CDE III is also essential for chromosome segregation and centromere structures associated with kinetochore assembly (Hyman and Sorger 1995). Does cohesion rely on kinetochore function as opposed to centromere sequence? Tanaka and coworkers (1999) found that mutations in kinetochore proteins, including Ndc10p, Cse4p, and Mif2p, abolish Mcd1p/Sccl1p deposition at ectopic centromeres (deposition at endogenous centromeres was only minimally affected). These results suggest that Mcd1p/Sccl1p is recruited to, and participates in, cohesion at yeast centromeres but that this recruitment relies on kinetochore function, not centromere sequence.

In addition to the centromere, numerous chromosome arm loci have been identified as sites of cohesin deposition or cohesion (Blat and Kleckner 1999; Megee and Koshland 1999; Megee et al. 1999; Tanaka et al. 1999). Using a functional assay for cohesion, inclusion of *URA3* was found to partially compensate for reduced minichromosome pairing upon centromere excision (Megee and Koshland 1999). *URA3* and *LEU2* were also found to function as sites of Mcd1p/Sccl1p deposition at their endogenous contexts (chromosomes V and III, respectively) (Megee and Koshland 1999; Megee et al. 1999). Other deposition sites for chromosome V include a region encompassing *DMC1* and another just downstream from *YER184C* (Tanaka et al. 1999). A site of cohesin deposition also has been identified proximal to the 5S gene within each rDNA repeat on chromosome XII (S. Laloraya and D. Koshland, pers. comm.). Although all of these sites have been resolved to within a few hundred base pairs, a motif that directs cohesin binding has not been identified, nor has a cor-

relation of deposition to intergenic or intragenic loci been obtained. To date, the best correlation between cohesin deposition and DNA sequence resides in nucleotide bias: Chromatin regions that bind cohesins are high in A/T content, including the CDE II region of the centromere. Regions that are high in G/C content appear to demarcate cohesin association, possibly acting as a barrier to cohesin spreading (Blat and Kleckner 1999; Megee et al. 1999; Tanaka et al. 1999).

Given that cohesion between sister chromatids must be maintained through a significant portion of the cell cycle, it is surprising to find evidence that the cohesin-chromatin interaction is quite dynamic. First, ectopic insertion of a centromere into a site previously devoid of detectable Mcd1p/Sccl1p levels results in deposition not only at the centromere, but also into flanking cold sequences (Megee et al. 1999; Tanaka et al. 1999). Cohesin spreading over time may help explain the robust and expansive signal observed for centromeres relative to arm sites (Blat and Kleckner 1999; Megee et al. 1999; Tanaka et al. 1999). Arm sites may also be competent to direct the spread of cohesin: When a region of high Mcd1p/Sccl1p deposition was divided into eight short DNA sequences, only two of these sequences allowed for deposition when integrated to a new locus (Tanaka et al. 1999). Note that the inability of the remaining six segments to direct binding could instead relate to an inhibitory effect emanating from the flanking sequences (see below). Second, cohesin spreading occurs symmetric to flanking sequences but can also spread in an asymmetric fashion, indicating that flanking sequences can facilitate or inhibit cohesin spreading (Megee et al. 1999). Third, cohesins that spread to sequences adjacent to the centromere dissociate from those regions upon excision of the centromere, even when the centromere is excised as late as metaphase (Megee et al. 1999). Thus, cohesin's association with DNA must be actively maintained (at least at sites of cohesin spreading). Finally, cohesion must accommodate chromosome condensation. Cytological studies reveal that human chromosomes in late prophase appear as cylinders with unresolved sister chromatids, suggesting that a significant volume of sister chromatids has cohesion. As condensation proceeds, the individual sisters become more defined until, late in metaphase, each chromatid is highly resolved and paired along a narrowed axial region (Sumner 1991). This axis of cohesion does not simply appear to be the condensed sum of all the previous cohesion sites, because cells maintained in metaphase eventually lose cohesion along the arms. These results suggest that the majority of cohesion sites are lost before metaphase, consistent with studies performed in *Xenopus* and *Sordaria* (Losada et al. 1998; van Heemst et al. 1999). Insight as to how the dynamic process of cohesion and condensation may coexist was

obtained by the finding that cohesion factors Mcd1p/Sccl1p, Ctf7p/Eco1p, and *CHL12/CTF18* are required for aspects of chromosome condensation (Guacci et al. 1997; Skibbens et al. 1999) (J. Hanna et al., pers. comm.). Further study is required to determine the extent to which cohesion can be remodeled in response to DNA replication/repair, condensation, transcription, or other chromatin-altering processes.

Establishing Cohesion

The mechanism by which the products of DNA replication are identified as sisters remains elusive. Three factors, Ctf7p/Eco1p, Sccl2p, and Sccl4p, have been identified as acting early in the process of cohesion (Skibbens et al. 1999; Toth et al. 1999; Ciosk et al. 2000). Sccl2p and Sccl4p appear to function in cohesin deposition. First, Sccl2p (Mis4p in *S. pombe*) (Furuya et al. 1998) and Sccl4p combine to form a complex and function during S-phase when cohesion is established. Second, the structural cohesins, Mcd1p/Sccl1p and Irr1p/Sccl3p, fail to stably associate with chromatin in *scc2* or *scc4* mutant cells (Toth et al. 1999; Ciosk et al. 2000), consistent with a deposition function for the Sccl2p-Sccl4p complex. The molecular mechanism by which this deposition occurs is unknown. For instance, while cohesin and deposition (Sccl2p and Sccl4p) complexes associate with chromatin, they infrequently colocalize and do not stably associate with one another, at least not in stoichiometric measures (Toth et al. 1999; Ciosk et al. 2000).

One model put forward in the literature posits that structural cohesins are preassembled into a soluble complex and then deposited onto the chromatin (Tanaka et al. 1999; Toth et al. 1999; Ciosk et al. 2000). For instance, all four cohesins physically associate to form a soluble complex and have been shown to coassociate with at least one discrete chromosomal locus (Tanaka et al. 1999; Toth et al. 1999). Further evidence for the requirement of preassembly before deposition is that Sccl3p fails to associate with chromatin in cells harboring mutations in either *MCD1/SCC1*, *SMC1*, or *SMC3* (or the deposition factor *SCC2*) while Mcd1p/Sccl1p fails to associate with chromatin in cells harboring mutations in *IRR1/SCC3* (or *SCC2*). A chromatin association for Mcd1p/Sccl1 was not detected in *SMC1* or *SMC3* mutant cells (Toth et al. 1999). These findings have been interpreted to show an interdependence of cohesin subunit association with chromatin: Deposition occurs in the form of an intact cohesin complex.

Formally, an alternate model is that cohesin deposition occurs by the stepwise addition of cohesins. It has been well established that Mcd1p/Sccl1p is upregulated and begins to associate with chromatin in late G1/early S-phase (Guacci et al. 1997; Michaelis et al. 1997; Toth et al. 1999). However, Smc1p associates with chromatin before Mcd1p/Sccl1p is upregulated

and before Mcd1p/Sccl1p or Irr1p/Sccl3p associate with chromatin (Ciosk et al. 2000). That SMC protein association with chromatin may be the first step in cohesin deposition makes sense: Only SMC family members (shown for various constructs of cohesins Smc1p and Smc3p and condensin Smc2p) have been shown to directly bind DNA. Smc1p and Smc3p C-terminal fragments preferentially bind DNA substrates that are A/T rich or contain secondary structures. Of interest is that yeast centromere DNA, a prominent cohesion site that is A/T rich and contains inverted palindromic repeats, is one of many efficient competitors of Smc1p substrate binding (Akhmedov et al. 1998, 1999). Eukaryotic SMCs most likely assemble into heterodimers (Hirano 1998; Jessberger et al. 1998; Nasmyth 1999). Note that bacterial SMC proteins are singly represented in the genome and thus assemble into homodimers (Melby et al. 1998). Thus, a likely scenario for eukaryotes is that an Smc1p-Smc3p complex associates with DNA, and that this deposition allows for the subsequent loading of Mcd1p/Sccl1p and Irr1p/Sccl3p. In *Drosophila*, the cohesion factor, MEI-S332, associates with chromatin after S-phase, providing evidence across evolutionary lines that cohesion between sister chromatids may occur in a multistep fashion (Kerrebrock et al. 1992; Tang et al. 1998).

Surprisingly, the combined activities of deposition (Sccl2p and Sccl4p) and cohesin (Mcd1p/Sccl1p, Irr1p/Sccl3p, Smc1p, and Smc3p) factors are still insufficient to establish cohesion. For instance, studies in yeast and *Xenopus* reveal that cohesins can bind chromatin (presumably via the Sccl2p-Sccl4p complex) before, during, after, or in the absence of DNA replication. However, when cohesin deposition is excluded from S-phase, sister chromatid cohesion is abolished (Losada et al. 1998; Uhlmann and Nasmyth 1998). Characterization of Ctf7p/Eco1p unveiled a third class of cohesion factors: those that function in cohesion establishment. Like *SCC2* and *SCC4*, *CTF7/ECO1* is required only during S-phase. However, *CTF7/ECO1* mutant cells contain separated sister chromatids even though cohesin proteins associate with chromatin and appear properly targeted to cohesion sites (Skibbens et al. 1999; Toth et al. 1999; R. Skibbens and D. Koshland, unpubl.). Fundamentally, cohesion establishment, as mediated by Ctf7p/Eco1p, must include a mechanism in addition to cohesin assembly and deposition.

The finding that *CTF7/ECO1* interacts genetically with *POL30* and *CHL12/CTF18* provided one of the first molecular clues that the processes of DNA replication and cohesion establishment are intimately coupled (Skibbens et al. 1999). *POL30* encodes the DNA polymerase cofactor proliferating cell nuclear antigen (PCNA) (Bauer and Burgers 1990). PCNA trimers form a tauoidal sliding clamp that associates with and promotes DNA polymerase processivity at the replica-

tion fork (Kelman 1997). PCNA is loaded onto dsDNA by a replication factor complex (RFC) (Cullmann et al. 1995; Jonsson and Hubscher 1997). Attesting to the specificity of the above interactions is that *CHL12/CTF18* encodes an RFC-like protein (Spencer et al. 1990; Kouprina et al. 1993,1994). Because cohesion is established during S-phase (Sumner 1991; Selig et al. 1992; Guacci et al. 1994; Uhlmann and Nasmyth 1998; Skibbens et al. 1999; Toth et al. 1999), the sum of these results suggests a model in which cohesion establishment is directly coupled to DNA replication and takes place in association with PCNA at the replication fork where sister chromatids are closely apposed (Fig. 2). This model is attractive because it helps explain how cohesion establishment ensures the specificity of chromatid pairing while at the same time precludes the catastrophic pairing of nonsisters or repetitive sequences within a chromatid. Mis4p, the fission yeast homolog of budding yeast Scc2p, also interacts with the DNA replication machinery (Furuya et al. 1998), providing support across evolutionary lines that cohesion establishment and DNA replication are coupled.

Cementing the link between DNA replication/repair and cohesion are two recent findings, the first that *TRF4* encodes a DNA polymerase (Pol κ), which functions in sister chromatid cohesion (Wang et al. 2000). Pharmacological studies reveal that Pol κ (*TRF4*) shows sensitivities shared by B-polymerases, a class of DNA replication/repair enzymes (Pelletier et al. 1994; Wang et al. 2000). Whereas *TRF4-TRF5* double mutant cells fail to complete DNA synthesis, *TRF4* mutant cells complete the bulk of S-phase but contain a high frequency of separated sisters. It is not known how the Pol κ polymerase function of *TRF4* facilitates cohesion,

although *TRF4* interactions with *SMC1* have been reported (Castano et al. 1996). The second finding comes from fission yeast. *ESO1*, the fission yeast homolog of *CTF7/ECO1*, has been isolated and is required for sister chromatid cohesion, showing that *CTF7/ECO1* is conserved in function across evolution (Tanaka et al. 2000b). Intriguingly, *ESO1* is much larger than its budding yeast counterpart and appears to be the fusion of budding yeast *RAD30* as its N-terminus and *CTF7/ECO1* as its C-terminus (Skibbens et al. 1999; Toth et al. 1999; Tanaka et al. 2000b). *RAD30* encodes yet another DNA polymerase (Pol η) that functions in DNA repair (Johnson et al. 1999). *RAD30* function in cohesion has yet to be tested. Interestingly, mutations in one of the human homologs of *RAD30* (Pol η) have been identified in patients afflicted with xeroderma pigmentosum, a disease associated with UV-induced oncogenesis (Woodgate 1999).

Good things can come in threes, and characterizations of the three newest cohesion factors (*CTF8*, *CHL12/CTF18*, and *CTF4/CHL15/POB1*) provide exciting insights into the mechanism of cohesion. *CTF8*, a chromosome transmission fidelity mutant, is synthetically lethal with known structural (*MCD1/SCC1* and *SMC3*) deposition (*SCC2*) and establishment (*CTF7/ECO1*) cohesion factors. Given these interactions, it is not surprising that *CTF8* mutant cells are defective in sister chromatid cohesion (M. Mayer, S. Gygi, and P. Hieter, pers. comm.). *CHL12/CTF18*, implicated in cohesion by genetic interactions with *CTF7/ECO1* (Skibbens et al. 1999), also functions in cohesion (J. Hanna, E. Kroll, V. Lundblad, and F. Spencer, pers. comm.). In a remarkable convergence of independent endeavors, numerous methodologies show that *CHL12/CTF18* and *CTF8* interact with each other and with bona fide RFC components (*RFC4* and/or *RFC3*) (J. Hanna, pers. comm.; M. Mayer, S. Gygi, and P. Hieter, pers. comm.). Note that while *CHL12/CTF18* contains a region conserved among RFC factors (and most notably, *RFC1*), *CHL12/CTF18* does not appear to associate with the RFC complex (*RFC1-RFC5*) that functions in processive DNA replication (J. Hanna, pers. comm.). Previous studies showed that RFC components (*RFC2-RFC5*) combined with the DNA damage checkpoint protein *RAD24p*, which contains regions of homology with *RFC1* (Ellison and Stillman 1998; Green et al. 2000). The interactions described above extend this finding and suggest that there may be several such RFC-containing complexes, one of which may scan along DNA in association with the DNA replication fork to function in cohesion.




Figure 2 One possible mode of action for cohesion establishment. Structural cohesion factors Mcd1p/Scs1p, Irr1p/Scs3p, Smc1p, and Smc3p (yellow balls) are loaded as a complex or by stepwise addition onto chromatid via the Scc2p-Scc4p deposition complex (salmon rectangle). Deposition is shown to occur near the site of action of Ctf7p, but may occur anywhere before or after the replication fork. Establishment of sister chromatid cohesion requires Ctf7p/Eco1p (green triangle), which may move in association with PCNA at the DNA replication fork. Ctf8p, Ctf4p/Chl15p/Pob1p, Chl12p/Ctf18p, and the DNA polymerase Trf4p (Pol κ) are other cohesion factors that may associate with the DNA replication fork (not shown).

Whereas *CTF4/CHL15/POB1* has not yet been implicated in interactions with bona fide RFC components, this third-newest cohesion factor is required for cohesion, and genetically interacts with *CHL12/CTF18* and several genes implicated in DNA synthesis, including DNA polymerase α (Miles and Formosa 1992; Wittmeyer and Formosa 1995; Formosa and Nittis 1999; J. Hanna, E. Kroll, V. Lundblad, and F. Spencer, pers. comm.). Of interest is that both *CTF8* and *CHL12/CTF18* interact with telomere metabolism genes and function in telomere length regulation (M. Mayer, S. Gygi, and P. Hieter, pers. comm.; J. Hanna, E. Kroll, V. Lundblad, and F. Spencer, pers. comm.). Can the mechanisms of cohesion be central to so many different processes (telomere metabolism, chromosome condensation, DNA replication), or are the various phenotypes shown by cohesion mutant cells a secondary consequence of cohesion loss?

Whereas the challenges that face researchers interested in sister chromatid cohesion still loom large, the summit is much closer and the trails are clear. Specific milestones that we will likely find along the way include discerning that portion of the cohesion machinery that resides at the DNA replication fork as opposed to that which resides and functions on discrete loci after fork passage. In addition, it will be interesting to determine how mediators of DNA topology (DNA replication, transcription, silencing, etc.) influence cohesion and site selection. Coming full circle, cohesion factors may themselves remodel chromatin to facilitate the site-selection process (Hirano and Mitchison 1994; Saka et al. 1994; Hirano et al. 1997; Sutani and Yanagida 1997; Losada et al. 1998; Kimura et al. 1999). For instance, mutations in numerous cohesion factors have been shown to functionally alter or abolish centromere structures or higher-order assemblies on chromosome arms (Spencer et al. 1990; Castano et al. 1996; Guacci et al. 1997; Saitoh et al. 1997; Goshima et al. 1999; Skibbens et al. 1999; van Heemst et al. 1999; J. Hanna, E. Kroll, V. Lundblad, and F. Spencer, pers. comm.). Finally, if the cohesion machinery scans the DNA to identify sites for cohesin deposition, how are these complexes regulated and for what does it scan?

ACKNOWLEDGMENTS

The author thanks Drs. Meg Kenna, Vincent Guacci, Paul Megee, F. Spencer, M. Mayer, and laboratory members for their critical reading of the manuscript, their time, and helpful discussions. The author also thanks those researchers who kindly shared information before publication: S. Laloraya, V. Guacci, D. Koshland, J. Hanna, E. Kroll, V. Lundblad, F. Spencer, M. Mayer, S. Gygi, and P. Hieter.

REFERENCES

Akhmedov, A.T., Frei, C., Tsai-Pflugfelder, M., Kemper, B., Gasser, S.M., and Jessberger, R. 1998. Structural maintenance of

- chromosomes protein C-terminal domains bind preferentially to DNA with secondary structure. *J. Biol. Chem.* **273**: 24088–24094.
- Akhmedov, A.T., Gross, B., and Jessberger, R. 1999. Mammalian SMC3 C-terminal and coiled-coil protein domains specifically bind palindromic DNA, do not block DNA ends, and prevent DNA bending. *J. Biol. Chem.* **274**: 38216–38224.
- Bauer, G.A. and Burgers, P.M. 1990. Molecular cloning, structure and expression of the yeast proliferating cell nuclear antigen gene. *Nucleic Acids Res.* **18**: 261–265.
- Blat, Y. and Kleckner, N. 1999. Cohesins bind to preferential sites along yeast chromosome III, with differential regulation along arms versus the centric region. *Cell* **98**: 249–259.
- Castano, I.B., Brzoska, P.M., Sadoff, B.U., Chen, H., and Christman, M.F. 1996. Mitotic chromosome condensation in the rDNA requires TRF4 and DNA topoisomerase I in *Saccharomyces cerevisiae*. *Genes & Dev.* **10**: 2564–2576.
- Ciosk, R., Zachariae, W., Michaelis, C., Shevchenko, A., Mann, M., and Nasmyth, K. 1998. An ESP1/PDS1 complex regulates loss of sister chromatid cohesion at the metaphase to anaphase transition in yeast. *Cell* **93**: 1067–1076.
- Ciosk, R., Shirayama, M., Shevchenko, A., Tanaka, T., Toth, A., Shevchenko, A., and Nasmyth, K. 2000. Cohesin's binding to chromosomes depends on a separate complex consisting of Scc2 and Scc4. *Molecular Cell* **5**: 243–254.
- Cohen-Fix, O., Peters, J.M., Kirschner, M.W., and Koshland, D. 1996. Anaphase initiation in *Saccharomyces cerevisiae* is controlled by the APC-dependent degradation of the anaphase inhibitor Pds1p. *Genes & Dev.* **10**: 3081–3093.
- Cullmann, G., Fien, K., Kobayashi, R., and Stillman, B. 1995. Characterization of the five replication factor C genes of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **15**: 4661–4671.
- Dedon, P.C., Soult, J.A., Allis, C.D., and Gorovsky, M.A. 1991. A simplified formaldehyde fixation and immunoprecipitation technique for studying protein-DNA interactions. *Annal. Biochem.* **197**: 83–90.
- Donze, D., Adams, C.R., Rine, J., and Kamakaka, R.T. 1999. The boundaries of the silenced HMR domain in *Saccharomyces cerevisiae*. *Genes & Dev.* **13**: 698–708.
- Ellison, V. and Stillman, B. 1998. Reconstitution of recombinant human replication factor C (RFC) and identification of an RFC subcomplex possessing DNA-dependent ATPase activity. *J. Biol. Chem.* **273**: 5979–5987.
- Farr, K.A. and Cohen-Fix, O. 1999. The metaphase to anaphase transition: A case of productive destruction. *Eur. J. Biochem.* **263**: 14–19.
- Formosa, T. and Nittis, T. 1999. Dna2 mutants reveal interactions with DNA polymerase alpha and Ctf4, a Pol alpha accessory factor, and show that full Dna2 helicase activity is not essential for growth. *Genetics* **151**: 1459–1470.
- Furuya, K., Takahashi, K., and Yanagida, M. 1998. Faithful anaphase is ensured by Mis4, a sister chromatid cohesion molecule required in S phase and not destroyed in G1 phase. *Genes & Dev.* **12**: 3408–3418.
- Goshima, G., Saitoh, S., and Yanagida, M. 1999. Proper metaphase spindle length is determined by centromere proteins Mis12 and Mis6 required for faithful chromosome segregation. *Genes & Dev.* **13**: 1664–1677.
- Green, C.M., Erdjument-Bromage, H., Tempst, P., and Lowndes, N.F. 2000. A novel Rad24 checkpoint protein complex closely related to replication factor C. *Curr. Biol.* **10**: 39–42.
- Guacci, V., Hogan, E., and Koshland, D. 1994. Chromosome condensation and sister chromatid pairing in budding yeast. *J. Cell Biol.* **125**: 517–530.
- Guacci, V., Koshland, D., and Strunnikov, A. 1997. A direct link between sister chromatid cohesion and chromosome condensation revealed through the analysis of MCD1 in *S. cerevisiae*. *Cell* **91**: 47–57.
- Hirano, T. 1998. SMC protein complexes and higher-order chromosome dynamics. *Curr. Opin. Cell Biol.* **10**: 317–322.
- . 1999. SMC-mediated chromosome mechanics: A conserved scheme from bacteria to vertebrates? *Genes & Dev.* **13**: 11–19.

- Hirano, T. and Mitchison, T.J. 1994. A heterodimeric coiled-coil protein required for mitotic chromosome condensation in vitro. *Cell* **79**: 449–458.
- Hirano, T., Kobayashi, R., and Hirano, M. 1997. Condensin, chromosome condensation protein complexes containing XCAP-C, XCAP-E and a Xenopus homolog of the *Drosophila barren* protein. *Cell* **89**: 511–521.
- Holloway, S.L., Glotzer, M., King, R.W., and Murray, A.W. 1993. Anaphase is initiated by proteolysis rather than by the inactivation of maturation-promoting factor. *Cell* **73**: 1393–1402.
- Hoyt, M.A., Totis, L., and Roberts, B.T. 1991. *S. cerevisiae* genes required for cell cycle arrest in response to loss of microtubule function. *Cell* **66**: 507–517.
- Hyman, A.A. and Sorger, P.K. 1995. Structure and function of kinetochores in budding yeast. *Ann. Rev. Cell. Dev. Biol.* **11**: 471–495.
- Irniger, S., Piatti, S., Michaelis, C., and Nasmyth, K. 1995. Genes involved in sister chromatid separation are needed for B-type cyclin proteolysis in budding yeast. *Cell* **81**: 269–278.
- Jessberger, R., Riwar, B., Baechtold, H., and Akhmedov, A.T. 1996. SMC proteins constitute two subunits of the mammalian recombination complex RC-1. *EMBO J.* **15**: 4061–4068.
- Jessberger, R., Frei, C., and Gasser, S.M. 1998. Chromosome dynamics: The SMC protein family. *Curr. Opin. Genet. Dev.* **8**: 254–259.
- Johnson, R.E., Prakash, S., and Prakash, L. 1999. Efficient bypass of a thymine-thymine dimer by yeast DNA polymerase, Polh. *Science* **283**: 1001–1004.
- Jonsson, Z.O. and Hubscher, U. 1997. Proliferating cell nuclear antigen: More than a clamp for DNA polymerases. *BioEssays* **19**: 967–975.
- Kelman, Z. 1997. PCNA: Structure, functions and interactions. *Oncogene* **14**: 629–640.
- Kerrebrock, A.W., Miyazaki, W.Y., Birnby, D., and Orr-Weaver, T.L. 1992. The *Drosophila mei-5332* gene promotes sister-chromatid cohesion in meiosis following kinetochore differentiation. *Genetics* **130**: 827–841.
- Kimura, K., Rybenkov, V.V., Crisona, N.J., Hirano, T., and Cozzarelli, N.R. 1999. 13S condensin actively reconfigures DNA by introducing global positive writhe: Implications for chromosome condensation. *Cell* **98**: 239–248.
- Kouprina, N., Tsouladze, A., Koryabin, M., Hieter, P., Spencer, F., and Larionov, V. 1993. Identification and genetic mapping of CHL genes controlling mitotic chromosome transmission in yeast. *Yeast* **9**: 11–19.
- Kouprina, N., Kroll, E., Kirillov, A., Bannikov, B., Zakharyev, V., and Larionov, V. 1994. CHL12, a gene essential for the fidelity of chromosome transmission in the yeast *Saccharomyces cerevisiae*. *Genetics* **138**: 1067–1079.
- Kurlandzka, A., Rytka, J., Gromadka, R., and Murawski, M. 1995. A new essential gene located on *Saccharomyces cerevisiae* chromosome IX. *Yeast* **11**: 885–890.
- Li, R. and Murray, A.W. 1991. Feedback control of mitosis in budding yeast. *Cell* **66**: 519–531.
- Li, X. and Nicklas, R.B. 1997. Tension-sensitive kinetochore phosphorylation and the chromosome distribution checkpoint in praying mantid spermatocytes. *J. Cell Science* **110**: 537–545.
- Losada, A., Hirano, M., and Hirano, T. 1998. Identification of Xenopus SMC protein complexes required for sister chromatid cohesion. *Genes & Dev.* **12**: 1986–1997.
- McNeill, P.A. and Berns, M.W. 1981. Chromosome behavior after laser microirradiation of a single kinetochore in mitotic PtK2 cells. *J. Cell Biol.* **88**: 543–553.
- Megee, P.C. and Koshland, D. 1999. A functional assay for centromere-associated sister chromatid cohesion. *Science* **285**: 254–257.
- Megee, P.C., Mistrot, C., Guacci, V., and Koshland, D. 1999. The centromeric sister chromatid cohesion site directs Mcd1p binding to adjacent sequences. *Mol. Cell* **4**: 445–450.
- Melby, T.E., Ciampaglio, C.N., Briscoe, G., and Erickson, H.P. 1998. The symmetrical structure of structural maintenance of chromosomes (SMC) and MukB proteins: Long, antiparallel coiled coils, folded at a flexible hinge. *J. Cell Biol.* **142**: 1595–1604.
- Meluh, P.B. and Koshland, D. 1997. Budding yeast centromere composition and assembly as revealed by in vivo cross-linking. *Genes & Dev.* **11**: 3401–3412.
- Michaelis, C., Ciosk, R., and Nasmyth, K. 1997. Cohesins: Chromosomal proteins that prevent premature separation of sister chromatids. *Cell* **91**: 35–45.
- Miles, J. and Formosa, T. 1992. Evidence that POB1, a *Saccharomyces cerevisiae* protein that binds to DNA polymerase alpha, acts in DNA metabolism in vivo. *Mol. Cell. Biol.* **12**: 5724–5735.
- Moore, D.P. and Orr-Weaver, T.L. 1998. Chromosome segregation during meiosis: Building an unambivalent bivalent. *Curr. Top. Dev. Biol.* **37**: 263–299.
- Nasmyth, K. 1999. Separating sister chromatids. *Trends Biochem. Sci.* **24**: 98–104.
- Nasmyth, K., Peters, J.M., and Uhlmann, F. 2000. Splitting the chromosome: Cutting the ties that bind sister chromatids. *Science* **288**: 1379–1385.
- Nicklas, R.B. 1997. How cells get the right chromosomes. *Science* **275**: 632–637.
- Nicklas, R.B., Ward, S.C., and Gorbsky, G.J. 1995. Kinetochore chemistry is sensitive to tension and may link mitotic forces to a cell cycle checkpoint. *J. Cell Biol.* **130**: 929–939.
- Pelletier, H., Sawaya, M.R., Kumar, A., Wilson, S.H., and Kraut, J. 1994. Structures of ternary complexes of rat DNA Polymerase beta, a DNA template-primer, and ddCTP. *Science* **264**: 1891–1903.
- Peters, J.M. 1999. Subunits and substrates of the anaphase-promoting complex. *Exp. Cell Res.* **248**: 339–349.
- Rieder, C.L. 1982. The formation, structure, and composition of the mammalian kinetochore and kinetochore fiber. *Int. Rev. Cytol.* **79**: 1–58.
- Rieder, C.L., Davison, E.A., Jensen, L.C., Cassimeris, L., and Salmon, E.D. 1986. Oscillatory movements of monooriented chromosomes and their position relative to the spindle pole result from the ejection properties of the aster and half-spindle. *J. Cell Biol.* **103**: 581–591.
- Saitoh, S., Takahashi, K., and Yanagida, M. 1997. Mis6, a fission yeast inner centromere protein, acts during G1/S and forms specialized chromatin required for equal segregation. *Cell* **90**: 131–143.
- Saka, Y., Sutani, T., Yamashita, Y., Saitoh, S., Takeuchi, M., Nakaseko, Y., and Yanagida, M. 1994. Fission yeast cut3 and cut14, members of a ubiquitous protein family, are required for chromosome condensation and segregation in mitosis. *EMBO J.* **13**: 4938–4952.
- Schwab, M., Lutum, A.S., and Seufert, W. 1997. Yeast Hct1 is a regulator of Clb2 cyclin proteolysis. *Cell* **90**: 683–693.
- Selig, S., Okumura, K., Ward, D.C., and Cedar, H. 1992. Delineation of DNA replication time zones by fluorescence in situ hybridization. *EMBO J.* **11**: 1217–1225.
- Skibbens, R.V., Rieder, C.L., and Salmon, E.D. 1995. Kinetochore motility after severing between sister centromeres using laser microsurgery: evidence that kinetochore directional instability and position is regulated by tension. *J. Cell Science* **108**: 2537–2548.
- Skibbens, R.V., Corson, L.B., Koshland, D., and Hieter, P. 1999. Ctf7p is essential for sister chromatid cohesion and links mitotic chromosome structure to the DNA replication machinery. *Genes & Dev.* **13**: 307–319.
- Spencer, F., Gerring, S.L., Connelly, C., and Hieter, P. 1990. Mitotic chromosome transmission fidelity mutants in *Saccharomyces cerevisiae*. *Genetics* **124**: 237–249.
- Straight, A.F., Belmont, A.S., Robinett, C.C., and Murray, A.W. 1996. GFP tagging of budding yeast chromosomes reveals that protein-protein interactions can mediate sister chromatid cohesion. *Curr. Biol.* **6**: 1599–1608.
- Strunnikov, A.V. and Jessberger, R. 1999. Structural maintenance of chromosomes (SMC) proteins: Conserved molecular properties for multiple biological functions. *Eur. J. Biochem.* **263**: 6–13.

- Strunnikov, A.V., Larionov, V.L., and Koshland, D. 1993. SMC1: An essential yeast gene encoding a putative head-rod-tail protein is required for nuclear division and defines a new ubiquitous protein family. *J. Cell Biol.* **123**: 1635–1648.
- Sumner, A.T. 1991. Scanning electron microscopy of mammalian chromosomes from prophase to telophase. *Chromosoma* **100**: 410–418.
- Surana, U., Amon, A., Dowzer, C., McGrew, J., Byers, B., and Nasmyth, K. 1993. Destruction of the CDC28/CLB mitotic kinase is not required for the metaphase to anaphase transition in budding yeast. *EMBO J.* **12**: 1969–1978.
- Sutani, T. and Yanagida, M. 1997. DNA renaturation activity of the SMC complex implicated in chromosome condensation. *Nature* **388**: 798–801.
- Tanaka, T., Cosma, M.P., Wirth, K., and Nasmyth, K. 1999. Identification of cohesin association sites at centromeres and along chromosome arms. *Cell* **98**: 847–858.
- Tanaka, T., Fuchs, J., Loidl, J., and Nasmyth, K. 2000a. Cohesin ensures bipolar attachment of microtubules to sister centromeres and resists their precocious separation. *Nat. Cell Biol.* **2**: 492–499.
- Tanaka, K., Yonekawa, T., Kawasaki, Y., Kai, M., Furuya, K., Iwasaki, M., Murakami, H., Yanagida, M., and Okayama, H. 2000b. Fission yeast Eso1p is required for establishing sister chromatid cohesion during S phase. *Mol. Cell. Biol.* **20**: 3459–3469.
- Tang, T.T.L., Bickel, S.E., Young, L.M., and Orr-Weaver, T.L. 1998. Maintenance of sister-chromatid cohesion at the centromere by the *Drosophila* MEI-S332 protein. *Genes & Dev.* **12**: 3843–3856.
- Toth, A., Ciosk, R., Uhlmann, F., Galova, M., Schleiffer, A., and Nasmyth, K. 1999. Yeast cohesin complex requires a conserved protein, Eco1p(Ctf7), to establish cohesion between sister chromatids during DNA replication. *Genes & Dev.* **13**: 320–333.
- Uhlmann, F. and Nasmyth, K. 1998. Cohesion between sister chromatids must be established during DNA replication. *Curr. Biol.* **8**: 1095–1101.
- Uhlmann, F., Lottspeich, F., and Nasmyth, K. 1999. Sister-chromatid separation at anaphase onset is promoted by cleavage of the cohesin subunit Scc1. *Nature* **400**: 37–42.
- van Heemst, D., James, F., Poggeler, S., Berteaux-Lecellier, V., and Zickler, D. 1999. Spo76p is a conserved chromosome morphogenesis protein that links the mitotic and meiotic programs. *Cell* **98**: 261–271.
- Visintin, R., Prinz, S., and Amon, A. 1997. CDC20 and CDH1: A family of substrate-specific activators of APC-dependent proteolysis. *Science* **278**: 460–463.
- Wang, Z., Castano, I.B., De Las Penas, A., Adams, C., and Christman, M.F. 2000. Pol kappa: A DNA polymerase required for sister chromatid cohesion. *Science* **289**: 774–779.
- Waters, J.C., Skibbens, R.V., and Salmon, E.D. 1996. Oscillating mitotic newt lung cell kinetochores are, on average, under tension and rarely push. *J. Cell Science* **109**: 2823–2831.
- Waters, J.C., Chen, R.H., Murray, A.W., and Salmon, E.D. 1998. Localization of Mad2 to kinetochores depends on microtubule attachment, not tension. *J. Cell Biol.* **141**: 1181–1191.
- Waters, J.C., Chen, R.H., Murray, A.W., Gorbsky, G.J., Salmon, E.D., and Nicklas, R.B. 1999. Mad2 binding by phosphorylated kinetochores links error detection and checkpoint action in mitosis. *Curr. Biol.* **9**: 649–652.
- Wittmeyer, J. and Formosa, T. 1995. Identifying DNA replication complex components using protein affinity chromatography. *Methods Enzymol.* **262**: 415–430.
- Woodgate, R. 1999. A plethora of lesion-replicating DNA polymerases. *Genes & Dev.* **13**: 2191–2195.