Unlocking the brain for better learning

Julie Miwa

Assistant Professor
Biological Sciences
Lehigh University
Exploring the Brain

November 4th, 2013

Our ability to learn changes with age

Example: language

Later in life we can learn language, but it is harder and we might retain an accent

The brain changes as it ages

Younger brains are more adaptable to new information- they have greater plasticity

Language Center “Broca’s area” activity in bilingual people



Kim, Relkin, Lee, Hirsch, Nature, 1997

Cortical Reorganization



Figure 1. Two-dimensional, flattened views of *z*-score statistical parameter maps for visual cortex BOLD responses in early and late blind subjects. Maps in *A* and *B* were obtained, respectively, when subjects generated verbs to Braille read or heard nouns. A

Visual Cortex Activity in Early and Late Blind People

The Journal of Neuroscience, May 15, 2003, 23(10):4005–4011 -4005

fMRI

Physical changes occur in our brain as we age

The brain gets wired by early experiences

Different regions/functions have different sensitive periods

How does this happen?

It has been known that the sensitive period for robust plasticity – the “critical period closes”, but the mechanism for this is unknown



The lynx1 gene, discovered as a graduate student, up-regulates at the closing of the Critical Period in the visual cortex of mice



I removed the lynx1 gene through genetic engineering…



Creating lynx1KO mice

Morishita, Miwa, Heintz, Hensch, Science, 2010
Higgins, Science, (2010)

Classical Fear Conditioning (Associative Learning)



Training

24 hours



Associative “cued” learning



Contextual learning

Improved Learning By Increased Cholinergic Tone



lynx1KO mice exhibit enhanced **associative** learning

Miwa et al, 51, Neuron, 2006

Another animal study- Critical Period plasticity



Monkey visual cortex

The visual cortex gets input from two eyes

Each eye goes to one column in the visual cortex, creating a banded pattern (right/left)

Input from two eyes compete for cortical territory in the visual cortex

Critical Period Plasticity



If you close one eye, monocular deprivation (MD), the cortical territory receiving more light input will expand

If you don't correct cross-eyes early in humans (one eye loses the competition), you can lose depth perception

David Hubel and Torsten Wiesel received the Nobel prize for this work in 1981

The robust plasticity of the critical period was extended past the normal time frame.



P19-P33 in mice

Morishita, Miwa, Heintz, Hensch, Science, (2010)

The robust plasticity of the critical period was extended past the normal time frame.



Morishita, Miwa, Heintz, Hensch, Science, (2010)

P19-P33 in mice

This showed that the lynx1 gene acts as a molecular brake on Critical Period plasticity



P19-P33 in mice

How does lynx suppress learning?

Lynx1 binds nicotinic receptors of the cholinergic system

Lynx1 is similar to snake toxin proteins, which bind on muscle and inhibit



Miwa et al., 70, Neuron 2011

L...C...C...C...C...C...C...C...CC...LCN lynx1
L...C...C...C...C...C...C...C...CC...LCN Ly6
L...C...C...C...C...C...C...C...CC...LCN ThB
l....C.....C...C...C...C...CC...LCN α-btxn
l....C.....C...C...C...C...CC...LCN Cbtxn

Miwa et al., 23, Neuron 1999

Lynx1 modulates the cholinergic system

cholinergic neurons
- •localized discretely
- •project widely
- •release the neurotransmitter acetylcholine diffusely

•The cholinergic system can raise brain excitability

The cholinergic system contributes to learning

It operates on a gradient of activation



Optimized Cholinergic Tone



We can also raise cholinergic tone through pharmacology



Cholinergic agents that raise cholinergic tone can lead to enhanced critical period plasticity in normal adult mice



P19-P33 in mice

This demonstrates that the mechanisms for robust plasticity are still present in the adult brain

Cholinergic activation mediates this plasticity

BUT, that plasticity is suppressed by the presence of lynx modulators

We can also raise cholinergic tone through lynx dosage

overactivation

neurodegeneration
epilepsies
addiction

optimal

learning and memory
synaptic plasticity
neurotransmitter release
enhancement of attention

learning deficits
dementias
Alzheimer's disease

0 copies of lynx

1 copy of lynx

2 copies of lynx





Question: Can we remove the brake (lynx1) and recapture youthful plasticity in humans?



P19-P33 in mice

WHAT

What therapeutic advantages can we achieve if so?

Language learning in stroke

Learning deficits in dementias (Alzheimer’s disease)

Traumatic brain injury

HOW

How could we manipulate the lynx1 gene or lynx1 protein for learning benefits?

Further genetic manipulation- let animal grow up normally, and remove the lynx1 gene in adulthood

Environmental manipulations of lynx1 (Miwa et al., Physiology, 2012)

Individual variation in the lynx1 gene in the natural population

Closed ended learners **Open ended learners**



What is a SNP? Single Nucleotide Polymorphism



Sequence differences at a defined place in the person's gene
Many are scattered throughout many genes (the genome)
These are inherited mutations
They can exist in the population at different frequencies (1-49%)
They are used in forensic science to identify individuals (DNA fingerprinting)

expression levels (amount of protein)



Regulation (turn the gene on/ off more readily)

the amino acid sequence (protein function)

What if one SNP sequence made the lynx protein less effective?

Have better cognitive flexibility?

GACTTA

GAGTTA



Would that person learn better?

Be more creative?

The gene superfamily arose by gene duplication and Divergence

Gave rise to snake venom toxin protein through accelerated evolution

The lynx Family



lynx genes are highly conserved

Exist in all animals from C. elegans to humans

Most bind to nicotinic receptors

lynx Expression Patterns





lynx1

lynx2

amydgala

lynx Binding Specificity

	$\alpha4\beta2$	$\alpha7$	muscle	$\alpha4\beta4$	$\alpha3\beta4$
lynx1	+++	+++	+	+++	-
lynx2	+++	+	+++	+++	-
lynx5	-	-	-	+	-

lynx Expression Patterns





lynx1

lynx2

Cognition and learning

fear and anxiety

WHY

If lynx1 suppresses cholinergic activity, why have it?

Control cholinergic tone in discrete areas

Acknowledgements

HHMI/Rockefeller University
Nathaniel Heintz

Library screen
Siobhan Kuhar
Lei Feng

In vitro functional and binding studies
Ines Ibanez-Tallon

lynx family members
Ayse Tekinay

Rockefeller University
Hippocampal slices:
Constantine Pavlides

Structural Modeling
Andrej Sali
Roberto Sanchez

Lynx2 electrophysiology
Paul Greengard
Yi Nong

Yale School of Medicine
Marina Picciotto
Reiko Maki Fitzsimonds
Sarah King
Barbara Caldarone
Tanya Stevens

Caltech

Henry Lester
Doreen Rhee
Rell Parker
Mark Starbird
Po Ku
Atsuko Kobayashi
Weston Nichols
Cheng Xiao

Harvard University

Takao Hensch
Hirofumi Morishita

Other:
Steve Sine- Mayo Clinic
Hai-Long Wang
Lorna Role-Columbia University
Gregg Crabtree

Lehigh Students

Kate Oliver- Presidential scholar
Adam Van Handel- Eckhardt scholar
Tim Yeh
Sana Ali- sophomore
Chris Hoke
Kyra Feuer- BDSI

Former students

Carly Garrison- Presidential scholar
Kasarah Ackerman- graduated 2014
Courtney Meyer- President of Lehigh's Neuroscience club
Chelsea Serrano
Jacquelin Botello

Julie Miwa
Biological Sciences

lynx Acts As A Molecular Brake



Addition of lynx shifts dose response curve to the right



lynx1 removal shifts dose response curve to the left



Oocytes

Co-expression of lynx with α4β2 nAChRs:

- reduces sensitive to agonist
- increases desensitization kinetics
- slows recovery from desensitization



Brain slice recordings

Ibanez-Tallon et al., 33, Neuron, 2002

Miwa et al., Neuron, 51, 2006