



Why Don't These Drugs Work Anymore?

Biosciences in the 21st Century

Dr. Amber Rice

November 17, 2014

Estimates of Burden of Antibacterial Resistance

European Union *population 500m*

25,000 deaths per year

2.5m extra hospital days

Overall societal costs
(€ 900 million, hosp. days)
Approx. €1.5 billion per year



Source: ECDC 2007

Thailand *population 70m*

>38,000 deaths

>3.2m hospital days

Overall societal costs
US\$ 84.6–202.8 mill. direct
>US\$1.3 billion indirect




Source: Pumart et al 2012

United States *population 300m*

>23,000 deaths

>2.0m illnesses

Overall societal costs
Up to \$20 billion direct
Up to \$35 billion indirect



Source: US CDC 2013


Global information is insufficient to show complete disease burden impact and costs

Why don't these drugs work anymore?

- Drug resistance: a case study
- How does resistance evolve?
 - Three mechanisms of evolution
- Examples of “superbugs”
- Avoiding more widespread resistance


Drug resistance: a case study

- AZT (azidothymidine) approved as a treatment for HIV in 1987.



Drug resistance: a case study


- After several months of treatment, higher concentrations of AZT were needed.



Drug resistance: a case study

- In most cases, patients became resistant to AZT within 6 months!

(b) Resistance in 39 patients, checked at different times



What happened?


Why might a drug stop working so quickly?

Why don't these drugs work anymore?


- Drug resistance: a case study
- How does resistance evolve?
 - Three mechanisms of evolution
- Examples of “superbugs”
- Avoiding more widespread resistance

Back to our case study: the evolution of resistance

Susceptible



Resistant




Mechanisms of evolution

1. Natural selection: differential reproductive success
 - Non-random
 - Not forward-looking, can only work with existing variation
 - Only adaptive mechanism of evolution



Back to our case study: the evolution of resistance by natural selection



Mechanisms of evolution: sources of variation

2. Mutation: a change in DNA sequence, gene order, or chromosome number


- Random
- Increases genetic variation within populations
- Types of mutations:
 - Point mutations
 - Insertions
 - Deletions
 - Gene duplications
 - Chromosomal inversions
 - Polyploidy



Mechanisms of evolution: sources of variation


3. Gene flow (or migration): movement of genes between populations

- Increases genetic variation within populations
- Makes populations more similar to each other



Evolution of resistance: gene flow introduces variation

- Bacteria can also pick-up resistance genes through **horizontal gene transfer**



Are the necessary “ingredients”
present for the evolution of
antimicrobial resistance by natural
selection?

Evolution of resistance


- Heritable variation for resistance
 - 30,000 year-old bacterial DNA recovered from Yukon permafrost
 - Genes for resistance to several antibiotics, including tetracycline and vancomycin (D'Costa et al. 2011, Nature)
 - So, resistant strains:
 - can pre-date use of the antimicrobial drug.
 - may arise by random mutation or even gene transfer after the drug is in use.

Evolution of resistance

- Differential reproduction
 - Widespread use of antibiotics creates **strong selection** for resistant strains.
 - Antibiotics over-prescribed by doctors
 - Antibiotics used in agriculture and commercial products
 - Use of *any* anti-microbial drug, not only antibiotics, creates strong selection for resistance.

Back to our case study: new treatments informed by evolution

- By understanding how resistance evolves, researchers could design new treatments.



Why are drug cocktails more effective?

- With a single drug, only 1 mutation can confer resistance.
 - Easy. HIV has large populations, a short generation time, and a high mutation rate.
- For resistance to drug cocktails,
- More mutations needed for resistance → lower probability the mutations will occur together in one virion


Why don't these drugs work anymore?

- Drug resistance: a case study
- How does resistance evolve?
 - Three mechanisms of evolution
- **Examples of “superbugs”**
- Avoiding more widespread resistance

MRSA

Methicillin-resistant *Staphylococcus aureus*


- Spread in hospitals and with close physical contact (e.g., among inmates, athletes)
- 33% worldwide have Staph, ~1% MRSA
- Painful skin conditions, even bacterial pneumonia and blood infections
- Can be fatal
- Resistant to entire class of penicillin-like antibiotics
- In 2002, vancomycin-resistant strain found



MDR-TB

Multidrug-resistant Tuberculosis

- TB is major cause of death worldwide.
 - 2 million TB-related deaths each year
- 440,000 MDR-TB cases each year
 - 150,000 deaths
- 2nd line drugs have more side-effects, cost up to 100x more!



Malaria

- Caused by *Plasmodium spp.* protozoan
- Transmitted by mosquito
- Tropical and sub-tropical regions
- Fever, muscle & back pain, vomiting, anemia...
- Brain damage in children
- Nearly 1 million deaths each year
- Drugs used for treatment and for prevention
- Resistance to cheapest and most commonly used drugs is widespread
- Resistance to newer drugs is emerging



World Health Organization (WHO)

U.S. National Institute of Allergy and Infectious Diseases (NIAID)

Why don't these drugs work anymore?


- Drug resistance: a case study
- How does resistance evolve?
 - Three mechanisms of evolution
- Examples of “superbugs”
- **Avoiding more widespread resistance**

Avoiding more widespread resistance

1. Avoid contracting infections
2. Minimize transmission of resistant microbes
3. Improve use of antimicrobial drugs
 - Take only when appropriate (i.e., don't take an antibiotic for the flu!)
 - Use antibacterial soaps/cleaners ONLY around people with weakened immune systems
 - Avoid broad-spectrum antibiotics if possible.
 - Take ALL of the medication
 - Reduce agricultural use of antibiotics

Why can reducing inappropriate use of antimicrobial drugs combat resistance?

- Resistance is sometimes costly for microbes.




Why can reducing inappropriate use of antimicrobial drugs combat resistance?

- Use of specific antibiotics (not broad-spectrum)
 - Some antibiotics target a greater number of bacterial species.
 - Often used when diagnosis is unclear.
 - Why is it better to prescribe an antibiotic that targets fewer species?
 - Selection for resistance will act only on the species that are targeted by the drug.

Why can reducing inappropriate use of antimicrobial drugs combat resistance?

- Taking ALL of the medication increases the chance of exterminating the microbial population before resistance evolves.
 - If you stop early, you may get sick again or stay sick for longer.
 - Longer illness → more bacterial generations → greater chance of mutation for resistance arising
 - Even if resistant microbes DO arise, immune system may successfully fight them if population is small.
 - Stopping the drug lets the population grow larger.

Why can reducing inappropriate use of antimicrobial drugs combat resistance?



- Resistant bacteria escape livestock, spread to humans

Current research aims

- What is the mechanism of resistance?
- How do microbes acquire and pass on resistance genes?
- Development of better diagnostic tests to avoid the need for “broad spectrum” antibiotics
- Development of new drugs/vaccines

Over the last 30 years, no major new types of antibiotics have been developed

