

LIMIT THEORY FOR GEOMETRIC STATISTICS OF POINT PROCESSES HAVING FAST DECAY OF CORRELATIONS

BY B. BŁASZCZYSZYN, D. YOGESHWARAN * AND J. E. YUKICH †

Inria/ENS, Paris; Indian Statistical Institute, Bangalore and Lehigh University, Bethlehem

Let \mathcal{P} be a simple, stationary point process on \mathbb{R}^d having fast decay of correlations, i.e., its correlation functions factorize up to an additive error decaying faster than any power of the separation distance. Let $\mathcal{P}_n := \mathcal{P} \cap W_n$ be its restriction to windows $W_n := [-\frac{1}{2}n^{1/d}, \frac{1}{2}n^{1/d}]^d \subset \mathbb{R}^d$. We consider the statistic $H_n^\xi := \sum_{x \in \mathcal{P}_n} \xi(x, \mathcal{P}_n)$ where $\xi(x, \mathcal{P}_n)$ denotes a score function representing the interaction of x with respect to \mathcal{P}_n . When ξ depends on local data in the sense that its radius of stabilization has an exponential tail, we establish expectation asymptotics, variance asymptotics, and central limit theorems for H_n^ξ and, more generally, for statistics of the re-scaled, possibly signed, ξ -weighted point measures $\mu_n^\xi := \sum_{x \in \mathcal{P}_n} \xi(x, \mathcal{P}_n) \delta_{n^{-1/d}x}$, as $W_n \uparrow \mathbb{R}^d$. This gives the limit theory for non-linear geometric statistics (such as clique counts, the number of Morse critical points, intrinsic volumes of the Boolean model, and total edge length of the k -nearest neighbors graph) of α -determinantal point processes (for $-1/\alpha \in \mathbb{N}$) having fast decreasing kernels, including the β -Ginibre ensembles, extending the Gaussian fluctuation results of Soshnikov [68] to non-linear statistics. It also gives the limit theory for geometric U-statistics of α -permanental point processes (for $1/\alpha \in \mathbb{N}$) as well as the zero set of Gaussian entire functions, extending the central limit theorems of Nazarov and Sodin [50] and Shirai and Takahashi [67], which are also confined to linear statistics. The proof of the central limit theorem relies on a factorial moment expansion originating in [11, 12] to show the fast decay of the correlations of ξ -weighted point measures. The latter property is shown to imply a condition equivalent to Brillinger mixing and consequently yields the asymptotic normality of μ_n^ξ via an extension of the cumulant method.

CONTENTS

1	Introduction and main results	2
1.1	Admissible point processes having fast decay of correlations	6
1.2	Admissible score functions	9

*Research supported by DST-INSPIRE faculty award, CPDA from the Indian Statistical Institute and TOPOSYS grant.

†Research supported in part by NSF grant DMS-1406410

MSC 2010 subject classifications: Primary:60F05, 60D05; Secondary:60G55, 52A22, 05C80

Keywords and phrases: Point processes having fast decay of correlations, determinantal point process, permanental point process, Gaussian entire functions, Gibbs point process, U-statistics, stabilization, difference operators, cumulants, Brillinger mixing, central limit theorem

scribe a global geometric feature of a structure on \mathcal{X} in terms of local contributions $\xi(x, \mathcal{X})$.

It is frequently the case in stochastic geometry, statistical physics, and spatial statistics that one seeks the large n limit behavior of

$$(1.2) \quad H_n^\xi := H_n^\xi(\mathcal{P}) := \sum_{x \in \mathcal{P}_n} \xi(x, \mathcal{P}_n)$$

where ξ is an appropriately chosen score function, \mathcal{P} is a simple, stationary point process on \mathbb{R}^d , and \mathcal{P}_n is the restriction of \mathcal{P} to $W_n := [-\frac{1}{2}n^{1/d}, \frac{1}{2}n^{1/d}]^d$. For example if \mathcal{P}_n is either a Poisson or binomial point process and if ξ is either a local U -statistic or an exponentially stabilizing score function, then the limit theory for H_n^ξ is established in [6, 22, 38, 41, 55, 57, 60, 62]. If \mathcal{P}_n is a rarified Gibbs point process on W_n and ξ is exponentially stabilizing, then [65, 69] treat the limit theory for H_n^ξ .

It is natural to ask whether the limit theory of these papers extends to more general input satisfying a notion of ‘asymptotic independence’ for point processes. Recall that if $\xi \equiv 1$ and if \mathcal{P} is an α -determinantal point process with $\alpha = -1/m$ or an α -permanental point process with $\alpha = 2/m$ for some m in the set of positive integers \mathbb{N} (respectively \mathcal{P} is the zero set of a Gaussian entire function), then remarkable results of Soshnikov [68], Shirai and Takahashi [67] (respectively Nazarov and Sodin [50]), show that the counting statistic $\mathcal{P}_n(W_n) := \sum_{x \in \mathcal{P}_n} \mathbf{1}[x \in W_n]$ is asymptotically normal. One may ask whether asymptotic normality of H_n^ξ still holds when ξ is either a local U -statistic or an exponentially stabilizing score function. We answer these questions affirmatively. Loosely speaking, subject to a mild growth condition on $\text{Var} H_n^\xi$, our approach shows that H_n^ξ is asymptotically normal whenever \mathcal{P} is a point process having fast decay of correlations.

Heuristically, when the score functions depend on ‘local data’ and when the input is ‘asymptotically independent’, one might expect that the statistics H_n^ξ obey a strong law and a central limit theorem. The notion of dependency on ‘local data’ for score functions is formalized via stabilization in [6, 22, 55, 57, 60]. Here we formalize the idea of asymptotically independent input \mathcal{P} via the notion of ‘fast decay of correlation functions’. We thereby extend the limit theory of the afore-mentioned papers to input having fast decay of correlation functions. A point process \mathcal{P} on \mathbb{R}^d has fast decay of correlations if for all $p, q \in \mathbb{N}$ and all $x_1, \dots, x_{p+q} \in \mathbb{R}^d$, its correlation functions $\rho^{(p+q)}(x_1, \dots, x_{p+q})$ factorize into $\rho^{(p)}(x_1, \dots, x_p)\rho^{(q)}(x_{p+1}, \dots, x_{p+q})$ up to an additive error decaying faster than any power of the separation distance

$$(1.3) \quad s := d(\{x_1, \dots, x_p\}, \{x_{p+1}, \dots, x_{p+q}\}) := \inf_{i \in \{1, \dots, p\}, j \in \{p+1, \dots, p+q\}} |x_i - x_j|$$

Given input \mathcal{P} having fast decay of correlations, an interesting feature of the measures μ_n^ξ is that their variances are at most of order $\text{Vol}_d(W_n)$, the volume of the window W_n (Theorem 1.12). This holds also for the statistic $\hat{H}_n^\xi := \sum_{x \in \mathcal{P}_n} \xi(x, \mathcal{P})$, which involves summands having no boundary effects. An interesting feature of this statistic is that if its variance is $o(\text{Vol}_d(W_n))$ then it has to be $O(\text{Vol}_{d-1}(\partial W_n))$, where ∂W_n denotes the boundary of W_n and $\text{Vol}_{d-1}(\cdot)$ stands for the $(d-1)$ th intrinsic volume (Theorem 1.15). In other words, if the fluctuations of \hat{H}_n^ξ are not of volume order, then they are at most of surface order.

Coming back to our set-up, when a functional $H_n^\xi(\mathcal{P})$ is expressible as a sum of local U -statistics or, more generally, as a sum of exponentially stabilizing score functions ξ , then a key step towards proving the central limit theorem is to show that the correlation functions of the ξ -weighted measures defined via Palm expectations $\mathbb{E}_{x_1, \dots, x_k}$ (cf Section 1.1) and given by

$$(1.6) \quad m^{(k_1, \dots, k_{p+q})}(x_1, \dots, x_{p+q}; n) := \mathbb{E}_{x_1, \dots, x_{p+q}}(\xi(x_1, \mathcal{P}_n)^{k_1} \dots \xi(x_{p+q}, \mathcal{P}_n)^{k_{p+q}}) \\ \times \rho^{(p+q)}(x_1, \dots, x_{p+q}),$$

similar to those of the input process \mathcal{P} , approximately factorize into

$$m^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n) m^{(k_{p+1}, \dots, k_{p+q})}(x_{p+1}, \dots, x_{p+q}; n),$$

uniformly in $n \leq \infty$, up to an additive error decaying faster than any power of the separation distance s , defined at (1.3). Here x_1, \dots, x_{p+q} are distinct points in W_n and $k_1, \dots, k_{p+q} \in \mathbb{N}$. This result, spelled out in Theorem 1.11, is at the heart of our approach. We then give two proofs of the central limit theorem (Theorem 1.13) for the purely atomic random measures (1.4) via the cumulant method, and as a corollary, derive the asymptotic normality of $H_n^\xi(\mathcal{P})$ and $\int f d\mu_n^\xi$, f a test function, as $n \rightarrow \infty$. The proof of expectation and variance asymptotics (Theorem 1.12) mainly relies upon the refined Campbell theorem.

In contrast to the afore-mentioned works, our proof of the fast decay of correlations of the ξ -weighted measures depends heavily on a factorial moment expansion for expected values of functionals of a general point process \mathcal{P} . This expansion, which originates in [11, 12], is expressed in terms of iterated difference operators of the considered functional on the null configuration of points and integrated against factorial moment measures of the point process. It is valid for general point processes, in contrast to the Fock space representation of Poisson functionals, which involves the same difference operators but is deeply related to chaos expansions [39]. Further connections with the literature are discussed in the remarks following Theorems 1.14 and 1.15.

Our interest in these issues was stimulated by similarities in the methods of [42], [5, 6, 65] and [50]. The articles [6, 65] prove central limit theorems for stabilizing

the k -point correlation function (or k th joint intensity) and is characterized by the relation

$$(1.7) \quad \alpha^{(k)}(B_1 \times \cdots \times B_k) = \mathbb{E} \left(\prod_{1 \leq i \leq k} \mathcal{P}(B_i) \right) = \int_{B_1 \times \cdots \times B_k} \rho^{(k)}(x_1, \dots, x_k) dx_1 \dots dx_k,$$

where B_1, \dots, B_k are mutually disjoint bounded Borel sets in \mathbb{R}^d . Since \mathcal{P} is simple, we may put $\rho^{(k)}$ to be zero on the diagonals of $(\mathbb{R}^d)^k$, that is on the subsets of $(\mathbb{R}^d)^k$ where two or more coordinates coincide. The disjointness assumption is crucial as illustrated by the following useful relation: For any bounded Borel set $B \subset \mathbb{R}^d$ and $k \geq 1$, we have

$$(1.8) \quad \alpha^{(k)}(B^k) = \mathbb{E} \left(\mathcal{P}(B)(\mathcal{P}(B)-1) \dots (\mathcal{P}(B)-k+1) \right) = \int_{B^k} \rho^{(k)}(x_1, \dots, x_k) dx_1 \dots dx_k.$$

Heuristically, the k th Palm measure P_{x_1, \dots, x_k} of \mathcal{P} is the probability distribution of \mathcal{P} conditioned on $\{x_1, \dots, x_k\} \subset \mathcal{P}$. More formally, if $\alpha^{(k)}$ is locally finite, there exists a family of probability distributions P_{x_1, \dots, x_k} on $(\mathcal{N}, \mathcal{B})$, unique up to an $\alpha^{(k)}$ -null set of $(\mathbb{R}^d)^k$, called the k th Palm measures of \mathcal{P} , and satisfying the disintegration formula

$$(1.9) \quad \mathbb{E} \left(\sum_{(x_1, \dots, x_k) \in \mathcal{P}^{(k)}} f(x_1, \dots, x_k; \mathcal{P}) \right) = \int_{(\mathbb{R}^d)^k} \int_{\mathcal{N}} f(x_1, \dots, x_k; \mu) P_{x_1, \dots, x_k}(d\mu) \alpha^{(k)}(dx_1 \dots dx_k)$$

for any (say non-negative) measurable function f on $(\mathbb{R}^d)^k \times \mathcal{N}$. Formula (1.9) is also known as the refined Campbell theorem.

To simplify notation, write $\int_{\mathcal{N}} f(x_1, \dots, x_k; \mu) P_{x_1, \dots, x_k}(d\mu) = \mathbb{E}_{x_1, \dots, x_k}(f(x_1, \dots, x_k; \mathcal{P}))$, where $\mathbb{E}_{x_1, \dots, x_k}$ is the expectation corresponding to the Palm probability $\mathbb{P}_{x_1, \dots, x_k}$ on a canonical probability space on which \mathcal{P} is also defined. To further simplify notation, denote by $\mathbb{P}_{x_1, \dots, x_k}^!$ the reduced Palm probabilities and their expectation by $\mathbb{E}_{x_1, \dots, x_k}^!$, which satisfies $\mathbb{E}_{x_1, \dots, x_k}^!(f(x_1, \dots, x_k; \mathcal{P})) = \mathbb{E}_{x_1, \dots, x_k}(f(x_1, \dots, x_k; \mathcal{P} \setminus \{x_1, \dots, x_k\}))$ ¹.

All Palm probabilities (expectations) are meaningfully defined only for $\alpha^{(k)}$ almost all $x_1, \dots, x_k \in \mathbb{R}^d$. Consequently, all expressions involving these measures should be understood in the $\alpha^{(k)}$ a.e. sense and suprema should likewise be understood as essential suprema with respect to $\alpha^{(k)}$.

The following definition is reminiscent of the so-called weak exponential decrease of correlations introduced in [42] and subsequently used in [5, 44, 50].

DEFINITION 1.1 (ω -mixing correlation functions). *The correlation functions*

¹ It can be shown that $\mathbb{P}_{x_1, \dots, x_k}(x_1, \dots, x_k \in \mathcal{P}) = 1$ for $\alpha^{(k)}$ a.e. $x_1, \dots, x_k \in \mathbb{R}^d$.

of a point process \mathcal{P} are ω -mixing if there exists a decreasing function $\omega : \mathbb{N} \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for all $n \in \mathbb{N}$, $\lim_{x \rightarrow \infty} \omega(n, x) = 0$ and for all $p, q \in \mathbb{N}$, $x_1, \dots, x_{p+q} \in \mathbb{R}^d$, we have

$$|\rho^{(p+q)}(x_1, \dots, x_{p+q}) - \rho^{(p)}(x_1, \dots, x_p)\rho^{(q)}(x_{p+1}, \dots, x_{p+q})| \leq \omega(p+q, s),$$

where $s := d(\{x_1, \dots, x_p\}, \{x_{p+1}, \dots, x_{p+q}\})$ is as at (1.3).

By an admissible point process \mathcal{P} on \mathbb{R}^d , $d \geq 2$, we mean that \mathcal{P} is simple, stationary (i.e., $\mathcal{P} + x \stackrel{d}{=} \mathcal{P}$ for all $x \in \mathbb{R}^d$, where $\mathcal{P} + x$ denotes the translation of \mathcal{P} by the vector x), with non-null and finite intensity $\rho^{(1)}(\mathbf{0}) = \mathbb{E}(\mathcal{P}(W_1))$, and has k -point correlation functions of all orders $k \in \mathbb{N}$. By a fast decreasing function $\phi : \mathbb{R}^+ \rightarrow [0, 1]$ we mean ϕ satisfies $\lim_{x \rightarrow \infty} x^m \phi(x) = 0$ for all $m \geq 1$.

DEFINITION 1.2 (Admissible point process having fast decay of correlations). *Let \mathcal{P} be an admissible point process. \mathcal{P} is said to have fast decay of correlations if its correlation functions are ω -mixing as in Definition 1.1 with $\omega(n, x) = C_n \phi(c_n x)$ for some correlation decay constants $c_n, C_n \in (0, \infty)$ and a fast decreasing function $\phi : \mathbb{R}^+ \rightarrow [0, 1]$, called a correlation decay function.*

More explicitly, an admissible point process has *fast decay of correlations*, if for all $p, q \in \mathbb{N}$ and all $(x_1, \dots, x_{p+q}) \in (\mathbb{R}^d)^{p+q}$

$$(1.10) \quad |\rho^{(p+q)}(x_1, \dots, x_{p+q}) - \rho^{(p)}(x_1, \dots, x_p)\rho^{(q)}(x_{p+1}, \dots, x_{p+q})| \leq C_{p+q}\phi(c_{p+q}s),$$

where $s := d(\{x_1, \dots, x_p\}, \{x_{p+1}, \dots, x_{p+q}\})$ is as at (1.3) and C_k, c_k, ϕ are as in Definition 1.2. Without loss of generality, we assume that c_k is non-increasing in k , and that $C_k \in [1, \infty)$ is non-decreasing in k . As a by-product of our proof of the asymptotic normality of μ_n^ξ in (1.4), we establish that the fast decay of correlations of \mathcal{P} implies that it is Brillinger mixing; cf Remark (vi) in Section 1.4 and Remarks at the end of Section 4.4.2.

Admissible point processes having fast decay of correlations are ubiquitous and include certain determinantal, permanental, and Gibbs point processes, as explained in Section 2.2. The k -point correlation functions of admissible point processes having fast decay of correlations are bounded i.e.,

$$(1.11) \quad \sup_{(x_1, \dots, x_k) \in (\mathbb{R}^d)^k} \rho^{(k)}(x_1, \dots, x_k) \leq \kappa_k < \infty,$$

for some constants κ_k , which without loss of generality are assumed non-decreasing in k . Also without loss of generality, assume $\kappa_0 := \max\{\rho^{(1)}(\mathbf{0}), 1\}$. For station-

ary \mathcal{P} with intensity $\rho^{(1)}(\mathbf{0}) \in (0, \infty)$ we have that (1.10) implies (1.11) with

$$(1.12) \quad \kappa_k \leq (\rho^{(1)}(\mathbf{0}))^k + \sum_{i=2}^k C_i (\rho^{(1)}(\mathbf{0}))^{k-i} \leq k C_k \kappa_0^k.$$

The bound (1.12) helps to determine when point processes having fast decay of correlations also have exponential moments, as in Section 2.1.

1.2. Admissible score functions. Throughout we restrict to translation-invariant score functions $\xi : \mathbb{R}^d \times \mathcal{N} \rightarrow \mathbb{R}$, i.e., those which are measurable in each coordinate, $\xi(x, \mathcal{X}) = 0$ if $x \notin \mathcal{X} \in \mathcal{N}$, and for all $y \in \mathbb{R}^d$, satisfy $\xi(\cdot + y, \cdot + y) = \xi(\cdot, \cdot)$.

We introduce classes (A1) and (A2) of *admissible* score and input pairs (ξ, \mathcal{P}) . Specific examples of admissible input pairs of both classes are provided in Sections 2.2 and 2.3. The first class allows for admissible input \mathcal{P} as in Definition 1.2 whereas the second considers admissible input \mathcal{P} having fast decay of correlations (1.10), subject to $c_k \equiv 1$ and growth conditions on the decay constants C_k and the decay function ϕ .

DEFINITION 1.3 (Class (A1) of admissible score and input pairs (ξ, \mathcal{P})). *Admissible input \mathcal{P} consists of admissible point processes having fast decay of correlations as in Definition 1.2. Admissible score functions are of the form*

$$(1.13) \quad \xi(x, \mathcal{X}) := \frac{1}{k!} \sum_{\mathbf{x} \in \mathcal{X}^{(k-1)}} h(x, \mathbf{x}),$$

for some $k \in \mathbb{N}$ and a symmetric, translation-invariant function $h : \mathbb{R}^d \times (\mathbb{R}^d)^{k-1} \rightarrow \mathbb{R}$ such that $h(x_1, \dots, x_k) = 0$ whenever either $\max_{2 \leq i \leq k} |x_i - x_1| > r$ for some given $r > 0$ or when $x_i = x_j$ for some $i \neq j$. When $k = 1$, we set $\xi(x, \mathcal{X}) = h(x)$. Further, assume

$$\|h\|_\infty := \sup_{\mathbf{x} \in \mathbb{R}^{d(k-1)}} |h(\mathbf{0}, \mathbf{x})| < \infty.$$

The interaction range for h is at most r , showing that the functionals H_n^ξ defined at (1.2) generated via scores (1.13) are local U -statistics of order k as in [62]. Before introducing a more general class of score functions, we recall [6, 41, 55, 57, 60] a few definitions formalizing the notion of the local dependence of ξ on its input. Let $B_r(x) := \{y : |y - x| \leq r\}$ denote the ball of radius r centered at x and $B_r^c(x)$ its complement.

DEFINITION 1.4 (Radius of stabilization). *Given a score function ξ , input \mathcal{X} , and $x \in \mathcal{X}$, define the radius of stabilization $R^\xi(x, \mathcal{X})$ to be the smallest $r \in \mathbb{N}$*

there exists $\hat{c} \in [1, \infty)$ such that for all $r \in (0, \infty)$

$$(1.18) \quad |\xi(x, \mathcal{X} \cap B_r(x))| \mathbf{1}[\text{card}(\mathcal{X} \cap B_r(x)) = n] \leq (\hat{c} \max(r, 1))^n.$$

The condition $c_k \equiv 1$ is equivalent to $c_* := \inf c_k > 0$. This follows since we may replace the fast decreasing function $\phi(\cdot)$ by $\phi(c_* \times \cdot)$, with $c_k \equiv 1$ for this new fast decreasing function. Score functions of class (A1) also satisfy the power growth condition (1.18) since in this case the left hand side of (1.18) is at most $\|h\|_\infty n^{(k-1)}/k$. Thus the generalization from (A1) to (A2) consists in replacing local U-statistics by exponentially stabilizing score functions satisfying the power growth condition. This is done at the price of imposing stronger conditions on the input process, requiring in particular that it has finite exponential moments, as explained in Section 2.1.

1.3. Fast decay of correlations of the ξ -weighted measures. The following p -moment condition involves the score function ξ and the input \mathcal{P} . We shall describe in Section 2.1 ways to control the p -moments of input pairs of class (A1) and (A2).

DEFINITION 1.8 (Moment condition). *Given $p \in [1, \infty)$, say that the pair (ξ, \mathcal{P}) satisfies the p -moment condition if*

$$(1.19) \quad \sup_{1 \leq n \leq \infty} \sup_{1 \leq p' \leq [p]} \sup_{x_1, \dots, x_{p'} \in W_n} \mathbb{E}_{x_1, \dots, x_{p'}} |\xi(x_1, \mathcal{P}_n)|^p \leq \tilde{M}_p < \infty$$

for some constant $\tilde{M}_p := \tilde{M}_p^\xi \in [1, \infty)$, where \sup signifies *ess sup* with respect to $\alpha^{(p)}$. Without loss of generality we assume that \tilde{M}_p is increasing in p for all p such that (1.19) holds.

We next consider the decay of the functions at (1.6), the so-called correlation functions of the ξ -weighted measures at (1.5). These functions indeed play the same role as the k -point correlation functions of the simple point process \mathcal{P} . When $\xi \equiv 1$ they obviously reduce to the correlation functions of \mathcal{P} . For general ξ and $k_i \equiv 1$ they are densities (‘mixed moment densities’ in the language of [6]) of the higher-order moment measures of the ξ -weighted measures with all distinct arguments. In the case of repeated arguments, the moment measures of a simple point process ‘collapse’ to appropriate lower dimensional ones. This is neither the case for non-simple point processes nor for our ξ -weighted measures, where general exponents k_i are required to properly take into account repeated arguments.

When $k_i \equiv 1$ for all $1 \leq i \leq p$, we write $m_{(p)}(x_1, \dots, x_p; n)$ instead of $m^{(1, \dots, 1)}(x_1, \dots, x_p; n)$. Abbreviate $m^{(k_1, \dots, k_p)}(x_1, \dots, x_p; \infty)$ by $m^{(k_1, \dots, k_p)}(x_1, \dots, x_p)$. These functions exist whenever (1.19) is satisfied for p set to $k_1 + \dots + k_p$ and provided the p -point correlation function $\rho^{(p)}$ exists. As for the input process \mathcal{P} we

1.4. **Main results.** We give the limit theory for the measures $\mu_n^\xi, n \geq 1$, defined at (1.4). Given a score function ξ on admissible input \mathcal{P} we set ²

$$(1.22) \quad \sigma^2(\xi) := \mathbb{E}_0 \xi^2(\mathbf{0}, \mathcal{P}) \rho^{(1)}(\mathbf{0}) + \int_{\mathbb{R}^d} (m_{(2)}(\mathbf{0}, x) - m_{(1)}(\mathbf{0})^2) dx.$$

The following result provides expectation and variance asymptotics for $\mu_n^\xi(f)$, with f belonging to the space $\mathcal{B}(W_1)$ of bounded measurable functions on W_1 .

THEOREM 1.12. *Let \mathcal{P} be an admissible point process on \mathbb{R}^d .*

(i) *If ξ satisfies exponential stabilization (1.15) and if (ξ, \mathcal{P}) satisfies the p -moment condition (1.19) for some $p \in (1, \infty)$ then for all $f \in \mathcal{B}(W_1)$*

$$(1.23) \quad \left| n^{-1} \mathbb{E} \mu_n^\xi(f) - \mathbb{E}_0 \xi(\mathbf{0}, \mathcal{P}) \rho^{(1)}(\mathbf{0}) \int_{W_1} f(x) dx \right| = O(n^{-1/d}).$$

If ξ only satisfies stabilization (1.14) and the p -moment condition (1.19) for some $p \in (1, \infty)$, then the right hand side of (1.23) is $o(1)$.

(ii) *Assume that the second correlation function $\rho^{(2)}$ of \mathcal{P} exists and is bounded as in (1.11), that ξ satisfies (1.14), and that (ξ, \mathcal{P}) satisfies the p -moment condition (1.19) for some $p \in (2, \infty)$. If the second-order correlations of the ξ -weighted measures decay fast, i.e. satisfy (1.21) with $p = q = k_1 = k_2 = 1$ and all $n \in \mathbb{N} \cup \{\infty\}$, then for all $f \in \mathcal{B}(W_1)$*

$$(1.24) \quad \lim_{n \rightarrow \infty} n^{-1} \text{Var} \mu_n^\xi(f) = \sigma^2(\xi) \int_{W_1} f(x)^2 dx \in [0, \infty),$$

whereas for all $f, g \in \mathcal{B}(W_1)$

$$(1.25) \quad \lim_{n \rightarrow \infty} n^{-1} \text{Cov}(\mu_n^\xi(f), \mu_n^\xi(g)) = \sigma^2(\xi) \int_{W_1} f(x)g(x) dx.$$

We remark that (1.23) and (1.24) together show convergence in probability

$$n^{-1} \mu_n^\xi(f) \xrightarrow{P} \mathbb{E}_0 \xi(\mathbf{0}, \mathcal{P}) \rho^{(1)}(\mathbf{0}) \int_{W_1} f(x) dx \text{ as } n \rightarrow \infty.$$

The proof of variance asymptotics (1.24) requires fast decay of the second-order correlations of the ξ -weighted measures. Fast decay of *all* higher-order correlations as in Definition 1.10 yields Gaussian fluctuations of $\mu_n^\xi, n \geq 1$, under moment conditions on the atom sizes (i.e., under moment conditions on ξ) and a variance lower bound. Let N denote a mean zero normal random variable

²For a stationary point process \mathcal{P} , its Palm expectation \mathbb{E}_0 (and consequently $m_{(1)}(\mathbf{0}), m_{(2)}(\mathbf{0}, x)$) is meaningfully defined e.g. via the Palm-Matthes approach.

with variance 1. We write $f(n) = \Omega(g(n))$ when $g(n) = O(f(n))$, i.e., when $\liminf_{n \rightarrow \infty} |f(n)/g(n)| > 0$.

THEOREM 1.13. *Let \mathcal{P} be an admissible point process on \mathbb{R}^d and let the pair (ξ, \mathcal{P}) satisfy the p -moment condition (1.19) for all $p \in (1, \infty)$. If the correlations of the ξ -weighted measures at (1.5) decay fast as in Definition 1.10 and if $f \in \mathcal{B}(W_1)$ satisfies*

$$(1.26) \quad \text{Var} \mu_n^\xi(f) = \Omega(n^\nu)$$

for some $\nu \in (0, \infty)$, then as $n \rightarrow \infty$

$$(1.27) \quad \frac{\mu_n^\xi(f) - \mathbb{E} \mu_n^\xi(f)}{\sqrt{\text{Var} \mu_n^\xi(f)}} \xrightarrow{\mathcal{D}} N.$$

Combining Theorem 1.11 and Theorem 1.13 yields the following theorem, which is well-suited for off-the-shelf use in applications, as seen in Section 2.3.

THEOREM 1.14. *Let (ξ, \mathcal{P}) be an admissible pair of class (A1) or (A2) such that the p -moment condition (1.19) holds for all $p \in (1, \infty)$. If $f \in \mathcal{B}(W_1)$ satisfies condition (1.26) for some $\nu \in (0, \infty)$, then $\mu_n^\xi(f)$ is asymptotically normal as in (1.27), as $n \rightarrow \infty$.*

Theorems 1.12 and 1.13 are proved in Section 4. We next compare our results with those in the literature. Point processes mentioned below are defined in Section 2.2.

Remarks.

(i) *Theorem 1.12.* In the case of Poisson and binomial input, the limits (1.23) and (1.24) are shown in [59] and [6, 55], respectively (the binomial point processes are not the restriction of an infinite point process to windows, but rather a re-scaled binomial point process on $[0, 1]^d$). In the case of Gibbsian input, the limits (1.23) and (1.24) are established in [65]. Theorem 1.12 shows these limits hold for general stationary input. The paper [70] gives a weaker version of Theorem 1.12 for specific ξ and for $f = \mathbf{1}[x \in W_1]$. In full generality, the convergence rate (1.23) is new.

(ii) *Theorems 1.13 and 1.14.* Under condition (1.26), Theorems 1.13 and 1.14 provide a central limit theorem for non-linear statistics of either α -determinantal and α -permanental input ($|\alpha|^{-1} \in \mathbb{N}$) with a fast-decaying kernel as at (2.7), the zero set \mathcal{P}_{GEF} of a Gaussian entire function, or rarified Gibbsian input. When $\xi \equiv 1$, then $\mu_n^\xi(f)$ reduces to the linear statistic $\sum_{x \in \mathcal{P}_n} f(x)$. These theorems extend the

central limit theorem for linear statistics of \mathcal{P}_{GEF} as established in [50]. When the input is determinantal with a fast decaying kernel as at (2.7), then Theorems 1.13 and 1.14 also extend the main result of Soshnikov [68], whose pathbreaking paper gives a central limit theorem for linear statistics for any determinantal input, provided the variance grows as least as fast as a power of the expectation. Proposition 5.7 of [67] shows central limit theorems for linear statistics of α -determinantal point processes with $\alpha = -1/m$ or α -permanental point processes with $\alpha = 2/m$ for some $m \in \mathbb{N}$. During the revision of this article, we noticed the recent work [53]. This paper shows that when the kernel satisfies (2.7) with $\omega(s) = o(s^{-(d+\epsilon)/2})$ and when $|\xi|$ is bounded with a deterministic radius of stabilization, then H_n^ξ at (1.2) is asymptotically normal. The generality of the score functionals and point processes considered in our article necessitates assumptions on the determinantal kernel which are more restrictive than those of [53, 68].

(iii) *Variance lower bounds.* To prove asymptotic normality it is customary to require variance lower bounds as at (1.26); [50] and [68] both require assumptions of this kind. Showing condition (1.26) is a separate problem and it fails in general. For example, the variance of the point count of some determinantal point processes, including the GUE point process, grows at most logarithmically. This phenomena is especially pronounced in dimensions $d = 1, 2$. Additionally, given input \mathcal{P}_{GEF} and $\xi \equiv 1$, the bound (1.26) may fail even when f is a smooth cutoff that equals one in a neighborhood of the origin (cf. Prop. 5.2 of [49]). On the other hand, if $\xi \equiv 1$, and if the kernel K for a determinantal point process satisfies $\int_{\mathbb{R}^d} |K(\mathbf{0}, x)|^2 dx < K(\mathbf{0}, \mathbf{0}) = \rho^{(1)}(\mathbf{0})$, then recalling the definition of $\sigma^2(\xi)$ at (1.22), we have $\sigma^2(\xi) = \sigma^2(1) = \rho^{(1)}(\mathbf{0}) - \int_{\mathbb{R}^d} |K(\mathbf{0}, x)|^2 dx > 0$. In the case of rarified Gibbsian input, the bound (1.26) holds with $\nu = 1$, as shown in of [69, Theorem 1.1]. Theorem 1.14 allows for surface-order variance growth, which arises for linear statistics $\sum_{x \in \mathcal{P}_n} \xi(x)$ of determinantal point processes; see [24, (4.15)].

(iv) *Poisson, binomial, and Gibbs input.* When \mathcal{P} is Poisson or binomial input and when ξ is a functional which stabilizes exponentially fast as at (1.15), then μ_n^ξ is asymptotically normal (1.27) under moment conditions on ξ ; see the survey [72]. When \mathcal{P} is a rarified Gibbs point process with ‘ancestor clans’ decaying exponentially fast, and when ξ is an exponentially stabilizing functional, then μ_n^ξ satisfies normal convergence (1.27), as established in [65, 69].

(v) *Mixing conditions.* Central limit theorems for geometric functionals of mixing point processes (random fields) are established in [2, 17, 34, 30, 32, 31, 53]. The geometric functionals considered in these papers are different than the ones considered here; furthermore the relation between the mixing conditions in these papers and ω -mixing correlation functions as in Definition 1.1 is unclear. Though correlation functions are simpler than mixing coefficients, which depend on σ -algebras

generated by the point processes, our decay rates appear more restrictive than those needed in afore-mentioned papers. A careful investigation of the relations between the various notions of mixing and fast decay of correlations lies beyond the scope of our limit results and will be treated in a separate paper. In the case of point processes on discrete spaces, such a study is easier, c.f. [61].

(vi) *Brillinger mixing and fast decay of correlations.* Brillinger mixing [34, Section 3.5] is defined via finiteness of integrals of the *reduced cumulant measures* (see Section 4.3.2). The very definition of Brillinger mixing implies volume-order growth of cumulants; the converse follows using the ideas in the proof of [9, Theorem 3.2]. The key to proving our announced central limit theorems is to show that the fast decay of correlations of the ξ -weighted measures (1.5) implies volume-order growth of cumulants and hence Brillinger mixing; see the remarks at the beginning of Section 4.3 and also those and at the end of Section 4.4.2.

(vii) *Multivariate central limit theorem.* We may use the Cramér-Wold device to extend Theorems 1.12 and 1.14 to the multivariate setting as follows. Let (ξ, \mathcal{P}) be a pair satisfying the hypotheses of Theorems 1.12 and 1.14. If $f_i \in \mathcal{B}(W_1)$, $1 \leq i \leq k$, satisfy the variance limit (1.24) with $\sigma^2(\xi) > 0$, then as $n \rightarrow \infty$ the fidis

$$\left(\frac{\mu_n^\xi(f_1) - \mathbb{E}\mu_n^\xi(f_1)}{\sqrt{n}}, \dots, \frac{\mu_n^\xi(f_k) - \mathbb{E}\mu_n^\xi(f_k)}{\sqrt{n}} \right)$$

converge to that of a centred Gaussian field having covariance kernel $f, g \mapsto \sigma^2(\xi) \int_{W_1} f(x)g(x)dx$.

(viii) *Deterministic radius of stabilization.* It may be shown that our main results go through without the condition (1.17) if the radius of stabilization $R^\xi(x, \mathcal{P})$ is bounded by a non-random (deterministic) constant and if (1.16) and (1.18) are satisfied. However we are unable to find any interesting examples of point processes satisfying (1.10) but not (1.17).

(ix) *Fast decay of the correlation of the ξ -weighted measures; Theorem 1.11.* Though the cumulant method is common to [6, 65, 50] and this article, a distinguishing and novel feature of our approach is the proof of fast decay of correlations of the ξ -weighted measures (1.21), and consequently their Brillinger mixing, for a wide class of functionals and point processes. As mentioned in the introduction, the proof of this result is via factorial moment expansions, which differs from the approach of [6, 65, 50] (see the remarks at the beginning of Section 3). Fast decay of correlations of the ξ -weighted measures (1.21) appears to be of independent interest. It features in the proofs of moderate deviation principles and laws of the iterated logarithms for stabilizing functionals of Poisson point process [4], [21]. Fast decay of correlations (1.21) yields volume order cumulant bounds, useful in establishing concentration inequalities as well as moderate deviations, as explained in [26, Lemma 4.2].

(x) *Normal approximation.* Difference operators (which appear in our factorial moment expansions) are also a key tool in the Malliavin-Stein method [51, 52]. This method yields presumably optimal rates of normal convergence for various statistics (including many considered in Section 2.3) in stochastic geometric problems [38, 41, 62, 37]. However, these methods currently apply only to functionals defined on Poisson and binomial point processes. It is an open question whether a refined use of these methods would yield rates of convergence in our central limit theorems.

(xi) *Cumulant bounds.* As mentioned, we establish that the k th order cumulants for $\langle f, \mu_n^\xi \rangle$ grow at most linearly in n for $k \geq 1$. Thus, under assumption (1.26), the cumulant C_n^k for $\langle f, \mu_n^\xi \rangle / \sqrt{\text{Var}\langle f, \mu_n^\xi \rangle}$ satisfies $C_n^k \leq D(k)n^{1-(\nu k/2)}$, with $D(k)$ depending only on k . For $k = 3, 4, \dots$ and $\nu > 2/3$, we have $C_n^k \leq D(k)/(\Delta(n))^{k-2}$, where $\Delta(n) := n^{(3\nu-2)/2}$. When $D(k)$ satisfies $D(k) \leq (k!)^{1+\gamma}$, γ a constant, we obtain the Berry-Esseen bound (cf. [26, Lemma 4.2])

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P} \left(\frac{\mu_n^\xi(f) - \mathbb{E}\mu_n^\xi(f)}{\sqrt{\text{Var}\mu_n^\xi(f)}} \leq t \right) - \mathbb{P}(N \leq t) \right| = O(\Delta(n)^{-1/(1+2\gamma)}).$$

Determining conditions on input pairs (ξ, \mathcal{P}) insuring the bounds $\nu > 2/3$ and $D(k) \leq (k!)^{1+\gamma}$, γ a constant, is beyond the scope of this paper. When \mathcal{P} is Poisson input, this issue is addressed by [21].

We next consider the case when the fluctuations of $H_n^\xi(\mathcal{P})$ are not of volume-order, that is to say $\sigma^2(\xi) = 0$. Though this may appear to be a degenerate condition, interesting examples involving determinantal point processes or zeros of GEF in fact satisfy $\sigma^2(1) = 0$. Such point processes are termed ‘super-homogeneous point processes’ [50, Remark 5.1]. Put

$$(1.28) \quad \widehat{H}_n^\xi(\mathcal{P}) := \sum_{x \in \mathcal{P}_n} \xi(x, \mathcal{P}).$$

The summands in $\widehat{H}_n^\xi(\mathcal{P})$, in contrast to those of $H_n^\xi(\mathcal{P})$, are not sensitive to boundary effects. We shall show that under volume-order scaling the asymptotic variance of $\widehat{H}_n^\xi(\mathcal{P})$ also equals $\sigma^2(\xi)$. However, when $\sigma^2(\xi) = 0$ we derive surface-order variance asymptotics for $\widehat{H}_n^\xi(\mathcal{P})$. Though a similar result should plausibly hold for $H_n^\xi(\mathcal{P})$, a proof seems beyond the scope of the current paper. Letting Vol_d denote the d -dimensional Lebesgue volume, for $y \in \mathbb{R}^d$ and $W \subset \mathbb{R}^d$, put

$$(1.29) \quad \gamma_W(y) := \text{Vol}_d(W \cap (\mathbb{R}^d \setminus W - y)).$$

By [44, Lemma 1(a)], we are justified in writing $\gamma(y) := \lim_{n \rightarrow \infty} \gamma_{W_n}(y)/n^{(d-1)/d}$.

5.2.III.]). By (1.12) an admissible point process having fast decay of correlations has exponential moments provided

$$(2.3) \quad \sum_{k=1}^{\infty} \frac{C_k t^k}{(k-1)!} < \infty, \quad t \in \mathbb{R}^+.$$

Note that input of type (A2) has exponential moments since by (1.16), we have $C_k = O(k^{ak})$, $a \in [0, 1)$, making (2.3) summable. For pairs (ξ, \mathcal{P}) of type (A2) with radius of stabilization bounded by $r_0 \in [1, \infty)$, by (1.18) the p -moment in (1.19) is consequently controlled by a finite exponential moment, i.e., for $x_1, \dots, x_{p'} \in W_n$,

$$(2.4) \quad \mathbb{E}_{x_1, \dots, x_{p'}} |\xi(x_1, \mathcal{P}_n)|^p \leq \mathbb{E}_{x_1, \dots, x_{p'}} (\hat{c}r_0)^{p\mathcal{P}(B_{r_0}(x_1))}.$$

Finally, if \mathcal{P} has exponential moments under its stationary probability \mathbb{P} , the same is true under $\mathbb{P}_{x_1, \dots, x_k}$ for $\alpha^{(k)}$ almost all x_1, \dots, x_k ³.

(ii) For pairs (ξ, \mathcal{P}) of type (A1), the p -moment (1.19) satisfies for $x_1, \dots, x_{p'} \in W_n$

$$(2.5) \quad \mathbb{E}_{x_1, \dots, x_{p'}} |\xi(x_1, \mathcal{P}_n)|^p \leq \left(\frac{\|h\|_{\infty}}{k} \right)^p \mathbb{E}_{x_1, \dots, x_{p'}} [(\mathcal{P}(B_r(x_1)))^{(k-1)p}].$$

We next show that (2.5) may be controlled by moments of Poisson random variables. For any Borel set $B \subset (\mathbb{R}^d)^k$, the definition of factorial moment measures gives $\alpha^{(k)}(B) \leq \kappa_k \text{Vol}_{dk}(B)$. Since moments may be expressed as a linear combination of factorial moments, for $k \in \mathbb{N}$ and a bounded Borel subset $B \subset \mathbb{R}^d$, using (1.8) we have

$$(2.6) \quad \mathbb{E}[(\mathcal{P}(B))^k] = \sum_{j=0}^k \left\{ \begin{matrix} k \\ j \end{matrix} \right\} \alpha^{(j)}(B^j) \leq \kappa_k \sum_{j=0}^k \left\{ \begin{matrix} k \\ j \end{matrix} \right\} \text{Vol}_{jd}(B)^j = \kappa_k \mathbb{E}(\text{Po}(\text{Vol}_d(B))^k),$$

where $\left\{ \begin{matrix} k \\ j \end{matrix} \right\}$ stand for the *Stirling numbers of the second kind*, $\text{Po}(\lambda)$ denotes a Poisson random variable with mean λ and where κ_j 's are non-decreasing in j . Thus by (1.12), an admissible point process having fast decay of correlations has all moments, as in (2.2). If \mathcal{P} has all moments under its stationary probability \mathbb{P} , the same is true under $\mathbb{P}_{x_1, \dots, x_k}$ for $\alpha^{(k)}$ almost all x_1, \dots, x_k (by the same arguments as in Footnote 3).

³ Indeed, if $\mathbb{E}_{x_1, \dots, x_k} [\rho^{\mathcal{P}(B_r(x_1))}] = \infty$ for $x_1, \dots, x_k \in B'$ for some bounded $B' \in \mathbb{R}^d$ such that $\alpha^{(k)}(B'^k) > 0$ then $\mathbb{E}_{x_1, \dots, x_k} [\rho^{\mathcal{P}(B_r(x_1))}] \leq \mathbb{E}_{x_1, \dots, x_k} [\rho^{\mathcal{P}(B'_r)}] = \infty$ with $B'_r = B' \oplus B_r(\mathbf{0}) = \{y' + y : y' \in B', y \in B_r(\mathbf{0})\}$ the r -parallel set of B' . Integrating with respect to $\alpha^{(k)}$ in B'^k , by the Campbell formula $\mathbb{E}[(\mathcal{P}(B'_r))^k \rho^{\mathcal{P}(B'_r)}] = \infty$, which contradicts the existence of exponential moments under \mathbb{P} .

locally trace class, a point process \mathcal{P} is said to be α -*permanental*⁵ if its correlation functions satisfy

$$(2.9) \quad \rho^{(k)}(x_1, \dots, x_k) = \sum_{\pi \in S_k} \alpha^{k-\nu(\pi)} \prod_{i=1}^k K(x_i, x_{\pi(i)})$$

where S_k stands for the usual symmetric group and $\nu(\cdot)$ denotes the number of cycles in a permutation. The right hand side is the α -permanent of the matrix $((K(x_i, x_j))_{i,j \leq k})$. The special cases $\alpha = 0$ and $\alpha = 1$ respectively give the Poisson point process with intensity $K(\mathbf{0}, \mathbf{0})$ and the permanental point process with kernel K . In what follows, we assume $\alpha = 1/m$ for $m \in \mathbb{N}$, i.e. $1/\alpha$ is a positive integer. Existence of such α -permanental point processes is guaranteed by [67, Theorem 1.2]. The property of these point processes most important to us is that an α -permanental point process with kernel K is a superposition of $1/\alpha$ i.i.d. copies of a permanental point process with kernel αK (see [8, Section 4.10]). Also from definition (2.9), we obtain

$$\rho^{(k)}(x_1, \dots, x_k) \leq \|K\|^k \alpha^k \sum_{\pi \in S_k} (\alpha^{-1})^{\nu(\pi)},$$

and so we can take $\kappa_k = \prod_{i=0}^{k-1} (j\alpha + 1) \|K\|^k$ for an α -permanental point process. The following result is a consequence of the upcoming Proposition 2.3 and the identity (2.8) for decay constants of a permanental point process with kernel αK .

PROPOSITION 2.1. *Let $\alpha = 1/m$ for some $m \in \mathbb{N}$ and let \mathcal{P}_α be the stationary α -permanental point process with a kernel K which is Hermitian, non-negative definite and locally trace class. Assume also that $|K(x, y)| \leq \omega(|x - y|)$ for some fast-decreasing ω . Then \mathcal{P}_α is an admissible point process having fast decay of correlations with correlation decay constants $C_k = km^{1-k(m-1)} m! (k!)^m \|K\|^{km-1}$, $c_k = 1$ and decay function $\phi = \omega$.*

Zero set of Gaussian entire function (GEF). A Gaussian entire function $f(z)$ is the sum $\sum_{j \geq 0} X_j \frac{z^j}{\sqrt{j!}}$, where X_j are i.i.d. with the standard normal density on the complex plane. The zero set $f^{-1}(\{0\})$ gives rise to the point process $\mathcal{P}_{GEF} := \sum_{x \in f^{-1}(\{0\})} \delta_x$ on \mathbb{R}^2 . The point process \mathcal{P}_{GEF} is an admissible point process having fast decay of correlations [50, Theorem 1.4] and exhibits local repulsion of points. Though \mathcal{P}_{GEF} satisfies condition (1.17), it is unclear whether (1.16) holds. By [36, Theorem 1], $\mathcal{P}_{GEF}(B_r(\mathbf{0}))$ has exponential moments.

Moment conditions. For $p \in [1, \infty)$, we show that the p -moment condition (1.19)

⁵In contrast to terminology in [8, 67], here we distinguish the two cases (i) $\alpha \geq 0$ (α -permanental) and (ii) $\alpha \leq 0$ (α -determinantal)

stationary determinantal point process has a fast-decreasing kernel as at (2.7), then [14, Lemma 1.3] in the supplemental file shows that it is an admissible point process having fast decay of correlations satisfying (1.16) with decay function $\phi = \omega$, with ω as at (2.7), and correlation decay constants

$$(2.15) \quad C_k := k^{1+(k/2)} \|K\|^{k-1}, c_k \equiv 1.$$

Consequently, ϕ satisfies the requisite exponential decay (1.17) whenever ω itself satisfies (1.17).

The Ginibre ensemble of eigenvalues of $N \times N$ matrices with independent standard complex Gaussian entries is a leading example of a determinantal point process. The limit of the Ginibre ensemble as $N \rightarrow \infty$ is the Ginibre point process (or the infinite Ginibre ensemble), here denoted \mathcal{P}_{GIN} . It is the prototype of a stationary determinantal point process and has the following kernel : For $z_1, z_2 \in \mathbb{C}$,

$$K(z_1, z_2) := \exp(z_1 \bar{z}_2) \exp\left(-\frac{|z_1|^2 + |z_2|^2}{2}\right) = \exp\left(i \operatorname{Im}(z_1 \bar{z}_2) - \frac{|z_1 - z_2|^2}{2}\right).$$

More generally, for $0 < \beta \leq 1$, the β -Ginibre (determinantal) point process (see [27]) has kernel

$$K_\beta(z_1, z_2) := \exp\left(\frac{1}{\beta} z_1 \bar{z}_2\right) \exp\left(-\frac{|z_1|^2 + |z_2|^2}{2\beta}\right), \quad z_1, z_2 \in \mathbb{C}.$$

When $\beta = 1$, we obtain \mathcal{P}_{GIN} and as $\beta \rightarrow 0$ we obtain the Poisson point process. Thus the β -Ginibre point process interpolates between the Ginibre and Poisson point processes. Identifying the complex plane with \mathbb{R}^2 we see that all β -Ginibre point processes are admissible point processes having fast decay of correlations satisfying (1.16) and (1.17).

Moment Conditions. Let $p \in [1, \infty)$ and let \mathcal{P} be a stationary determinantal point process with a continuous and fast-decreasing kernel. We now show that the p -moment condition (1.19) holds for pairs (ξ, \mathcal{P}) of class (A1) or (A2), provided ξ has a deterministic radius of stabilization, say $r_0 \in [1, \infty)$. First, for all $(x_1, \dots, x_p) \in (\mathbb{R}^d)^p$, all increasing $F : \mathbb{N} \rightarrow \mathbb{R}^+$ and all bounded Borel sets B we have [27, Theorem 2]

$$\mathbb{E}_{x_1, \dots, x_p}^! (F(\mathcal{P}(B))) \leq \mathbb{E}(F(\mathcal{P}(B))).$$

Thus using (2.4), the above inequality and stationarity of \mathcal{P} , we get that for any bounded stabilizing score function ξ of class (A2),

$$(2.16) \quad \begin{aligned} & \sup_{1 \leq n \leq \infty} \sup_{1 \leq p' \leq [p]} \sup_{x_1, \dots, x_{p'} \in W_n} \mathbb{E}_{x_1, \dots, x_{p'}} |\xi(x_1, \mathcal{P}_n)|^p \\ & \leq \mathbb{E}(\hat{c}r_0)^{p\mathcal{P}(B_{r_0}(\mathbf{0})) + p^2} < \infty. \end{aligned}$$

2.2.3. *Additional input examples.* For additional examples of admissible point processes having fast decay of correlations, we refer to the arxiv version of this paper [15, Section 2.3]. We shall discuss but one example here.

Superpositions of i.i.d. point processes. A natural operation on point processes generating new point processes consists of independent superposition. We show that this operation preserves fast decay of correlations.

Let $\mathcal{P}_1, \dots, \mathcal{P}_m, m \in \mathbb{N}$, be i.i.d. copies of an admissible point process \mathcal{P} with correlation functions ρ and having fast decay of correlations. Let ρ_0 denote the correlation functions of the point process $\mathcal{P}_0 := \cup_{i=1}^m \mathcal{P}_i$. For any $k \geq 1$ and distinct $x_1, \dots, x_k \in \mathbb{R}^d$ the following relation holds

$$(2.18) \quad \rho_0^{(k)}(x_1, \dots, x_k) = \sum_{\sqcup_{i=1}^m S_i = [k]} \prod_{i=1}^m \rho(S_i),$$

where \sqcup stands for disjoint union and where we abbreviate $\rho^{(|S_i|)}(x_j : j \in S_i)$ by $\rho(S_i)$. Here S_i may be empty, in which case we set $\rho(\emptyset) = 1$. From (2.18), we have that \mathcal{P}_0 is an admissible point process with intensity $m\rho^{(1)}(\mathbf{0})$. Further, we take $\kappa_k(\mathcal{P}_0) = (\kappa_k)^m m^k$. The proof of the next proposition, which shows that \mathcal{P}_0 has fast decay of correlations, is in the supplemental file (cf. [14, Proposition 1.8]).

PROPOSITION 2.3. *Let $m \in \mathbb{N}$ and $\mathcal{P}_1, \dots, \mathcal{P}_m$ be i.i.d. copies of an admissible point process \mathcal{P} having fast decay of correlations with decay function ϕ and correlation decay constants C_k and c_k . Then $\mathcal{P}_0 := \cup_{i=1}^m \mathcal{P}_i$ is an admissible point process having fast decay of correlations with decay function ϕ and correlation decay constants $m^k m! (\kappa_k)^{m-1} C_k$ and c_k . Further, if \mathcal{P} is admissible input of type (A2) with $\kappa_k \leq \lambda^k$ for some $\lambda \in (0, \infty)$, then \mathcal{P}_0 is also admissible input of type (A2).*

We have already used this proposition in the context of fast decay of correlations of α -permanental and determinantal point processes.

2.3. Applications. Having provided examples of admissible point processes, one may use Theorems 1.12 and 1.14 to deduce the limit theory for geometric and topological statistics of these point processes. Examples include statistics arising in combinatorial and differential topology, integral geometry, and computational geometry. As fully explained in Section 2.3 of [15], one may deduce expectation and variance asymptotics and central limit theorems for statistics of random Čech complexes, Morse critical points, as well as statistics of germ-grain models generated by admissible point processes. The results described in Section 2.3 of [15] are not exhaustive and include functionals in stochastic geometry already discussed in e.g. [6, 60]. There are further applications to (i) random packing models on input hav-

Edge-lengths of k -nearest neighbour graphs. Statistics of the Voronoi tessellation as well as of graphs in computational geometry such as the k -nearest neighbors graph and sphere of influence graph may be expressed as sums of exponentially stabilizing score functionals [57] and hence via Theorems 1.12 and 1.14, we may deduce the limit theory for these statistics. To illustrate, we establish a weak law of large numbers, variance asymptotics, and a central limit theorem for the total edge-length of the k -nearest neighbors graph on a α -determinantal point process $\mathcal{P} := \mathcal{P}_\alpha$ with $-1/\alpha \in \mathbb{N}$ and a fast-decreasing kernel as in (2.7). As noted in Proposition 2.2, such an α -determinantal point process is of class (A2) as in Definition 1.7.

As shown in [14, Corollary 1.10] of the supplemental file, we may explicitly upper bound void probabilities for \mathcal{P} , allowing us to deduce exponential stabilization for score functions on \mathcal{P} . This is a recurring phenomena, and it is often the case that to show exponential stabilization of statistics, it suffices to control the Palm probability content of large Euclidean balls. This opens the way towards showing that other relevant statistics of random graphs exhibit exponential stabilization on \mathcal{P} . This includes intrinsic volumes of faces of Voronoi tessellations [64, Section 10.2], edge-lengths in a radial spanning tree [66, Lemma 3.2], proximity graphs including the Gabriel graph, and Morse critical points.

Given locally finite $\mathcal{X} \subset \mathbb{R}^d$ and $k \in \mathbb{N}$, the (undirected) k -nearest neighbors graph $NG(\mathcal{X})$ is the graph with vertex set \mathcal{X} obtained by including an edge $\{x, y\}$ if y is one of the k nearest neighbors of x and/or x is one of the k nearest neighbors of y . In the case of a tie we may break the tie via some pre-defined total order (say lexicographic order) on \mathbb{R}^d . For any finite $\mathcal{X} \subset \mathbb{R}^d$ and $x \in \mathcal{X}$, we let $\mathcal{E}(x)$ be the edges e in $NG(\mathcal{X})$ which are incident to x . Defining

$$\xi_L(x, \mathcal{X}) := \frac{1}{2} \sum_{e \in \mathcal{E}(x)} |e|,$$

we write the total edge length of $NG(\mathcal{X})$ as $L(NG(\mathcal{X})) = \sum_{x \in \mathcal{X}} \xi_L(x, \mathcal{X})$. Let $\sigma^2(\xi_L)$ be as at (1.22), with ξ put to be ξ_L .

THEOREM 2.5. *Let $\mathcal{P} := \mathcal{P}_\alpha$ be a stationary α -determinantal point process on \mathbb{R}^d with $-1/\alpha \in \mathbb{N}$ and a fast-decreasing kernel K as at (2.7). We have*

$$\left| \frac{\mathbb{E}L(NG(\mathcal{P}_n))}{n} - \mathbb{E}_0 \xi_L(\mathbf{0}, \mathcal{P}) K(\mathbf{0}, \mathbf{0}) \right| = O(n^{-1/d})$$

and

$$\lim_{n \rightarrow \infty} \frac{\text{Var}L(NG(\mathcal{P}_n))}{n} = \sigma^2(\xi_L).$$

expectation of functionals of point processes. Notice that (1.21) holds for any exponentially stabilizing score function ξ satisfying the p -moment condition (1.19) for all $p \in [1, \infty)$ on a Poisson point process \mathcal{P} . Indeed if $x, y \in \mathbb{R}^d$ and $r_1, r_2 > 0$ satisfy $r_1 + r_2 < |x - y|$ then $\xi(x, \mathcal{P})\mathbf{1}[R^\xi(x, \mathcal{P}) \leq r_1]$ and $\xi(y, \mathcal{P})\mathbf{1}[R^\xi(y, \mathcal{P}) \leq r_2]$ are independent random variables. This yields the fast decay (1.21) with $k_1 = \dots = k_{p+q} = 1$ and $\tilde{C}_n \leq c_1^n$ with c_1 a constant, as in [6, Lemma 5.2]. On the other hand, if \mathcal{P} is rarified Gibbsian input and ξ is exponentially stabilizing, then [65, Lemma 3.4] shows the fast decay bound (1.21) with $k_1 = \dots = k_{p+q} = 1$. These methods depend on quantifying the region of spatial dependencies of Gibbsian points via exponentially decaying diameters of their ancestor clans. Such methods apparently neither extend to determinantal input nor to the zero set \mathcal{P}_{GEF} of a Gaussian entire function. On the other hand, for \mathcal{P}_{GEF} and for $\xi \equiv 1$, the paper [50] uses the Kac-Rice-Hammersley formula and complex analysis tools to show (1.21) with $k_1 = \dots = k_{p+q} = 1$. All three proofs are specific to either the underlying point process or to the score function ξ . The following more general and considerably different approach includes these results as special cases.

3.1. Difference operators and factorial moment expansions. We introduce some notation and collect auxiliary results required for an application of the much-needed factorial moment expansions [11, 12] for general point processes. Equip \mathbb{R}^d with a total order \prec defined using the lexicographical ordering of the polar coordinates. For $\mu \in \mathcal{N}$ and $x \in \mathbb{R}^d$, define the measure $\mu|_x(\cdot) := \mu(\cdot \cap \{y : y \prec x\})$. Note that since μ is a locally finite measure and the ordering is defined via polar coordinates $\mu|_x$ is a finite measure for all $x \in \mathbb{R}^d$. Let o denote the null-measure i.e., $o(B) = 0$ for all Borel subsets B of \mathbb{R}^d . For a measurable function $\psi : \mathcal{N} \rightarrow \mathbb{R}$, $l \in \mathbb{N} \cup \{0\}$, and $x_1, \dots, x_l \in \mathbb{R}^d$, we define the factorial moment expansion (FME) kernels [11, 12] as follows. For $l \geq 1$,

(3.1)

$$D_{x_1, \dots, x_l}^l \psi(\mu) = \sum_{i=0}^l (-1)^{l-i} \sum_{J \subset \binom{[l]}{i}} \psi(\mu|_{x_*} + \sum_{j \in J} \delta_{x_j}) = \sum_{J \subset [l]} (-1)^{l-|J|} \psi(\mu|_{x_*} + \sum_{j \in J} \delta_{x_j}),$$

where $\binom{[l]}{j}$ denotes the collection of all subsets of $[l] := \{1, \dots, l\}$ with cardinality j and $x_* := \min\{x_1, \dots, x_l\}$, with the minimum taken with respect to the order \prec . For $l = 0$, put $D^0 \psi(\mu) := \psi(o)$. Note that $D_{x_1, \dots, x_l}^{(l)} \psi(\mu)$ is a symmetric function of x_1, \dots, x_l .⁶

We say that ψ is \prec -continuous at ∞ if for all $\mu \in \mathcal{N}$ we have

$$\lim_{x \uparrow \infty} \psi(\mu|_x) = \psi(\mu).$$

⁶For $x_l \prec x_{l-1} \prec \dots \prec x_1$ the functional $D_{x_1, \dots, x_l}^l \psi(\mu)$ is equal to the iterated difference operator: $D_{x_1}^1 \psi(\mu) = \psi(\mu|_{x_1} + \delta_{x_1}) - \psi(\mu|_{x_1})$, $D_{x_1, \dots, x_l}^l \psi(\mu) = D_{x_l}^1 (D_{x_1, \dots, x_{l-1}}^{l-1} \psi(\mu))$.

$(1, \infty)$ and with \mathcal{P} having exponential moments. Then for distinct $x_1, \dots, x_p \in \mathbb{R}^d$, non-negative integers k_1, \dots, k_p and $n \leq \infty$ the functional ψ^\dagger at (3.6) admits the FME

$$\begin{aligned} \mathbb{E}_{x_1, \dots, x_p} [\psi_{k_1, \dots, k_p}(x_1, \dots, x_p; \mathcal{P}_n)] &= \mathbb{E}_{x_1, \dots, x_p}^\dagger [\psi_{k_1, \dots, k_p}^\dagger(x_1, \dots, x_p; \mathcal{P}_n)] \\ &= \psi_{k_1, \dots, k_p}^\dagger(x_1, \dots, x_p; o) \\ (3.7) \quad &+ \sum_{l=1}^{\infty} \frac{1}{l!} \int_{\mathbb{R}^{dl}} D_{y_1, \dots, y_l}^l \psi_{k_1, \dots, k_p}^\dagger(x_1, \dots, x_p; o) \rho_{x_1, \dots, x_p}^{(l)}(y_1, \dots, y_l) dy_1 \dots dy_l. \end{aligned}$$

When (ξ, \mathcal{P}) is of type (A1), the series (3.7) has at most $(k-1) \sum_{i=1}^p k_i$ non-zero terms, where k is as in (1.13).

PROOF. Throughout we fix non-negative integers k_1, \dots, k_p and suppress them when writing ψ^\dagger ; i.e., $\psi^\dagger(x_1, \dots, x_p; \mathcal{P}_n) := \psi_{k_1, \dots, k_p}^\dagger(x_1, \dots, x_p; \mathcal{P}_n)$. The bounded radius of stabilization for ξ implies ψ^\dagger is \prec -continuous at ∞ .

Consider first ψ^\dagger at (3.6) with ξ as in case (ii); later we consider the simpler case (i). We show the validity of the expansion (3.7) as follows. Let $y_1, \dots, y_l \in \mathbb{R}^d$. The difference operator D_{y_1, \dots, y_l}^l vanishes as soon as $y_k \notin \cup_{i=1}^p B_r(x_i)$ for some $k \in \{1, \dots, l\}$, that is to say

$$(3.8) \quad D_{y_1, \dots, y_l}^l \psi^\dagger(x_1, \dots, x_p; \mu) = 0.$$

To prove this, set $\mu_J := \mu|_{y_*} + \sum_{j \in J} \delta_{y_j}$ for $J \subset [l]$ and $y_* := \min\{y_1, \dots, y_l\}$, with the minimum taken with respect to \prec order. From (3.1) we obtain

$$\begin{aligned} D_{y_1, \dots, y_l}^l \psi^\dagger(x_1, \dots, x_p; \mu) &= \sum_{J \subset [l], k \notin J} (-1)^{l-|J|} \psi^\dagger(x_1, \dots, x_p; \mu_J) \\ &+ \sum_{J \subset [l], k \notin J} (-1)^{l-|J|-1} \psi^\dagger(x_1, \dots, x_p; \mu_{J \cup \{k\}}) = 0, \end{aligned}$$

where the last equality follows by noting that for $J \subset [l]$ with $k \notin J$, $\psi^\dagger(x_1, \dots, x_p; \mu_J) = \psi^\dagger(x_1, \dots, x_p; \mu_{J \cup \{k\}})$ because $R^\xi(x, \mathcal{P}) \in [1, r]$ by assumption.

Henceforth we put

$$(3.9) \quad K_p := \sum_{i=1}^p k_i, \quad K_q := \sum_{i=1}^q k_{p+i}, \quad K := \sum_{i=1}^{p+q} k_i.$$

Consider now $y_1, \dots, y_l \in \cup_{i=1}^p B_r(x_i)$. For $J \subset [l]$, from $1 \leq R^\xi(x, \mathcal{P}) \leq r$ and (1.18) we have

$l \in (K_p(k-1), \infty)$ we have

$$(3.13) \quad D_{y_1, \dots, y_l}^l \psi^l(x_1, \dots, x_p; \mu) = 0 \quad \forall y_1, \dots, y_l \in \mathbb{R}^d,$$

as shown in [62, Lemma 3.3] for Poisson point processes (the proof for general simple counting measures μ is identical). This implies that conditions (3.2) for $l \in (K_p(k-1), \infty)$ and (3.3) are trivially satisfied for ψ^l as at (3.6). Now, we need to verify the condition (3.2) for $l \in [1, K_p(k-1)]$. For $y_1, \dots, y_l \in \mathbb{R}^d$, set as before $\mu_J = \mu|_{y_*} + \sum_{j \in J} \delta_{y_j}$ for $J \subset [l]$ and $y_* := \min\{y_1, \dots, y_l\}$, with the minimum taken with respect to the order \prec . Since ξ has a bounded stabilization radius, by (3.8) and (2.5), we have

$$(3.14) \quad \begin{aligned} \psi^l(x_1, \dots, x_p; \mu_J) &\leq \prod_{i=1}^p \|h\|_\infty^{k_i} (\mu(\cup_{i=1}^p B_r(x_i)) + |J| + p)^{k_i(k-1)} \\ &\leq \|h\|_\infty^{K_p} (\mu(\cup_{i=1}^p B_r(x_i)) + |J| + p)^{K_p(k-1)}. \end{aligned}$$

The number of subsets of $[l]$ is 2^l and so by (3.1), we obtain

$$(3.15) \quad \begin{aligned} |D_{y_1, \dots, y_l}^l \psi^l(x_1, \dots, x_p; \mu)| &\leq \|h\|_\infty^{K_p} \sum_{J \subset [l]} (\mu(\cup_{i=1}^p B_r(x_i)) + |J| + p)^{K_p(k-1)} \\ &\leq \|h\|_\infty^{K_p} 2^l (\mu(\cup_{i=1}^p B_r(x_i)) + l + p)^{K_p(k-1)}. \end{aligned}$$

Consider $\psi^l(x_1, \dots, x_p; \mathcal{P}_n)$ with ψ^l defined as above. Using the refined Campbell theorem (1.9), the bound (3.15), and following the calculations as in (3.12), we obtain

$$\begin{aligned} &\frac{1}{l!} \int_{\mathbb{R}^{dl}} (\mathbb{E}_{x_1, \dots, x_p}^!)_{y_1, \dots, y_l} [|D_{y_1, \dots, y_l}^l \psi^l(x_1, \dots, x_p; \mathcal{P}_n)| \rho_{x_1, \dots, x_p}^{(l)}(y_1, \dots, y_l)] dy_1 \dots dy_l \\ &\leq \|h\|_\infty^{K_p} 2^l \mathbb{E}_{x_1, \dots, x_p} [\mathcal{P}(\cup_{i=1}^p B_r(x_i))^l (\mathcal{P}(\cup_{i=1}^p B_r(x_i)) + l + p)^{K_p(k-1)}]. \end{aligned}$$

Since \mathcal{P} has all moments under the Palm measure (see Remark (ii) at the beginning of Section 2.1), the finiteness of the last term and hence the validity of the condition (3.2) for $l \in [1, K_p(k-1)]$ follows. This justifies the FME expansion (3.7), with finitely many non-zero terms, when ψ^l is the product of score functions of class (A1). \square

3.2. Proof of Theorem 1.11. First assume that (ξ, \mathcal{P}) is of class (A2). Later we consider the simpler case that (ξ, \mathcal{P}) is of class (A1). For fixed $p, q, k_1, \dots, k_{p+q} \in \mathbb{N}$, consider correlation functions $m^{(k_1, \dots, k_{p+q})}(x_1, \dots, x_{p+q}; n)$, $m^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n)$, and $m^{(k_{p+1}, \dots, k_{p+q})}(x_{p+1}, \dots, x_{p+q}; n)$ of the ξ -weighted measures at (1.6). We abbreviate $\psi_{k_1, \dots, k_p}(x_1, \dots, x_p; \mu)$ by $\psi(x_1, \dots, x_p; \mu)$ as at (3.5), and similarly for

For any reals $A, B, \tilde{A}, \tilde{B}$, with $|\tilde{B}| \leq |B|$ we have $|AB - \tilde{A}\tilde{B}| \leq |A(B - \tilde{B})| + |(A - \tilde{A})\tilde{B}| \leq (|A| + |B|)(|B - \tilde{B}| + |A - \tilde{A}|)$. Hence, it follows that

$$\begin{aligned} & |m^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n) m^{(k_{p+1}, \dots, k_q)}(x_{p+1}, \dots, x_{p+q}; n) \\ & - \tilde{m}^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n) \tilde{m}^{(k_{p+1}, \dots, k_q)}(x_{p+1}, \dots, x_{p+q}; n)| \\ & \leq (c_1(K_p) + c_1(K_q)) \left(c_1(K_p) \varphi(a_{K_p} t)^{1/(K_p+1)} + c_1(K_q) \varphi(a_{K_q} t)^{1/(K_q+1)} \right) \\ & \leq c_2(K) \varphi(a_K t)^{1/(K+1)}, \end{aligned}$$

with $c_2(m) := 4(c_1(m))^2$ and where we note that $\varphi(a_m t)^{1/(m+1)}$ is also fast-decreasing by (1.15). The difference of correlation functions of the ξ -weighted measures is thus bounded by

$$\begin{aligned} & \left| m^{(k_1, \dots, k_{p+q})}(x_1, \dots, x_{p+q}; n) - m^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n) \right. \\ & \quad \left. \times m^{(k_{p+1}, \dots, k_{p+q})}(x_{p+1}, \dots, x_{p+q}; n) \right| \\ & \leq (c_1(K) + c_2(K)) \varphi(a_K t)^{1/(K+1)} + \left| \tilde{m}^{(k_1, \dots, k_{p+q})}(x_1, \dots, x_{p+q}; n) \right. \\ (3.19) \quad & \left. - \tilde{m}^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n) \tilde{m}^{(k_{p+1}, \dots, k_{p+q})}(x_{p+1}, \dots, x_{p+q}; n) \right|. \end{aligned}$$

The rest of the proof consists of bounding $|\tilde{m}^{(k_1, \dots, k_{p+q})} - \tilde{m}^{(k_1, \dots, k_p)} \tilde{m}^{(k_{p+1}, \dots, k_{p+q})}|$ by a fast-decreasing function of s . In this regard we will consider the expansion (3.7) with $\psi(x_1, \dots, x_p; \mathcal{P}_n)$ replaced by $\tilde{\psi}(x_1, \dots, x_p; \mathcal{P}_n)$ as at (3.17) and similarly for $\tilde{\psi}(x_{p+1}, \dots, x_{p+q}; \mathcal{P}_n)$ and $\tilde{\psi}(x_1, \dots, x_{p+q}; \mathcal{P}_n)$. By [14, Lemma 1.2] in the supplemental file, $\tilde{\xi}(x_i, \mathcal{P}_n)$, $1 \leq i \leq p$, have radii of stabilization bounded above by t and also satisfy the power-growth condition (1.18) since $|\tilde{\xi}| \leq |\xi|$. Thus the pair $(\tilde{\xi}, \mathcal{P})$ satisfies the assumptions of Lemma 3.2. The corresponding version of $\tilde{\psi}$, accounting for the fixed atoms of \mathcal{P}_n is

$$\tilde{\psi}^1(x_1, \dots, x_p; \mu) := \prod_{i=1}^p \tilde{\xi}(x_i, \mu + \sum_{i=1}^p \delta_{x_i})^{k_i}$$

and similarly for $\tilde{\psi}^1(x_{p+1}, \dots, x_q; \mathcal{P}_n)$ and $\tilde{\psi}^1(x_1, \dots, x_{p+q}; \mathcal{P}_n)$.

Put $B_{t,n}(x_i) := B_t(x_i) \cap W_n$. Applying (3.7), the multiplicative identity (2.11)

$\mathbb{E}_{x_1, \dots, x_p}^! [\psi^!(\mathcal{P}_n)]$ (cf. (3.6)), we obtain

$$\begin{aligned}
& \tilde{m}^{(k_1, \dots, k_p)}(x_1, \dots, x_p) \tilde{m}^{(k_{p+1}, \dots, k_q)}(x_{p+1}, \dots, x_{p+q}) \\
&= \mathbb{E}_{x_1, \dots, x_p}^! [\tilde{\psi}^!(x_1, \dots, x_p; \mathcal{P}_n)] \mathbb{E}_{x_{p+1}, \dots, x_{p+q}}^! [\tilde{\psi}^!(x_{p+1}, \dots, x_{p+q}; \mathcal{P}_n)] \\
& \quad \times \rho^{(p)}(x_1, \dots, x_p) \rho^{(q)}(x_{p+1}, \dots, x_{p+q}) \\
&= \sum_{l_1, l_2=0}^{\infty} \frac{1}{l_1! l_2!} \int_{(\cup_{i=1}^{l_1} B_{t,n}(x_i))^{l_1} \times (\cup_{i=1}^{l_2} B_{t,n}(x_{p+i}))^{l_2}} D_{y_1, \dots, y_{l_1}}^{l_1} \tilde{\psi}^!(x_1, \dots, x_p; o) \\
& \quad \times D_{z_1, \dots, z_{l_2}}^{l_2} \tilde{\psi}^!(x_{p+1}, \dots, x_{p+q}; o) \rho^{(l_1+p)}(x_1, \dots, x_p, y_1, \dots, y_{l_1}) \\
(3.22) \quad & \times \rho^{(l_2+q)}(x_{p+1}, \dots, x_{p+q}, z_1, \dots, z_{l_2}) dy_1 \dots dy_{l_1} dz_1 \dots dz_{l_2}.
\end{aligned}$$

Applying (3.1) once more for μ the null measure, this gives

$$\begin{aligned}
& \tilde{m}^{(k_1, \dots, k_p)}(x_1, \dots, x_p) \tilde{m}^{(k_{p+1}, \dots, k_q)}(x_{p+1}, \dots, x_{p+q}) \\
&= \sum_{l_1, l_2=0}^{\infty} \frac{1}{l_1! l_2!} \int_{(\cup_{i=1}^{l_1} B_{t,n}(x_i))^{l_1} \times (\cup_{i=1}^{l_2} B_{t,n}(x_{p+i}))^{l_2}} dy_1 \dots dy_{l_1} dz_1 \dots dz_{l_2} \\
& \quad \times \sum_{J_1 \subset [l_1], J_2 \subset [l_2]} (-1)^{l_1+l_2-|J_1|-|J_2|} \tilde{\psi}^!(x_1, \dots, x_p; \sum_{i \in J_1} \delta_{y_i}) \tilde{\psi}^!(x_{p+1}, \dots, x_{p+q}; \sum_{i \in J_2} \delta_{z_i}) \\
& \quad \times \rho^{(l_1+p)}(x_1, \dots, x_p, y_1, \dots, y_{l_1}) \rho^{(l_2+q)}(x_{p+1}, \dots, x_{p+q}, z_1, \dots, z_{l_2}) \\
&= \sum_{l=0}^{\infty} \sum_{j=0}^l \frac{1}{j!(l-j)!} \int_{(\cup_{i=1}^j B_{t,n}(x_i))^j \times (\cup_{i=1}^{l-j} B_{t,n}(x_{p+i}))^{l-j}} \sum_{J_1 \subset [j], J_2 \subset [l] \setminus [j]} (-1)^{l-|J_1|-|J_2|} \\
& \quad \times \tilde{\psi}^!(x_1, \dots, x_p; \sum_{i \in J_1} \delta_{y_i}) \tilde{\psi}^!(x_{p+1}, \dots, x_{p+q}; \sum_{i \in J_2} \delta_{y_i}) \\
& \quad \times \rho^{(j+p)}(x_1, \dots, x_p, y_1, \dots, y_j) \rho^{(l-j+q)}(x_{p+1}, \dots, x_{p+q}, y_{j+1}, \dots, y_l) dy_1 \dots dy_l \\
&= \sum_{l=0}^{\infty} \sum_{j=0}^l \frac{1}{j!(l-j)!} \int_{(\cup_{i=1}^j B_{t,n}(x_i))^j \times (\cup_{i=1}^{l-j} B_{t,n}(x_{p+i}))^{l-j}} \sum_{J \subset [l]} (-1)^{l-|J|} \tilde{\psi}^!(x_1, \dots, x_{p+q}; \sum_{i \in J} \delta_{y_i}) \\
(3.23) \quad & \times \rho^{(j+p)}(x_1, \dots, x_p, y_1, \dots, y_j) \rho^{(l-j+q)}(x_{p+1}, \dots, x_{p+q}, y_{j+1}, \dots, y_l) dy_1 \dots dy_l,
\end{aligned}$$

where we have used (3.21) in the last equality.

Now we estimate the difference of (3.20) and (3.23). Applying (1.10) and re-

where for $r \in \mathbb{R}$, $\lfloor r \rfloor$ is the greatest integer less than r . We compute

$$(3.27) \quad t^K \sum_{l=0}^{\infty} \frac{C_{l+K}}{l!} 4^l (\hat{c}t)^{lK} (K\theta_d t^d)^l \leq t^K \sum_{n=0}^{\infty} \sum_{\{l: \lfloor l(1-a) \rfloor = n\}} \frac{c_4 c_5^l l^{c_6} (t^{K+d})^l}{n!}$$

$$\leq t^K \sum_{n=0}^{\infty} \frac{c_4 c_5^n n^{c_6} (t^{K+d})^{(n+1)/(1-a)}}{(1-a)n!} \leq c_7 \exp(c_8 t^{(K+d)/(1-a)})$$

where c_7 and c_8 depend only on a , d and K .

Recalling from (3.16) that $t := (s/4)^{b(1-a)/(2(K+d))}$ we obtain

$$\sum_{l=0}^{\infty} \frac{C_{l+K}}{l!} 4^l (\hat{c}t)^{(l+1)K} (K\theta_d t^d)^l \leq c_7 \exp\left(c_8 \left(\frac{s}{4}\right)^{\frac{b}{2}}\right).$$

By (1.17), there is a constant c_9 depending only on a such that for all s we have $\phi(s) \leq c_9 \exp(-s^b/c_9)$. Combining this with (3.26) and (3.27) gives

$$|\tilde{m}^{(k_1, \dots, k_{p+q})} - \tilde{m}^{(k_1, \dots, k_p)} \tilde{m}^{(k_{p+1}, \dots, k_{p+q})}| \leq c_7 c_9 \exp\left(\frac{-(s/2)^b}{c_9} + c_8 \left(\frac{s}{4}\right)^{\frac{b}{2}}\right).$$

This along with (3.19) shows (1.21) when (ξ, \mathcal{P}) is an admissible pair of class (A2).

Now we establish (1.21) when (ξ, \mathcal{P}) is of class (A1). Let k be as in (1.13). Follow the arguments for case (A2) word for word using that $\sup_{x \in \mathcal{P}} R^\xi(x, \mathcal{P}) \leq r$. Notice that for $l \in ((k-1)K, \infty)$ the summands in (3.20) vanish. Likewise, when $l_1 \in ((k-1)K_p, \infty)$ and $l_2 \in ((k-1)K_q, \infty)$, the respective summands in (3.22) vanish. It follows that for $l \in ((k-1)K, \infty)$ the summands in (3.26) all vanish. The finiteness of \tilde{C}_K in expression (1.21) is immediate, without requiring decay rates for ϕ or growth bounds on C_k . Thus (1.21) holds when (ξ, \mathcal{P}) is of class (A1). \square

4. Proof of main results. We provide the proofs of Theorems 1.12, 1.15, and 1.13 in this order.

4.1. Proof of Theorem 1.12.

4.1.1. *Proof of expectation asymptotics (1.23).* The definition of the Palm probabilities gives

$$\mathbb{E}\mu_n^\xi(f) = \int_{W_n} f(n^{-1/d}u) \mathbb{E}_u^\xi(u, \mathcal{P}_n) \rho^{(1)}(u) \, du.$$

$$\begin{aligned} \text{Var}\mu_n^\xi(f) &= \mathbb{E} \sum_{x \in \mathcal{P}_n} f(n^{-1/d}x)^2 \xi^2(x, \mathcal{P}_n) \\ &\quad + \mathbb{E} \sum_{x, y \in \mathcal{P}_n, x \neq y} f(n^{-1/d}x) f(n^{-1/d}y) \xi(x, \mathcal{P}_n) \xi(y, \mathcal{P}_n) - \left(\mathbb{E} \sum_{x \in \mathcal{P}_n} f(n^{-1/d}x) \xi(x, \mathcal{P}_n) \right)^2 \end{aligned} \quad (4.1)$$

$$= \int_{W_n} f(n^{-1/d}u)^2 \mathbb{E}_u(\xi^2(u, \mathcal{P}_n)) \rho^{(1)}(u) du \quad (4.2)$$

$$+ \int_{W_n \times W_n} f(n^{-1/d}u) f(n^{-1/d}v) (m_{(2)}(u, v; n) - m_{(1)}(u; n) m_{(1)}(v; n)) dudv.$$

Since ξ satisfies the p -moment condition (1.19) for $p > 2$, we have that ξ^2 satisfies the p -moment condition for $p > 1$. Also, ξ and ξ^2 have the same radius of stabilization. Thus, the proof of expectation asymptotics, with ξ replaced by ξ^2 , shows that the first term in (4.1), multiplied by n^{-1} , converges to

$$\mathbb{E}_0 \xi^2(\mathbf{0}, \mathcal{P}) \rho^{(1)}(\mathbf{0}) \int_{W_1} f(x)^2 dx;$$

cf. expectation asymptotics (1.23). Setting $x = n^{-1/d}u$ and $z = v - u = v - n^{1/d}x$, the second term in (4.2), multiplied by n^{-1} , may be rewritten as

$$\begin{aligned} &\int_{W_1} \int_{W_n - n^{1/d}x} f(x + n^{-1/d}z) f(x) \\ &\quad \times [m_{(2)}(n^{1/d}x, n^{1/d}x + z; n) - m_{(1)}(n^{1/d}x; n) m_{(1)}(n^{1/d}x + z; n)] dz dx. \end{aligned} \quad (4.3)$$

Setting $\mathcal{P}_n^x := \mathcal{P} \cap (W_n - n^{1/d}x)$, the translation invariance of ξ and stationarity of \mathcal{P} yields

$$\begin{aligned} m_{(2)}(n^{1/d}x, n^{1/d}x + z; n) &= m_{(2)}(\mathbf{0}, z; \mathcal{P}_n^x) \\ m_{(1)}(n^{1/d}x; n) &= m_{(1)}(\mathbf{0}; \mathcal{P}_n^x) \\ m_{(1)}(n^{1/d}x + z; n) &= m_{(1)}(z; \mathcal{P}_n^x). \end{aligned}$$

Putting aside for the moment technical details, one expects that the above moments converge to $m_{(2)}(\mathbf{0}, z)$, $m_{(1)}(\mathbf{0})$ and $m_{(1)}(z) = m_{(1)}(\mathbf{0})$, respectively, when $n \rightarrow \infty$. Moreover, splitting the inner integral in (4.3) into two terms

$$\int_{W_n - n^{1/d}x} (\dots) dz = \int_{W_n - n^{1/d}x} \mathbf{1}[|z| \leq M] (\dots) dz + \int_{W_n - n^{1/d}x} \mathbf{1}[|z| > M] (\dots) dz \quad (4.4)$$

$|h_n^\xi(x, z)|$. To prove the convergence notice that

(4.6)

$$\begin{aligned} |m_{(2)}(\mathbf{0}, z; \mathcal{P}_n^x) - m_{(2)}(\mathbf{0}, z)| &= |\mathbb{E}_{\mathbf{0}, z}(X_n Y_n) - \mathbb{E}_{\mathbf{0}, z}(XY)| \rho^{(2)}(\mathbf{0}, z) \\ &\leq \kappa_2 (\mathbb{E}_{\mathbf{0}, z}|X_n Y_n - X_n Y| + \mathbb{E}_{\mathbf{0}, z}|X_n Y - XY|) \\ (4.7) \quad &\leq \kappa_2 (\mathbb{E}_{\mathbf{0}, z}(X_n^2) \mathbb{E}_{\mathbf{0}, z}(Y_n - Y)^2)^{1/2} + \kappa_2 (\mathbb{E}_{\mathbf{0}, z}(Y^2) \mathbb{E}_{\mathbf{0}, z}(X_n - X)^2)^{1/2}, \end{aligned}$$

where κ_2 bounds the second-order correlation function as at (1.11). We have already proved that $\mathbb{E}_{\mathbf{0}, z}(X_n^2)$, $\mathbb{E}_{\mathbf{0}, z}(Y^2)$ are bounded. Moreover

$$\begin{aligned} \mathbb{E}_{\mathbf{0}, z}(X_n - X)^2 &= \mathbb{E}_{\mathbf{0}, z}((X_n - X)^2 \mathbf{1}[X_n \neq X]) \\ &\leq \mathbb{E}_{\mathbf{0}, z}(X_n^2 \mathbf{1}[X_n \neq X]) + 2\mathbb{E}_{\mathbf{0}, z}(|X_n X| \mathbf{1}[X_n \neq X]) + \mathbb{E}_{\mathbf{0}, z}(X^2 \mathbf{1}[X_n \neq X]). \end{aligned}$$

The Hölder inequality gives for $p > 2$ and $2/p + 1/q = 1$,

$$\begin{aligned} \mathbb{E}_{\mathbf{0}, z}(X_n^2 \mathbf{1}[X_n \neq X]) &\leq (\mathbb{E}_{\mathbf{0}, z}(X_n^p))^{2/p} (\mathbb{P}_{\mathbf{0}, z}(X_n \neq X))^{1/q} \\ \mathbb{E}_{\mathbf{0}, z}(|X_n X| \mathbf{1}[X_n \neq X]) &\leq (\mathbb{E}_{\mathbf{0}, z}(X_n^p) \mathbb{E}_{\mathbf{0}, z}(X^p))^{1/p} (\mathbb{P}_{\mathbf{0}, z}(X_n \neq X))^{1/q} \\ \mathbb{E}_{\mathbf{0}, z}(X^2 \mathbf{1}[X_n \neq X]) &\leq (\mathbb{E}_{\mathbf{0}, z}(X^p))^{2/p} (\mathbb{P}_{\mathbf{0}, z}(X_n \neq X))^{1/q}. \end{aligned}$$

The p th moment of X_n and X under $\mathbb{E}_{\mathbf{0}, z}$ can be bounded by \tilde{M}_p using the p -moment condition (1.19) with $p > 2$ as in (4.5). Stabilization (1.14) with $l = 2$ gives

$$(4.8) \quad \mathbb{P}_{\mathbf{0}, z}(X_n \neq X) \leq \mathbb{P}_{\mathbf{0}, z}((\max(R^\xi(u, \mathcal{P}), R^\xi(u, \mathcal{P}_n)) > n^{1/d} d(x, \partial W_1))$$

$$(4.9) \quad \leq 2\varphi(a_2 n^{1/d} d(x, \partial W_1))$$

with the right-hand side converging to 0 for all $x \notin \partial W_1$. This proves that $\mathbb{E}_{\mathbf{0}, z}(X_n - X)^2$ and (by the very same arguments) $\mathbb{E}_{\mathbf{0}, z}(Y_n - Y)^2$ converge to 0 as $n \rightarrow \infty$ for all $x \notin \partial W_1$. Concluding this part of the proof, we have shown that the expression in (4.7) converges to 0 and thus $m_{(2)}(\mathbf{0}, z; \mathcal{P}_n^x)$ converges to $m_{(2)}(\mathbf{0}, z)$. Using similar arguments, we derive

$$\begin{aligned} |m_{(1)}(\mathbf{0}, \mathcal{P}_n^x) - m_{(1)}(\mathbf{0})| &= |\mathbb{E}_{\mathbf{0}}(X_n) - \mathbb{E}_{\mathbf{0}}(X)| \rho^{(1)}(\mathbf{0}) \\ &\leq \kappa_1 ((\mathbb{E}_{\mathbf{0}}(X_n)^2)^{1/2} + (\mathbb{E}_{\mathbf{0}}(X^2))^{1/2}) (\mathbb{P}_{\mathbf{0}}(X_n \neq X))^{1/2}, \end{aligned}$$

by the p -moment condition (1.19) and the stabilization property (1.14) for $p = 1$ one can show that $m_{(1)}(\mathbf{0}, \mathcal{P}_n^x)$ converges to $m_{(1)}(\mathbf{0})$ uniformly in x for all $x \in W_1 \setminus \partial W_1$. Exactly the same arguments assure convergence of $m_{(1)}(z, \mathcal{P}_n^x)$ to $m_{(1)}(z) = m_{(1)}(\mathbf{0})$. This concludes the proof of Lemma 4.1. \square

In order to complete the proof of variance asymptotics for general $f \in \mathcal{B}(W_1)$ (not necessarily continuous) we use arguments borrowed from the proof of [55, Theorem 2.1]. Recall that $x \in W_1$ is a Lebesgue point for f if $(\text{Vol}_d B_\varepsilon(x))^{-1} \int_{B_\varepsilon(x)} |f(z) -$

4.2. **Proof of Theorem 1.15.** The proof is inspired by the proofs of [44, Propositions 1 and 2]. By the refined Campbell theorem and stationarity of \mathcal{P} , we have

$$\begin{aligned} n^{-1}\text{Var}\widehat{H}_n^\xi(\mathcal{P}) &= \int_{W_n} \mathbb{E}_x \xi^2(x; \mathcal{P}) \rho^{(1)}(x) dx + \int_{W_n} \int_{W_n} [m_{(2)}(x, y) - m_{(1)}(x)m_{(1)}(y)] dy dx \\ (4.10) \quad &= \mathbb{E}_0 \xi^2(\mathbf{0}, \mathcal{P}) \rho^{(1)}(\mathbf{0}) + n^{-1} \int_{W_n} \int_{W_n} (m_{(2)}(x, y) - m_{(1)}(x)m_{(1)}(y)) dy dx. \end{aligned}$$

Writing $c(x, y) := m_{(2)}(x, y) - m_{(1)}(x)m_{(1)}(y)$, the double integral in (4.10) becomes ($z = y - x$)

$$\begin{aligned} n^{-1} \int_{W_n} \int_{W_n} (m_{(2)}(x, y) - m_{(1)}(x)m_{(1)}(y)) dy dx &= n^{-1} \int_{W_n} \int_{\mathbb{R}^d} c(\mathbf{0}, z) \mathbf{1}[x + z \in W_n] dz dx \\ &= n^{-1} \int_{W_n} \int_{\mathbb{R}^d} c(\mathbf{0}, z) \mathbf{1}[x \in W_n - z] dz dx. \end{aligned}$$

Write $\mathbf{1}[x \in W_n - z]$ as $1 - \mathbf{1}[x \in (W_n - z)^c]$ to obtain

$$\begin{aligned} n^{-1} \int_{W_n} \int_{W_n} (m_{(2)}(x, y) - m_{(1)}(x)m_{(1)}(y)) dy dx \\ = \int_{\mathbb{R}^d} c(\mathbf{0}, z) dz - n^{-1} \int_{\mathbb{R}^d} \int_{W_n} c(\mathbf{0}, z) \mathbf{1}[x \in \mathbb{R}^d \setminus (W_n - z)] dx dz. \end{aligned}$$

From (1.29), we have that $\gamma_{W_n}(z) := \text{Vol}_d(W_n \cap (\mathbb{R}^d \setminus (W_n - z)))$ and thus rewrite (4.10) as

$$(4.11) \quad n^{-1}\text{Var}\widehat{H}_n^\xi(\mathcal{P}) = \mathbb{E}_0 \xi^2(\mathbf{0}, \mathcal{P}) \rho^{(1)}(\mathbf{0}) + \int_{\mathbb{R}^d} c(\mathbf{0}, z) dz - n^{-1} \int_{\mathbb{R}^d} c(\mathbf{0}, z) \gamma_{W_n}(z) dz.$$

Now we claim that

$$\lim_{n \rightarrow \infty} n^{-1} \int_{\mathbb{R}^d} c(\mathbf{0}, z) \gamma_{W_n}(z) dz = 0.$$

Indeed, as noted in Lemma 1 of [44], for all $z \in \mathbb{R}^d$ we have $\lim_{n \rightarrow \infty} n^{-1} \gamma_{W_n}(z) = 0$. Since $n^{-1} c(\mathbf{0}, z) \gamma_{W_n}(z)$ is dominated by the fast-decreasing function $c(\mathbf{0}, z)$, the dominated convergence theorem gives the claimed limit. Letting $n \rightarrow \infty$ in (4.11) gives

$$(4.12) \quad \lim_{n \rightarrow \infty} n^{-1} \text{Var}\widehat{H}_n^\xi(\mathcal{P}) = \mathbb{E}_0 \xi^2(\mathbf{0}, \mathcal{P}) \rho^{(1)}(\mathbf{0}) + \int_{\mathbb{R}^d} c(\mathbf{0}, z) dz = \sigma^2(\xi),$$

where the last equality follows by the definition of $\sigma^2(\xi)$ in (1.22) and the finiteness follows by the fast-decreasing property of $c(\mathbf{0}, z, \mathcal{P})$ (which follows from the assumption of fast decay of the second mixed moment density).

nor on the localization properties of ξ . Semi-cluster measures for μ_n^ξ have the appealing property that they involve differences of measures on product spaces with product measures, and thus their Radon-Nikodym derivatives involve differences of correlation functions of the ξ -weighted measures.

In general, bounds on cumulant measures in terms of semi-cluster measures are not terribly informative. However, when ξ , together with \mathcal{P} , satisfy moment bounds and fast decay of correlations (1.21), then the situation changes. First, integrals of (S, T) semi-cluster measures on properly chosen subsets $W(S, T)$ of W_n^k , with (S, T) ranging over partitions of $\{1, \dots, k\}$, exhibit $O(n)$ growth. This is because the subsets $W(S, T)$ are chosen so that the Radon-Nikodym derivative of the (S, T) semi-cluster measure, being a difference of the correlation functions of the ξ -weighted measures, may be controlled by (1.21) for points $(v_1, \dots, v_k) \in W(S, T)$. Second, it conveniently happens that W_n^k is precisely the union of $W(S, T)$, as (S, T) ranges over partitions of $\{1, \dots, k\}$. Therefore, combining these observations, we see that every cumulant measure on W_n^k is a sum ranging over partitions (S, T) of $\{1, \dots, k\}$ of linear combinations of (S, T) semi-cluster measures on $W(S, T)$, each of which exhibits $O(n)$ growth.

Thus cumulant measures c_n^k exhibit growth *proportional to* $\text{Vol}_d(W_n)$ carrying \mathcal{P}_n , namely

$$(4.15) \quad \langle f^k, c_n^k \rangle = O(n), \quad f \in \mathcal{B}(W_1), \quad k = 2, 3, \dots$$

The remainder of Section 4.3 provides the details justifying (4.15).

Remarks on related work. (a) The estimate (4.15) first appeared in [6, Lemma 5.3], but the work of [21] (and to some extent [72]) was the first to rigorously control the growth of c_n^k on the diagonal subspaces, where two or more coordinates coincide. In fact Section 3 of [21] shows the estimate $\langle f^k, c_n^k \rangle \leq L^k (k!)^\beta n$, where L and β are constants independent of n and k . We assert that the arguments behind (4.15) are not restricted to Poisson input, but depend only on the fast decay of correlations (1.21) of the ξ -weighted measures and moment bounds (1.19). Since these arguments are not well known we present them in a way which is hopefully accessible and reasonably self-contained. Since we do not care about the constants in (4.15), we shall suitably adopt the arguments of [6, Lemma 5.3] and [72], taking the opportunity to make those arguments more rigorous. Indeed those arguments did not adequately explain the fast decay of the correlations of the ξ -weighted measures of the ξ -weighted measures on diagonal subspaces.

(b) The breakthrough paper [50] shows that the k th order cumulant for the *linear statistic* $\langle f, \sum_x \delta_{n^{-1/d_x}} \rangle / \sqrt{\text{Var} \langle f, \sum_x \delta_{n^{-1/d_x}} \rangle}$ vanishes as $n \rightarrow \infty$ and k large. This approach is extended to $\langle f, \mu_n^\xi \rangle$ in Section 4.4 thereby giving a second proof of the central limit theorem.

which equals zero unless equality holds among the coordinates in each non-empty subset of \mathcal{V} .

When μ is the atomic measure μ_n^ξ , we write M_n^k for $M^k(\mu_n^\xi)$. By the Campbell formula, considering repetitions in the k -fold product of \mathbb{R}^d , and putting $\tilde{y}_i := y_i(\mathcal{V})$ and $\mathcal{V} := (\mathcal{V}_1, \dots, \mathcal{V}_j)$ we have that

$$\begin{aligned} \langle f \otimes \dots \otimes f, M_n^k \rangle &= \mathbb{E}[\langle f, \mu_n^\xi \rangle \dots \langle f, \mu_n^\xi \rangle] \\ &= \sum_{j=1}^k \sum_{\mathcal{V}} \int_{(W_n)^j} \Pi_{i=1}^k f\left(\frac{y_i}{n^{1/d}}\right) \mathbb{E}_{\tilde{y}_1, \dots, \tilde{y}_j} [\Pi_{i=1}^j \xi^{|\mathcal{V}_i|}(\tilde{y}_i, \mathcal{P}_n)] \rho^{(j)}(\tilde{y}_1, \dots, \tilde{y}_j) \Pi_{i=1}^j dy_i(\mathcal{V}) \delta(\mathcal{V}). \end{aligned}$$

In other words, recalling Lemma 9.5.IV of [18] we get

$$(4.18) \quad dM_n^k(y_1, \dots, y_k) = \sum_{j=1}^k \sum_{\mathcal{V}} m^{(|\mathcal{V}_1|, \dots, |\mathcal{V}_j|)}(\tilde{y}_1, \dots, \tilde{y}_j; n) \Pi_{i=1}^j dy_i(\mathcal{V}) \delta(\mathcal{V}).$$

Cumulant measures. The k th cumulant measure $c_n^k := c^k(\mu_n)$ is defined analogously to the k th moment measure via

$$\langle f_1 \otimes \dots \otimes f_k, c^k(\mu_n) \rangle = c(\langle f_1, \mu_n \rangle \dots \langle f_k, \mu_n \rangle)$$

where $c(X_1, \dots, X_k)$ denotes the joint cumulant of the random variables X_1, \dots, X_k .

The existence of the cumulant measures c_n^l , $l = 1, 2, \dots$ follows from the existence of moment measures in view of the representation (4.17). Thus, we have the following representation for cumulant measures :

$$c_n^l = \sum_{T_1, \dots, T_p} (-1)^{p-1} (p-1)! M_n^{T_1} \dots M_n^{T_p},$$

where T_1, \dots, T_p ranges over all unordered partitions of the set $1, \dots, l$ (see p. 30 of [43]). Henceforth for $T_i \subset \{1, \dots, l\}$, let $M_n^{T_i}$ denote a copy of the moment measure M^{T_i} on the product space W^{T_i} . Multiplication denotes the usual product of measures: For T_1, T_2 disjoint sets of integers and for measurable $B_1 \subset (\mathbb{R}^d)^{T_1}$, $B_2 \subset (\mathbb{R}^d)^{T_2}$ we have $M_n^{T_1} M_n^{T_2}(B_1 \times B_2) = M_n^{T_1}(B_1) M_n^{T_2}(B_2)$. The first cumulant measure coincides with the expectation measure and the second cumulant measure coincides with the covariance measure.

Cluster and semi-cluster measures. We show that every cumulant measure c_n^k is a linear combination of products of moment and cluster measures. We first recall the definition of cluster and semi-cluster measures. A cluster measure $U_n^{S,T}$ on $W_n^S \times W_n^T$ for non-empty $S, T \subset \{1, 2, \dots\}$ is defined by

$$U_n^{S,T}(B \times D) = M_n^{S \cup T}(B \times D) - M_n^S(B) M_n^T(D)$$

v^S and v^T .

It is easy to see that for every $v := (v_1, \dots, v_k) \in W_n^k$, there is a partition (S, T) of $\{1, \dots, k\}$ such that $d(v^S, v^T) \geq D_k(v)/k^2$. If this were not the case then given $v := (v_1, \dots, v_k)$, the distance between any two components of v must be strictly less than $D_k(v)/k^2$ and we would get $\max_{i \leq k} \sum_{j=1}^k |v_i - v_j| \leq (k-1)kD_k/k^2 < D_k$, a contradiction. Thus W_n^k is the union of sets $W(S, T)$, $(S, T) \in \Xi(k)$, as asserted. We next describe the behavior of the differential $d(U_n^{S_1, T_1} M_n^{|S_2|} M_n^{|T_2|})$ on $W(S, T)$.

Semi-cluster measures on $W(S, T)$. Next, given $S_1 \subset S$ and $T_1 \subset T$, notice that $d(v^{S_1}, v^{T_1}) \geq d(v^S, v^T)$ where v^{S_1} denotes the projection of v^S onto $W_n^{S_1}$ and v^{T_1} denotes the projection of v^T onto $W_n^{T_1}$. Let $\Pi(S_1, T_1)$ be the partitions of S_1 into j_1 sets $\mathcal{V}_1, \dots, \mathcal{V}_{j_1}$, with $1 \leq j_1 \leq |S_1|$, and the partitions of T_1 into j_2 sets $\mathcal{V}_{j_1+1}, \dots, \mathcal{V}_{j_1+j_2}$, with $1 \leq j_2 \leq |T_1|$. Thus an element of $\Pi(S_1, T_1)$ is a partition of $S_1 \cup T_1$.

If a partition \mathcal{V} of $S_1 \cup T_1$ does not belong to $\Pi(S_1, T_1)$, then there is a partition element of \mathcal{V} containing points in S_1 and T_1 and thus, recalling (4.18), we have $\delta(\mathcal{V}) = 0$ on the set $W(S, T)$. Thus we make the crucial observation that, *on the set $W(S, T)$ the differential $d(M_n^{S_1 \cup T_1})$ collapses into a sum over partitions in $\Pi(S_1, T_1)$.* Thus $d(M_n^{S_1 \cup T_1})$ and $d(M_n^{S_1} M_n^{T_1})$ both involve sums of measures on common diagonal subspaces, as does their difference, made precise as follows.

LEMMA 4.2. *On the set $W(S, T)$ we have*

$$(4.21) \quad d(U_n^{S_1, T_1}) = \sum_{j_1=1}^{|S_1|} \sum_{j_2=1}^{|T_1|} \sum_{\mathcal{V} \in \Pi(S_1, T_1)} [\dots] \Pi_{i=1}^{j_1+j_2} dy_i(\mathcal{V}) \delta(\mathcal{V})$$

where

$$[\dots] := m^{(|\mathcal{V}_1|, \dots, |\mathcal{V}_{j_1}|, |\mathcal{V}_{j_1+1}|, \dots, |\mathcal{V}_{j_1+j_2}|)}(\tilde{y}_1, \dots, \tilde{y}_{j_1}, \tilde{y}_{j_1+1}, \dots, \tilde{y}_{j_1+j_2}; n) \\ - m^{(|\mathcal{V}_1|, \dots, |\mathcal{V}_{j_1}|)}(\tilde{y}_1, \dots, \tilde{y}_{j_1}; n) m^{(|\mathcal{V}_{j_1+1}|, \dots, |\mathcal{V}_{j_1+j_2}|)}(\tilde{y}_{j_1+1}, \dots, \tilde{y}_{j_1+j_2}; n).$$

The representations of $dM_n^{|S_2|}$ and $dM_n^{|T_2|}$ follow from (4.18), that is to say

$$(4.22) \quad dM_n^{|S_2|} = \sum_{j_3=1}^{|S_2|} \sum_{\mathcal{V} \in \Pi(S_2)} m^{(|\mathcal{V}_1|, \dots, |\mathcal{V}_{j_3}|)}(\tilde{y}_1, \dots, \tilde{y}_{j_3}; n) \Pi_{i=1}^{j_3} dy_i(\mathcal{V}) \delta(\mathcal{V}),$$

where $\Pi(S_2)$ runs over partitions of S_2 into j_3 sets, $1 \leq j_3 \leq |S_2|$. Similarly

$$(4.23) \quad dM_n^{|T_2|} = \sum_{j_4=1}^{|T_2|} \sum_{\mathcal{V} \in \Pi(T_2)} m^{(|\mathcal{V}_1|, \dots, |\mathcal{V}_{j_4}|)}(\tilde{y}_1, \dots, \tilde{y}_{j_4}; n) \Pi_{i=1}^{j_4} dy_i(\mathcal{V}) \delta(\mathcal{V}),$$

for the remaining indices $j \in \{1, \dots, k-1\}$ is similar. Write

$$\begin{aligned} & \int_{y_1 \in W_n} \dots \int_{y_k \in W_n} \tilde{\phi}\left(\frac{\tilde{c}_k D_k(y)}{k^2}\right) dy_1 \dots dy_k \\ &= \int_{y_1 \in W_n} \int_{w_2 \in W_{n-y_1}} \dots \int_{w_k \in W_{n-y_1}} \tilde{\phi}\left(\frac{\tilde{c}_k D_k(\mathbf{0}, w_2, \dots, w_k)}{k^2}\right) dy_1 dw_2 \dots dw_k. \end{aligned}$$

Now $D_k(\mathbf{0}, w_2, \dots, w_k) \geq \sum_{i=2}^k |w_i|$. Letting $e_k := \tilde{c}_k(k-1)/k^2$ gives

$$\begin{aligned} & \int_{y_1 \in W_n} \dots \int_{y_k \in W_n} \tilde{\phi}\left(\frac{\tilde{c}_k D_k(y)}{k^2}\right) dy_1 \dots dy_k \leq n \int_{w_2 \in \mathbb{R}^d} \dots \int_{w_k \in \mathbb{R}^d} \tilde{\phi}\left(\frac{e_k}{k-1} \sum_{i=2}^k |w_i|\right) dw_2 \dots dw_k \\ & \leq n \int_{w_2 \in \mathbb{R}^d} \dots \int_{w_k \in \mathbb{R}^d} \tilde{\phi}\left(\Pi_{i=2}^k |w_i|^{1/(k-1)}\right) dw_2 \dots dw_k = O(n), \end{aligned}$$

where the first inequality follows from the decreasing behavior of $\tilde{\phi}$, the second inequality follows from the arithmetic geometric mean inequality, and the last equality follows since $\tilde{\phi}$ is decreasing faster than any polynomial. We similarly bound the other summands for $j \in \{1, \dots, k-1\}$, completing the proof of Lemma 4.3. \square

4.3.4. Proof of Theorem 1.13. By the bound (4.20) and Lemma 4.3 we obtain (4.15). Letting C_n^k be the k th cumulant for $\langle f, \mu_n^\xi \rangle / \sqrt{\text{Var}\langle f, \mu_n^\xi \rangle}$, we obtain $C_n^1 = 0$, $C_n^2 = 1$, and for all $k = 3, 4, \dots$

$$C_n^k = O(n(\text{Var}\langle f, \mu_n^\xi \rangle)^{-k/2}).$$

Since $\text{Var}\langle f, \mu_n^\xi \rangle = \Omega(n^\nu)$ by assumption, it follows that if $k \in (2/\nu, \infty)$, then the k th cumulant tends C_n^k to zero as $n \rightarrow \infty$. By a classical result of Marcinkiewicz (see e.g. [68, Lemma 3]), we get that all cumulants C_n^k , $k \geq 3$, converge to zero as $n \rightarrow \infty$. This gives (4.14) as desired and completes the proof of Theorem 1.13. \square

4.4. Second proof of the central limit theorem. We now give a second proof of the central limit theorem which we believe is of independent interest. Even though this proof is also based on the cumulant method used in Section 4.3.1, we shall bound the cumulants using a different approach, using Ursell functions of the ξ -weighted measure and establishing a property equivalent to Brillinger mixing; see Remarks at the end of Section 4.4.2. Though much of this proof can be read independently of the proof in Section 4.3, we repeatedly use the definition of moments and cumulants from Section 4.3.2.

Our approach. We shall adapt the approach in [50, Sec. 4] replacing \mathcal{P}_{GEF} by our ξ -weighted measures, which are purely atomic measures. As noted in Sec-

tion $\gamma = \{\gamma(1), \dots, \gamma(l)\} \in \Pi(p)$ *refines* partition $\sigma = \{\sigma(1), \dots, \sigma(l_1)\} \in \Pi(p)$ if for all $i \in \{1, \dots, l\}$, $\gamma(i) \subset \sigma(j)$ for some $j \in \{1, \dots, l_1\}$. Otherwise, the partition γ is said to *mix* partition σ . Now using (4.25), we get for any $I \subsetneq \{1, \dots, p\}$

$$m^{(k_j:j \in I)}(x_j : j \in I) m^{(k_j:j \in I^c)}(x_j : j \in I^c) = \sum_{\substack{\gamma \in \Pi[p] \\ \gamma \text{ refines } \{I, I^c\}}} \prod_{i=1}^{|\gamma|} m_{\top}^{(k_j:j \in \gamma(i))}(x_j : j \in \gamma(i)),$$

and therefore, again in view of (4.25)

$$m_{\top}^{(k_1, \dots, k_p)}(x_1, \dots, x_p) = \sum_{\substack{\gamma \in \Pi[p], |\gamma| > 1 \\ \gamma \text{ mixes } \{I, I^c\}}} \prod_{i=1}^{|\gamma|} m_{\top}^{(k_j:j \in \gamma(i))}(x_j : j \in \gamma(i))$$

(4.30)

$$+ m^{(k_1, \dots, k_p)}(x_1, \dots, x_p) - m^{(k_j:j \in I)}(x_j : j \in I) m^{(k_j:j \in I^c)}(x_j : j \in I^c).$$

This extends the relation [50, last displayed formula in the proof of Claim 4.1] valid for point processes.

4.4.2. Fast decay of correlations and bounds for Ursell functions. We show now that fast decay of correlations (1.21) of the ξ -weighted measures implies some bounds on the Ursell functions of these measures. Since $m^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n)$ is invariant with respect to any joint permutation of its arguments (k_1, \dots, k_p) and (x_1, \dots, x_p) , fast decay of correlations (1.21) of the ξ -weighted measures may be rephrased as follows : There exists a fast-decreasing function $\tilde{\phi}$ and constants \tilde{C}_k , \tilde{c}_k , such that for any collection of positive integers k_1, \dots, k_p , $p \geq 2$, satisfying $k_1 + \dots + k_p = k$, for any nonempty, proper subset $I \subsetneq \{1, \dots, p\}$, for all $n \leq \infty$ and all configurations $x_1, \dots, x_p \in W_n$ of distinct points we have

$$(4.31) \quad \left| m^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n) - m^{(k_j:j \in I)}(x_j : j \in I; n) m^{(k_j:j \in I^c)}(x_j : j \in I^c; n) \right| \leq \tilde{C}_k \tilde{\phi}(\tilde{c}_k s),$$

where $s := d(\{x_j : j \in I\}, \{x_j : j \in I^c\})$.

Now we consider the bounds of Ursell functions of the ξ -weighted measures. Following the idea of [50, Claim 4.1] one proves that fast decay of correlations (1.21) of the ξ -weighted measures and the p -moment condition (1.19) imply that there exists a fast-decreasing function $\tilde{\phi}_{\top}$ and constants \tilde{C}_k^{\top} , \tilde{c}_k^{\top} , such that for any collection of positive integers k_1, \dots, k_p , $p \geq 2$, satisfying $k_1 + \dots + k_p = k$, for all $n \leq \infty$ and all configurations $x_1, \dots, x_p \in W_n$ of distinct points we have

$$(4.32) \quad |m_{\top}^{(k_1, \dots, k_p)}(x_1, \dots, x_p; n)| \leq \tilde{C}_k^{\top} \tilde{\phi}_{\top}(c_k^{\top} \text{diam}(x_1, \dots, x_p)),$$

where $\text{diam}(x_1, \dots, x_p) := \max_{i,j=1..p}(|x_i - x_j|)$. The proof uses the representa-

$\text{Var}\langle f, \mu_n^\xi \rangle = \Omega(n^\nu)$. We have

$$M_n^k := \mathbb{E}(\langle f, \mu_n^\xi \rangle)^k = \mathbb{E}\left(\sum_{x_i \in \mathcal{P}_n} f_n(x_i) \xi(x_i, \mathcal{P}_n)\right)^k,$$

where $f_n(\cdot) = f(\cdot/n^{1/d})$. Considering appropriately the repetitions of points x_i in the k th product of the sum and using the Campbell theorem at (1.9), one obtains

$$(4.35) \quad M_n^k = \sum_{\sigma \in \Pi[k]} \langle \bigotimes_{i=1}^{|\sigma|} f_n^{|\sigma(i)|} m^{(\sigma)}, \lambda_n^{|\sigma|} \rangle,$$

where λ_n^l denotes the Lebesgue measure on $(W_n)^l$ and \bigotimes denotes the tensor product of functions

$$\left(\bigotimes_{i=1}^p f_n^{k_j}\right)(x_1, \dots, x_p) = \prod_{i=1}^p (f_n)^{k_j}(x_j), \quad m^{(\sigma)}(x_1, \dots, x_{|\sigma|}; n) := m^{(|\sigma(1)|, \dots, |\sigma(|\sigma|)|)}(x_1, \dots, x_{|\sigma|}; n)$$

Using the above representation and (4.17) the k th cumulant $S_k(\mu_n^\xi(f))$ can be expressed as follows

$$(4.36) \quad S_k(\mu_n^\xi(f)) = \sum_{\gamma \in \Pi[k]} (-1)^{|\gamma|-1} (|\gamma|-1)! \sum_{\substack{\sigma \in \Pi[k] \\ \sigma \text{ refines } \gamma}} \prod_{i=1}^{|\gamma|} \langle \bigotimes_{j=1}^{|\gamma(i)/\sigma|} f_n^{(\gamma(i)/\sigma)(j)} m^{(\gamma(i)/\sigma)}, \lambda_n^{|\gamma(i)/\sigma|} \rangle$$

$$= \sum_{\sigma \in \Pi[k]} \sum_{\substack{\gamma \in \Pi[k] \\ \sigma \text{ refines } \gamma}} (-1)^{|\gamma|-1} (|\gamma|-1)! \prod_{i=1}^{|\gamma|} \langle \bigotimes_{j=1}^{|\gamma(i)/\sigma|} f_n^{(\gamma(i)/\sigma)(j)} m^{(\gamma(i)/\sigma)}, \lambda_n^{|\gamma(i)/\sigma|} \rangle,$$

where $\gamma(i)/\sigma$ is the partition of $\gamma(i)$ induced by σ . Note that for any partition $\sigma \in \Pi[k]$, with $|\sigma(j)| = k_j$, $j = 1, \dots, |\sigma| = p$, the inner sum in (4.36) can be rewritten as follows:

$$(4.37) \quad \sum_{\gamma \in \Pi[p]} (-1)^{|\gamma|-1} (|\gamma|-1)! \prod_{i=1}^{|\gamma|} \langle \bigotimes_{j \in \gamma(i)} f_n^{k_j} m^{(k_j: j \in \gamma(i))}, \lambda_n^{|\gamma(i)|} \rangle = \langle \bigotimes_{j=1}^p f_n^{k_j} m_{\top}^{(k_1, \dots, k_p)}, \lambda_n^p \rangle,$$

where the equality is due to (4.28). Consequently

$$(4.38) \quad S_k(\mu_n^\xi(f)) = \sum_{\sigma \in \Pi[k]} \langle \bigotimes_{j=1}^{|\sigma|} f_n^{|\sigma(j)|} m_{\top}^{(|\sigma(1)|, \dots, |\sigma(|\sigma|)|)}, \lambda_n^{|\sigma|} \rangle,$$

which extends the relation [50, Claim 4.3] valid for point processes. The formula (4.38), which expresses the k th cumulant in terms of the Ursell functions,

- [5] Yu. Baryshnikov and J. E. Yukich (2003), Gaussian fields and random packing, *J. of Stat. Phys.*, **111**, 443–463.
- [6] Yu. Baryshnikov and J. E. Yukich (2005), Gaussian limits for random measures in geometric probability. *Ann. Appl. Prob.*, **15**, no. 1A, 213–253.
- [7] K. Baumann and G. C. Hegerfeldt (1985), A noncommutative Marcinkiewicz theorem, *Publications of the Research Institute for Mathematical Sciences*, **21**, no. 1, 191–204.
- [8] J. Ben Hough, M. Krishnapur, Y. Peres, and B. Virág (2009), *Zeros of Gaussian Analytic Functions and Determinantal Point Processes*, Vol. 51, American Mathematical Society, Providence, RI.
- [9] C. Biscio and F. Lavancier (2016), Brillinger mixing of determinantal point processes and statistical applications, *Electronic J. of Statistics*, **10**, 582–607.
- [10] A. Björner (1995), Topological methods, In *Handbook of Combinatorics*, Vol. 2, 1819–1872.
- [11] B. Błaszczyszyn (1995), Factorial moment expansion for stochastic systems, *Stoch. Proc. Appl.*, **56**, 2, 321–335.
- [12] B. Błaszczyszyn, E. Merzbach and V. Schmidt (1997), A note on expansion for functionals of spatial marked point processes, *Stat. Probab. Letters*, **36**, 3, 299–306.
- [13] B. Błaszczyszyn and D. Yogeshwaran (2014), On comparison of clustering properties of point processes, *Adv. Appl. Probab.*, **46**, 1, 1–20.
- [14] B. Błaszczyszyn, D. Yogeshwaran and J. E. Yukich (2018), Supplement to “Limit theory for geometric statistics of point processes having fast decay of correlations”.
- [15] B. Błaszczyszyn, D. Yogeshwaran, and J. E. Yukich (2018), Limit theory for geometric statistics of point processes having fast decay of correlations, arXiv:1606.03988v3.
- [16] O. Bobrowski and R.J. Adler (2014), Distance functions, critical points, and the topology of random Čech complexes, *Homology Homotopy Appl.*, **16**, 2, 311–344.
- [17] A. Bulinski and E. Spodarev (2012), Central limit theorems for weakly dependent random fields. In *Stochastic Geometry, Spatial Statistics and Random Fields : Asymptotic methods*. Ed. Spodarev, E., Lect. Notes Math., vol. 2068.
- [18] D. J. Daley and D. Vere-Jones (2003), *An Introduction to the Theory of Point Processes*, Springer, NY.
- [19] L. Decreusefond, E. Ferraz, H. Randriam, and A. Vergne (2014), Simplicial Homology of Random Configurations, *Adv. in Appl. Probab.* **46**, 2, 1–20.
- [20] H. Edelsbrunner and J. L. Harer (2010), *Computational Topology : An Introduction*, American Mathematical Society, Providence, RI.
- [21] P. Eichelsbacher, M. Raic, and T. Schreiber (2015), Moderate deviations for stabilizing functionals in geometric probability, *Ann. Inst. Henri Poincaré Prob. Stat.*, **51**, No. 1, 89–128.
- [22] P. Eichelsbacher and C. Thäle (2014), New Berry-Esseen bounds for non-linear functionals of Poisson random measures, *Electron. J. Probab.* **102**(19).
- [23] R. Forman (2002), A user’s guide to discrete Morse theory, *Sém. Lothar. Combin.*, **48**.
- [24] P. J. Forrester and G. Honner (1999), Exact statistical properties of the zeros of complex random polynomials, *J. of Physics A: Math. and General*, **32**, 16, 2961.
- [25] V. Gershkovich and H. Rubinstein (1997), Morse theory for Min-type functions, *The Asian J. Math.* **1**, 696–715.
- [26] J. Grote and C. Thäle (2016), Gaussian polytopes: a cumulant-based approach, *preprint* arXiv:1602.06148.
- [27] A. Goldman (2010), The Palm measure and the Voronoi tessellation for the Ginibre point process, *Ann. Appl. Probab.*, **20**, No. 1, 90–128.
- [28] P. Hall (1988), *Introduction to the Theory of Coverage Processes*, John Wiley and Sons.
- [29] K-H. Hanisch (1982), On inversion formulae for n-fold Palm distributions of point processes in LCS-spaces, *Math. Nachr.*, **106**, 1, 171–179.
- [30] L. Heinrich (1994), Normal approximation for some mean-value estimates of absolutely regular

- tessellations, *Math. Meth. Stat.*, **3**, 1, 1–24.
- [31] L. Heinrich (2013), Asymptotic methods in statistics of random point processes, In *Stochastic Geometry, Spatial Statistics and Random Fields : Asymptotic methods*. Ed. Spodarev, E., Lect. Notes Math., Springer, vol. 2068.
- [32] L. Heinrich and I. Molchanov (1999), Central limit theorem for a class of random measures associated with germ-grain models, *Adv. Appl. Probab.*, **31**, 2, 283–314.
- [33] D. Hug, G. Last and M. Schulte (2016) Second order properties and central limit theorems for geometric functionals of Boolean models, *Ann. Appl. Probab.*, **26**, 1, 73–135.
- [34] G. Ivanoff (1982) Central limit theorems for point processes, *Stoch. Proc. Applications.*, **12**, 2, 171–186.
- [35] M. Kahle (2011), Random geometric complexes, *Discrete Comput. Geom.*, **45**, 3, 553–573.
- [36] M. Krishnapur (2006), Overcrowding estimates for zeroes of planar and hyperbolic Gaussian analytic functions, *J. Stat. Phys.*, **124**, No. 6, 1399–1423.
- [37] R. Lachièze-Rey and M. Schulte and J. E. Yukich (2017), Normal approximation for stabilizing functionals, *pre-print* arXiv:1702.00726.
- [38] R. Lachièze-Rey and G. Peccati (2015), New Kolmogorov bounds for functionals of binomial point processes, to appear in *Ann. Appl. Probab.*, arXiv:1505.04640.
- [39] G. Last and M. Penrose (2010), Poisson process Fock space representation, chaos expansion and covariance inequalities, *Probab. Th. Rel. Fields*, **150**, 663–690.
- [40] G. Last and M. Penrose (2012), Percolation and limit theory for the Poisson lilypond model, *Rand. Struct. Alg.*, **42**, 226–249.
- [41] G. Last, G. Peccati and M. Schulte (2016), Normal approximations on the Poisson space: Mehler’s formula, second order Poincaré inequalities and stabilization, *Probab. Theory Rel. Fields.*, **165**, 3, 667–723.
- [42] V. A. Malyshev (1975), The central limit theorem for Gibbsian random fields, *Soviet Math. Doklady* **16**, 1141–1145.
- [43] V. A. Malyshev and R. A. Minlos (1991), *Gibbs Random Fields*, Kluwer.
- [44] Ph. A. Martin and T. Yalcin (1980), The charge fluctuations in classical Coulomb systems, *J. Stat. Phys.*, **22**, No. 4, 435–463.
- [45] P. McCullagh and J. Møller (2006), The permanent process, *Adv. Appl. Probab.*, **38**, No. 4, 873–888.
- [46] J. Milnor (1963), *Morse Theory*, Princeton University Press.
- [47] J. Møller (2003), Shot noise Cox processes, *Adv. Appl. Probab.*, **35**, No. 3 614–640.
- [48] J. Møller and G.L. Torrisi (2005), Generalised shot noise Cox processes, *Adv. Appl. Probab.*, **37**, No. 1, 48–74.
- [49] F. Nazarov and M. Sodin (2011), Fluctuations in random complex zeros: asymptotic normality revisited, *Intern. Math. Research Notices*, **24**, 5720– 5759.
- [50] F. Nazarov and M. Sodin (2012), Correlation functions for random complex zeroes: Strong clustering and local universality, *Comm. Math. Phys.*, **310**, No. 1, 75–98.
- [51] I. Nourdin and G. Peccati (2012), *Normal Approximations with Malliavin Calculus : From Stein’s Method to Universality*, Cambridge Tracts in Mathematics, **192**, Cambridge University Press, Cambridge.
- [52] G. Peccati and M. Reitzner (Editors) (2016), *Stochastic Analysis for Poisson Point Processes: Malliavin Calculus, Wiener-It Chaos Expansions and Stochastic Geometry*, Springer-Verlag.
- [53] A. Poinas, B. Delyon, and F. Lavancier (2017), Mixing properties and central limit theorem for associated point processes, preprint, arXiv:1705.02276v2.
- [54] M. D. Penrose (2003), *Random Geometric Graphs*, Oxford Univ. Press, New York.
- [55] M. D. Penrose (2007), Gaussian limits for random geometric measures, *Elec. J. Probab.*, **12**, 989–1035.
- [56] M. D. Penrose (2007), Laws of large numbers in stochastic geometry with statistical applica-

- tions, *Bernoulli*, **13**, 1124-1150.
- [57] M. D. Penrose and J. E. Yukich (2001), Central limit theorems for some graphs in computational geometry, *Ann. Appl. Probab.*, **11**, 1005-1041.
- [58] M. D. Penrose and J. E. Yukich (2002), Limit theory for random sequential packing and deposition, *Ann. Appl. Probab.*, **12**, 272-301.
- [59] M. D. Penrose and J. E. Yukich (2003), Weak laws of large numbers in geometric probability, *Ann. Appl. Probab.*, **13**, 277-303.
- [60] M. D. Penrose and J. E. Yukich (2005), Normal approximation in geometric probability, *In Stein's Method and Applications*, Lect. Note Series, Inst. Math. Sci., Nat. Univ. of Singapore, **5**, A. Barbour and Louis Chen, Eds., 37-58.
- [61] T. Ram Reddy, S. Vadlamani and D. Yogeshwaran (2017), Central limit theorem for quasi-local statistics of spin models on Cayley graphs, *preprint* arXiv:1709.10424.
- [62] M. Reitzner and M. Schulte (2013), Central limit theorems for U -statistics of Poisson point processes, *Ann. Probab.*, **41**, No. 6, 3879–3909.
- [63] R. Schneider (1993), *Convex Bodies: the Brunn-Minkowski Theory*, Cambridge University Press, Cambridge.
- [64] R. Schneider and W. Weil (2008), *Stochastic and Integral geometry*, Probability and its Applications (New York), Springer-Verlag, Berlin.
- [65] T. Schreiber and J. E. Yukich (2013), Limit theorems for geometric functionals of Gibbs point processes, *Ann. Inst. Henri Poincaré Probab. Stat.*, **49**, 4, 1158 - 1182.
- [66] M. Schulte and C. Thäle (2016), Central limit theorems for the radial spanning tree. *Rand. Struct. Alg.*, online version.
- [67] T. Shirai and Y. Takahashi (2003), Random point fields associated with certain Fredholm determinants I: fermion, Poisson and boson point processes, *J. Funct. Anal.*, **205**, 2, 414–463.
- [68] A. Soshnikov (2002), Gaussian limit for determinantal random point fields, *Ann. Probab.*, **30**, 171-187.
- [69] A. Xia and J. E. Yukich (2015), Normal approximation for statistics of Gibbsian input in geometric probability, *Adv. Appl. Probab.*, **47**, 934-972.
- [70] D. Yogeshwaran and R. J. Adler (2015), On the topology of random complexes built over stationary point processes, *Ann. Appl. Probab.*, **25**, No. 6, 3338–3380.
- [71] J. E. Yukich (1998), *Probability Theory of Classical Euclidean Optimization Problems*, Lecture Notes in Mathematics, vol. 1675, Springer-Verlag.
- [72] J. E. Yukich (2013), Limit theorems in discrete stochastic geometry. In *Stochastic Geometry, Spatial Statistics and Random Fields : Asymptotic methods*. Ed. Spodarev, E., Lect. Notes Math., Springer, vol. 2068.
- [73] A. Zomorodian (2012), Topological data analysis. In *Advances in Applied and Computational Topology*, Proc. Sympos. Appl. Math., Amer. Math. Soc., **70**, 1–39.

INRIA/ENS, 2 RUE SIMONE IFF
CS 42112 75589 PARIS CEDEX 12.
FRANCE.

E-MAIL: Bartek.Blaszczyszyn@ens.fr

STATISTICS AND MATHEMATICS UNIT
INDIAN STATISTICAL INSTITUTE
BANGALORE - 560059

E-MAIL: d.yogesh@isibang.ac.in

DEPARTMENT OF MATHEMATICS
LEHIGH UNIVERSITY
BETHLEHEM PA 18015
E-MAIL: joseph.yukich@lehigh.edu