Chain Polynomials and Permutation Statistics

by

Mark Allan Skandera
M. S., Operations Research
University of California at Berkeley, 1992
Sc. B., Applied Mathematics
Brown University, 1991

Submitted to the Department of Mathematics
in partial fulfillment of the requirements for the degree of

Doctor of Philosophy

at the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

June 2000

© Mark Allan Skandera, MM. All rights reserved.
The author hereby grants to MIT permission to reproduce and distribute publicly paper and electronic copies of this thesis document in whole or in part, and to grant others the right to do so.

Author .....
Department of Mathematics
April 13, 2000

Certified by .....
Richard P. Stanley
Professor of Mathematics
Thesis Supervisor

Accepted by .....
Michael Sipser
Chairman, Applied Mathematics Committee

Accepted by .....
Thomasz S. Mrowka
Chairman, Department Committee on Graduate Students

Chain Polynomials and Permutation Statistics

by

Mark Allan Skandera

Submitted to the Department of Mathematics
on April 13, 2000, in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

Abstract

Motivated by results and conjectures involving f-vectors, h-vectors, and the sequences counting linear extensions of posets, we prove three new results which relate these sequences to permutation statistics and $(\mathbf{3}+\mathbf{1})$-free posets.

Our first result extends Dumont's permutation statistic to the rearrangement classes of arbitrary words, creating a generalized statistic which is Eulerian. As a consequence we show that for each distributive lattice $J(P)$ which is a product of chains, there is a poset Q such that the f-vector of Q is the h-vector of $J(P)$. This strengthens for products of chains a result of Stanley concerning the flag h-vectors of Cohen-Macaulay complexes [43, Cor. 4.5]. We conjecture that the strengthened result holds for all finite distributive lattices (Conjecture 2.7.1).

Our second result solves the open problem of defining a natural Eulerian permutation statistic whose joint distribution with INV on S_n is the same as that of des with MAJ. We define such a statistc stc, along with a set-valued function STC which is analogous to the descent set function D. We conjecture several properties of joint distributions involving the statistic stc, including the symmetry of the joint distribution of $(\mathrm{des}, \mathrm{stc})$ on S_n.

Lastly, we characterize $(\mathbf{3}+\mathbf{1})$-free posets in terms of their antiadjacency matrices. Generalizing a previous result for unit interval orders, this characterization relates $(\mathbf{3}+\mathbf{1})$-free posets to totally positive matrices and leads to a simple proof that the chain polynomial of a $(\mathbf{3}+\mathbf{1})$-free poset has only real zeros.

Thesis Supervisor: Richard P. Stanley
Title: Professor of Mathematics

Acknowledgments

I could not have written this thesis without the help of a number of people.

Richard Stanley undoubtedly helped the most by introducing me to algebraic combinatorics, suggesting research topics, lending me numerous books and articles, and offering endless patience as I struggled to ask the right questions about open problems. I am very fortunate to have been his advisee at MIT.

Jennifer Galovich, Ira Gessel, and Einar Steingrímsson taught me a great deal about permutation statistics, and inspired much of the work in Chapters 2 and 3.

Danny Kleitman was a great source of encouragement, and perhaps the first professor at MIT to believe in me and to help me believe in myself.

Other mathematicians offered suggestions involving specific proofs, helped me find relevant results in the literature, and brought related open questions to my attention. These are Sara Billey, Miklós Bóna, Tim Chow, Art Duval, Dominique Foata, Sergey Fomin, Patricia Hersh, Michael Kleber, Alex Postnikov, Don Rawlings, Vic Reiner, Gian-Carlo Rota, Rodica Simion, and John Stembridge. In addition, each of these people has been a great role model for me.

Finally, I must thank my parents for giving me since birth every advantage I needed to enter a Ph.D. program in mathematics.

Contents

Notation **9**

Introduction **11**

1 Three sequences associated to a poset **12**
1.1 Definitions of the three sequences 12
1.2 Distributive lattices and linear extensions of posets 15
1.3 The Neggers-Stanley conjectures 17
1.4 A theorem for balanced Cohen-Macaulay complexes 19
1.5 Three open problems 22

2 Dumont's statistic on words **24**
2.1 Eulerian permutation statistics 24
2.2 Word statistics 27
2.3 Generalizing Dumont's statistic 28
2.4 An application of Dumont's statistic 33
2.5 The chain map Ψ 35
2.6 Inverting the chain map 38
2.7 Open questions 41

3 Joint distributions of permutation statistics on S_n **43**
3.1 Introduction 43
3.2 Eulerian-Mahonian statistic pairs 44
3.3 Codes and major index tables 46
3.4 The statistic stc 52
3.5 The stc-set of a permutation 54
3.6 More equidistribution results 58
3.7 Open equidistribution questions 61

4 Posets whose chain polynomials have only real zeros **63**
4.1 Interval characterizations 63

4.2 Antiadjacency matrices of unit interval orders 64
4.3 The squared antiadjacency matrix 67
4.4 A characterization of $(\mathbf{3}+\mathbf{1})$-free posets 69
4.5 The altitude sequence 72
4.6 $(\mathbf{3}+\mathbf{1})$-free posets and distributive lattices 75

Bibliography **78**

Index **82**

Notation

$\mathbb{Z}$	the integers
$\mathbb{N}$	the nonnegative integers
$[n]$	the set of integers $\{1, 2, \ldots, n\}$
$\lvert T \rvert$	the cardinality of the set T
$\#T$	the cardinality of the set T
S_n	the symmetric group on n letters
$D(\pi)$	the descent set of π
$E(\pi)$	the excedance set of π
$LC(\pi)$	the (nonzero) letter set of code(π)
$\mathrm{des}(\pi)$	the number of descents in π
$\mathrm{exc}(\pi)$	the number of excedances in π
$\mathrm{dmc}(\pi)$	Dumont's statistic on π
$\textsc{maj}(\pi)$	the major index of π
$\textsc{inv}(\pi)$	the number of inversions in π
$\textsc{den}(\pi)$	Denert's statistic on π
$R(w)$	the set of rearrangements of the word w
$C(w)$	the set of codes of rearrangements of the word w
$w \cdot u$	the concatenation of words w, u
$w_1 \cdots w_m < u_1 \cdots u_m$	$w_i < u_i$ for $i = 1, \ldots, m$
$w_1 \cdots w_m \leq u_1 \cdots u_m$	$w_i \leq u_i$ for $i = 1, \ldots, m$
$w_1 \cdots w_m \lneqq u_1 \cdots u_m$	$w_1 \cdots w_m \leq u_1 \cdots u_m$ and $w_i < u_i$ for some $i \in [m]$
E_n	the set of words $\{w \in \mathbb{N}^n \mid w \leq (n-1) \cdots 1 \cdot 0\}$

$<_P$	is less than, in the poset P
$\mathbf{n} + \mathbf{m}$	the disjoint sum of an n-element chain and an m-element chain
$J(P)$	the distributive lattice corresponding to the poset P
$\mathcal{L}(P)$	the set of linear extensions of the poset P
$\mathcal{L}_k(P)$	the elements of $\mathcal{L}(P)$ which have exactly k descents
$\Delta(P)$	the order complex of the poset P
f_Σ	the f-vector of the simplicial complex Σ
f_P	the f-vector of the poset P
$f_P(x)$	the generating function of f_P
h_P	the h-vector of the poset P
$h_P(x)$	the generating function of h_P
Λ_x	the principal order ideal corresponding to element x in a poset
V_x	the principal dual order ideal corresponding to element x in a poset
$\langle A \rangle$	the order ideal corresponding to antichain A in a poset

Introduction

Many open problems in algebraic combinatorics concern the finite integer sequences $a = (a_0, \ldots, a_d)$ related to simplicial complexes, partially ordered sets (posets), and permutation statistics. Ideally, one would like to characterize the sequences which arise in various settings. For example, the Schützenberger-Kruskal-Katona theorem characterizes those sequences which are f-vectors of simplicial complexes. (See [48, p. 54], and also [4], [19].)

Precise characterization of sequences can be a very difficult problem, and authors often make progress toward characterization by proving and conjecturing related statements. (See [3], [4], [5], [12], [36], [37], [48, Ch. 2,3].) Typical results and conjectures relate the integers $a_0, \ldots, a_d$ by linear or quadratic inequalities. For instance, the sequence a is called *unimodal* if we have

$$a_0 \leq \cdots \leq a_j \geq \cdots \geq a_d, \text{ for some index } j,$$

and *log-concave* if we have

$$a_i^2 \geq a_{i-1}a_{i+1}, \text{ for } i = 1, \ldots, d-1.$$

In the case that a is a positive sequence, it is easy to see that log-concavity implies unimodality. A stronger property still is that the polynomial

$$a_0 + a_1 x + \cdots + a_d x^d$$

has only real zeros. (See [45].) Other results relate the sequences arising in one setting to those arising in another.

In Chapter 1, we define three sequences which have proven to be quite difficult to characterize. We state some known properties of these sequences, including a theorem of Stanley which relates the h-vectors of a particular class of simplicial complexes to the f-vectors of other complexes. The proof of a special case of this theorem and the famous Neggers-Stanley conjectures motivate three problems which we consider in greater detail in the remaining chapters.

Chapter 1

Three sequences associated to a poset

1.1 Definitions of the three sequences

Three integer sequences often associated to a finite poset are its f-vector, h-vector, and the sequence counting its linear extensions by descent. Let P be a finite poset on n elements. If x and y are two elements of P, we will write $x <_P y$ to indicate that x is less than y in P.

The *f-vector* of P counts *chains* by cardinality. A *chain* of P is an ordered k-tuple of elements $(x_1, \ldots, x_k)$ which satisfies

$$x_1 <_P \cdots <_P x_k.$$

We denote the f-vector of P by

$$f_P = (f_{-1}, f_0, \ldots, f_{d-1}),$$

where f_i is the number of $(i+1)$-element chains in P, and d is the maximum cardinality of a chain in P. By convention, we define $f_{-1} = 1$. There is considerable interest in characterizing the integer sequences which are f-vectors of posets, and in identifying the properties of P which guarantee f_P to be unimodal, log-concave, and which guarantee the generating function

$$f_P(x) = f_{-1} + f_0 x + \cdots + f_{d-1} x^d$$

to have only real zeros. We will call the polynomial $f_P(x)$ defined above the *chain polynomial* of P.

The *h-vector* of P,

$$h_P = (h_0, h_1, \ldots, h_d),$$



Figure 1-1: A poset whose h-vector isn't nonnegative.

is defined in terms of the f-vector by the equation

$$\sum_{i=0}^{d} f_{i-1}(x-1)^{d-i} = \sum_{i=0}^{d} h_i x^{d-i}.$$

From this definition, it is clear that knowing the h-vector of a poset is equivalent to knowing the f-vector. It is easy to verify that

$$h_k = \sum_{i=0}^{k} (-1)^{k-i} \binom{d-i}{k-i} f_{i-1}, \tag{1.1}$$

for $k = 0, \ldots, d$, and one may use this fact as an alternative definition of the h-vector. Another definition of the h-vector of a poset expresses its generating function in terms of a linear fractional transformation of the variable x in the chain polynomial. Defining the h-polynomial $h_P(x)$ by

$$h_P(x) = (1-x)^d f_P\left(\frac{x}{1-x}\right), \tag{1.2}$$

one easily verifies that the coefficients of $h_P(x)$ are the h-vector of P,

$$h_P(x) = h_0 + h_1 x + \cdots + h_d x^d.$$

Example 1.1.1. If P is the poset in Figure 1-1, then we have $f_P = (1, 4, 2)$ and $h_P = (1, 2, -1)$.

While the f-vector of a poset is positive by definition, this is not true of the h-vector. There is considerable interest in identifying the properties of P which guarantee the h-vector h_P to be nonnegative, unimodal, and log-concave, and which guarantee the h-polynomial $h_P(x)$ to have only real zeros.

Using (1.2) we observe two facts concerning f and h-vectors.

Observation 1.1.1. *Let P, P', and P'' be finite posets.*

1. *All zeros of $f_P(x)$ are real if and only if all zeros of $h_P(x)$ are real.*

2. *We have $f_P(x) = f_{P'}(x)f_{P''}(x)$ if and only if we have $h_P(x) = h_{P'}(x)h_{P''}(x)$.*

The polynomial $f_P(x)$ factors as $f_{P'}(x)f_{P''}(x)$ when P has the form $P = P' \oplus P''$. This poset is a disjoint union of the elements of P' and P'', in which we define $x <_P y$ whenever $x <_{P'} y$ or $x <_{P''} y$ or $x \in P'$, $y \in P''$. (See [44, Ch. 3].)

Now let us *label* P by some bijective function $\phi : P \to [n]$. If x and y are two elements of P which satisfy $x <_P y$ and are labelled as $\phi(x) = i$ and $\phi(y) = j$, we will abuse notation a bit and write $i <_P j$. We will say that P is *naturally labelled* if the map ϕ satisfies $\phi(x) < \phi(y)$ whenever $x <_P y$.

Our third important sequence counts linear extensions of P by descents, assuming a fixed labelling ϕ of P. A *linear extension* of P is a permutation $\pi = \pi_1 \cdots \pi_n$ satisfying $i < j$ whenever $\pi_i <_P \pi_j$. The set of such linear extensions is sometimes called the *Jordan-Hölder set* of P, and is denoted by $\mathcal{L}(P)$. A *descent* of a permutation π is a position i which satisfies $\pi_i > \pi_{i+1}$. We will denote the set of descents of π by $D(\pi)$ and the number of descents by $\mathrm{des}(\pi)$,

$$D(\pi) = \{i \,|\, \pi_i > \pi_{i+1}\},$$

$$\mathrm{des}(\pi) = \#\{i \,|\, \pi_i > \pi_{i+1}\}.$$

Counting descents in linear extensions of P, we partition the Jordan-Hölder set $\mathcal{L}(P)$ into disjoint subsets $\mathcal{L}_0(P), \ldots, \mathcal{L}_{n-1}(P)$, where $\mathcal{L}_k(P)$ is the subset of linear extensions of P having k descents,

$$\mathcal{L}_k(P) = \{\pi \in \mathcal{L}(P) \,|\, \mathrm{des}(\pi) = k\}.$$

Denoting the cardinality of $\mathcal{L}_k(P)$ by g_k, we define the integer sequence

$$g_P = (g_0, \ldots, g_{n-1}),$$

which counts linear extensions of P *by descent*. (See Example 1.2.1.) This sequence does not remain constant for all possible labellings of P. In particular, we have $g_0 = 1$ whenever P is naturally labelled and is $g_0 = 0$ otherwise.

It would be very interesting to characterize the integer sequences which arise in this manner, or to prove any properties of such sequences. In particular, it appears that for each finite labelled poset P, the sequence g_P is log-concave and unimodal, and the polynomial

$$g_P(x) = g_0 + g_1 x + \cdots + g_{n-1}x^{n-1}$$

has only real zeros. These properties have been proven only for special classes of posets. (See Section 1.3: the cases in which P is labelled naturally and arbitrarily correspond to Conjectures 1.3.2 and 1.3.3.)

Surprisingly, the sequences which count linear extensions of a poset by descent are closely related to f-vectors and h-vectors. We will describe an explicit relationship in the following section.

1.2 Distributive lattices and linear extensions of posets

A very important class of posets is the class of *distributive lattices*. (See [44, Ch. 3] for definitions.) Typically defined in terms of lattice join and meet operations, distributive lattices also may be defined in terms of antichains in posets. An *antichain* in a poset P is a subset of elements which are pairwise incomparable in P. For each antichain $X \subset P$ we define the corresponding *order ideal* $\langle X \rangle$ by

$$\langle X \rangle = \{x \in P \,|\, x \leq_P y \text{ for some } y \in X\},$$

and define a new poset $J(P)$ to be the set of order ideals of P, ordered by inclusion.

A distributive lattice is any poset $J(P)$ constructed in this way from an arbitrary poset P. We will denote by $\hat{0}$ and $\hat{1}$ the unique minimal and maximal elements of $J(P)$, which correspond to the order ideals containing no elements of P and all elements of P, respectively.

One important property of a distributive lattice $J(P)$ is that its h-vector is nonnegative. Furthermore, the h-vector of a distributive lattice has a simple combinatorial interpretation which links all three sequences in Section 1.1.

Theorem 1.2.1. *Let P be a naturally labelled finite poset on n elements, and let $h_{J(P)} = (h_0, \ldots, h_{n+1})$ be the h-vector of the distributive lattice $J(P)$. Then, $h_{J(P)}$ counts linear extensions of P by descents. That is, for $k = 0, \ldots, n+1$ we have*

$$h_k = |\mathcal{L}_k(P)|.$$

Proof. For each subset S of $[n-1]$, define $\alpha(S)$ to be the number of linear extensions of P whose descent set is contained in S, and define $\beta(S)$ to be the number of linear extensions of P whose descent set is equal to S.

$$\alpha(S) = \#\{\pi \in \mathcal{L}(P) \,|\, D(\pi) \subset S\},$$

$$\beta(S) = \#\{\pi \in \mathcal{L}(P) \,|\, D(\pi) = S\}.$$

The numbers $\alpha(S)$ and $\beta(S)$ are related by

$$\alpha(S) = \sum_{T \subset S} \beta(T),$$

or equivalently, using the principal of inclusion-exclusion, by

$$\beta(S) = \sum_{T \subset S} (-1)^{|S \smallsetminus T|} \alpha(T).$$

Thus we may express the cardinalities of the sets $\mathcal{L}_0(P), \ldots, \mathcal{L}_{n-1}(P)$ in terms of the numbers $\{\alpha(T) \,|\, T \subset [n-1]\}$,

$$\begin{aligned}
|\mathcal{L}_k(P)| &= \sum_{S \in \binom{[n-1]}{k}} \beta(S) \\
&= \sum_{S \in \binom{[n-1]}{k}} \sum_{T \subset S} (-1)^{|S \smallsetminus T|} \alpha(T) \\
&= \sum_{i=0}^{k} (-1)^{k-i} \binom{n-i-1}{k-i} \sum_{T \subset \binom{[n-1]}{i}} \alpha(T).
\end{aligned}$$

Trivially, $\mathcal{L}_k(P)$ is empty for k greater than $n-1$.

Furthermore, we may express the number $\alpha(T)$ in terms of the f-vector of the poset $J(P) \smallsetminus \{\hat{0}, \hat{1}\}$. Assume that the elements of T are $t_1 < \cdots < t_k$. It is not hard to see that $\alpha(T)$ counts chains in $J(P)$, the ranks of whose elements are are $t_1, \ldots, t_k$,

$$\alpha(T) = \#\{v_1 <_{J(P)} \cdots <_{J(P)} v_k \,|\, r(v_i) = t_i, \text{ for } i = 1, \ldots, k\}.$$

Denoting the f-vector of $J(P) \smallsetminus \{\hat{0}, \hat{1}\}$ by $f' = (f'_{-1}, f'_0, \ldots, f'_{n-2})$, we have

$$f'_{i-1} = \sum_{T \subset \binom{[n-1]}{i}} \alpha(T),$$

and therefore

$$|\mathcal{L}_k(P)| = \sum_{i=0}^{k} (-1)^{k-i} \binom{n-i-1}{k-i} f'_{i-1} = h'_k,$$

where $h' = (h'_0, \ldots, h'_{n-1})$ is the h-vector of $J(P) \smallsetminus \{\hat{0}, \hat{1}\}$. This is essentially the h-vector of $J(P)$, for by Observation 1.1.1 (2), we have

$$\begin{aligned}
f_{J(P)}(x) &= (1+x)^2 (f'_{-1} + f'_0 x + \cdots f'_{n-2} x^{n-1}), \\
h_{J(P)}(x) &= h'_0 + h'_1 x + \cdots h'_{n-1} x^{n-1} + 0x^n + 0x^{n+1}.
\end{aligned}$$

□

Example 1.2.1. Figure 1-2 shows a naturally labelled poset P and the corresponding distributive lattice $J(P)$. The chain polynomial and h-polynomial of $J(P)$ are given by

$$\begin{aligned}
f_{J(P)}(x) &= (1+x)^2 (1 + 6x + 10x^2 + 5x^3), \\
h_{J(P)}(x) &= 1 + 3x + x^2.
\end{aligned}$$



Figure 1-2: A poset P and the corresponding distributive lattice $J(P)$.

The linear extensions of P are $\mathcal{L}(P) = \{1234, 124/3, 13/24, 2/134, 2/14/3\}$. (We have marked descents by slashes.) Therefore, we see that $h_{J(P)}$ counts linear extensions of P by descent.

The astute reader will note from the above example that the h-polynomial $h_{J(P)}(x)$ is in fact the chain polynomial $f_Q(x)$ of another poset Q, whose elements and relations correspond to the "boxes" of the distributive lattice $J(P)$. This raises the question of identifying the distributive lattices whose h-vectors are the f-vectors of other posets. We will return to this question soon.

As we have implied in Section 1.1, the Jordan-Hölder set of a poset P depends upon a particular labelling of P. However, Theorem 1.2.1 implies the surprising fact that for any poset P, the cardinalities of the sets $\mathcal{L}_k(P)$ remain constant for all natural labellings of P. The reader may wish to switch the labels 1 and 2 on the poset P in Figure 1-2 to verify that the set $\mathcal{L}(P)$ changes, while the cardinalities $|\mathcal{L}_0(P)|, |\mathcal{L}_1(P)|, |\mathcal{L}_2(P)|$ do not.

1.3 The Neggers-Stanley conjectures

One interesting conjecture concerning f-vectors and h-vectors is due to Neggers [39, p. 114].

Conjecture 1.3.1. *(The Distributive Lattice Conjecture.) Let P be any poset on n elements, and let $J(P)$ be the corresponding finite distributive lattice. Then the polynomial*

$$f_{J(P)}(x) = f_{-1} + f_0 x + \cdots + f_n x^{n+1}$$

has only real zeros.

As we have have implied in the Introduction, the truth of this conjecture would imply the log-concavity and unimodality of the f-vector $f_{J(P)}$. Various proofs show that the conjecture holds for the special cases in which P is a disjoint sum of chains [42], a Ferrers poset [8], a Gaussian poset [8], and a poset of the form $P' + P''$, where P' and P'' are posets for which the conjecture holds. This last proof implies that the conjecture holds for the class of series-parallel posets. (See [44, Ch. 3] for definitions.) In addition, Stembridge has verified the conjecture for all posets P having eight or fewer elements [53]. A more general open problem is to determine whether the conjecture holds for polynomials of the form $f_L(x)$, where L is a *modular* lattice. No counterexamples are known.

By Observation 1.1.1 (1), Conjecture 1.3.1 is equivalent to the assertion that the polynomial

$$h_{J(P)}(x) = h_0 + h_1 x + \cdots + h_n x^n$$

has only real zeros. By Theorem 1.2.1, this polynomial counts linear extensions of P by descent. Therefore, the conjecture may be restated in terms of naturally labelled posets.

Conjecture 1.3.2. *Let P be any naturally labelled poset on n elements, and define the sequence $g_P = (g_0, \ldots, g_{n-1})$ by $g_i = |\mathcal{L}_i(P)|$. Then the polynomial*

$$g_P(x) = g_0 + g_1 x + \cdots + g_{n-1} x^{n-1}$$

has only real zeros.

Gasharov [32] has proven that the sequence g_P is unimodal in the case that P is a naturally labelled poset of rank less than three.

When the poset P in Conjecture 1.3.2 is *not* naturally labelled, we do not have the equality of sequences g_P and $h_{J(P)}$. Nevertheless, Stanley has conjectured that the same conclusion holds for arbitrary labellings of P. Thus we have the following stronger conjecture. (See [50, Problem 20].)

Conjecture 1.3.3. *(Poset Conjecture) Let P be an arbitrarily labelled poset on n elements, and define the sequence $g_P = (g_0, \ldots, g_{n-1})$ by $g_i = |\mathcal{L}_i(P)|$. Then the polynomial*

$$g_P(x) = g_0 + g_1 x + \cdots + g_{n-1} x^{n-1}$$

has only real zeros.

Brenti has proven that Conjecture 1.3.3 holds if P is a Ferrers poset with a column-strict labelling, or if P is a disjoint sum of chains [8]. Stembridge has verified the conjecture for all posets having seven or fewer elements [53].

Many equivalent forms of the Neggers-Stanley conjectures may be found in [8] and [56]. In these papers and elsewhere in the literature, the polynomials and sequences in question are named differently. The polynomial denoted by $W_P(x)$ corresponds to our polynomial $g_P(x)$. When P is naturally labelled, the polynomial denoted by $E_P(x)$ corresponds to our $f_{J(P)}(x)$, and otherwise, it corresponds to $(1+x)^d g_P(\frac{x}{1+x})$. In addition, the polynomials $W_P(x)$ and $E_P(x)$ sometimes differ from our polynomials by a multiplicative factor of x.

Recalling the coincidence from Example 1.2.1, in which the h-vector of one poset was the f-vector of another, we will consider a theorem in Section 1.4 which addresses a similar phenomenon.

1.4 A theorem for balanced Cohen-Macaulay complexes

The conditions for which the h-vector of one poset is the f-vector of another are unknown. It would be very interesting to obtain any information about this phenomenon. An existing result of this nature involves the f-vectors and h-vectors of simplicial complexes.

The f-vector of a simplicial complex generalizes that of a poset. If Σ is a $(d-1)$-dimensional simplicial complex, we define its f-vector to be the sequence

$$f_\Sigma = (f_{-1}, f_0, f_1, \ldots, f_{d-1}),$$

where f_i counts the number of i-dimensional faces of Σ. By convention, $f_{-1} = 1$. Since the chains of a poset P form a simplicial complex, known as the *order complex* of P and denoted by $\Delta(P)$, this new definition of the f-vector is compatible with the definition we gave in Section 1.1. That is, we have $f_P = f_{\Delta(P)}$. (See [44, p. 120].)

The h-vector of a simplicial complex generalizes that of a poset analogously. As in Section 1.1, we define the h-vector $h_\Sigma = (h_0, h_1, \ldots, h_d)$ in terms of f_Σ,

$$\sum_{i=0}^{d} f_{i-1}(x-1)^{d-i} = \sum_{i=0}^{d} h_i x^{d-i}.$$

For some conditions on a simplicial complex, one can show that its h-vector is the f-vector of another complex. Specifically, we have the following result due to Stanley [43, Cor. 4.5].

Theorem 1.4.1. *If Σ is a balanced Cohen-Macaulay complex, then its h-vector is the f-vector of some balanced simplicial complex Γ.*

(The original statement of this theorem asserts the stronger result that the *flag* h-vector of Σ is equal to the *flag* f-vector of Γ. We use the simpler version here to

emphasize its similarity to the theorems and conjectures in other sections of this paper.)

We define a simplicial complex to be *Cohen-Macaulay* if it satisfies a certain topological condition [48, p. 61], and *balanced* if we can color the vertices with d colors such that no face contains two vertices of the same color [48, p. 95]. The class of balanced Cohen-Macaulay complexes is quite important because it includes the order complexes of all distributive lattices. We have seen that the distributive lattices, in turn, contain information about all posets.

By placing an additional restriction on the complex Σ, one arrives at a special case of the theorem which has an elegant bijective proof. (See [6, Remark 6.6] and [20, Cor. 2.2].) Let us require that Σ be the order complex of a distributive lattice $J(P)$.

Theorem 1.4.2. *If $J(P)$ is any finite distributive lattice, then its h-vector is the f-vector of some balanced simplicial complex Γ.*

In this case, $h_\Sigma = h_{J(P)}$ counts the number of linear extensions of P by descent. Therefore, the theorem asserts that there is a bijective correspondence between linear extensions of P with k descents and $(k-1)$-faces of some simplicial complex $\Xi(P)$.

$$\mathcal{L}_k(P) \stackrel{1-1}{\longleftrightarrow} \{\sigma \,|\, \sigma \text{ a } (k-1)\text{-face of } \Xi(P)\}. \tag{1.3}$$

To prove Theorem 1.4.2, we will construct such a simplicial complex.

Definition 1.4.1. Let P be any poset, and define $\Xi(P)$ to be the collection of sets of the form $\{v_1, \ldots, v_k\}$ which satisfy

1. Each element $v_i = v_{i1} \cdots v_{in}$ belongs to $\mathcal{L}_1(P)$.
2. The k descent sets $D(v_i) = \{\ell_i\}$ for $i = 1, \ldots, k$ are distinct.
3. There is some linear extension π in $\mathcal{L}_k(P)$ with descent set $D(\pi) = \{\ell_1, \ldots, \ell_k\}$ such that we have

$$\{\pi_1, \ldots, \pi_{\ell_i}\} = \{v_{i1}, \ldots, v_{i\ell_i}\}, \text{ for } i = 1, \ldots, k. \tag{1.4}$$

It is not difficult to see that the collection $\Xi(P)$ of sets satisfies the conditions (1.3). Definition 1.4.1 (3) essentially defines a bijection $\Psi : \mathcal{L}(P) \to \Xi(P)$ which identifies a linear extension π with a set

$$\Psi(\pi) = \{v_1, \ldots, v_k\}$$

in $\Xi(P)$, where $k = \mathrm{des}(\pi)$. Let us call this bijection the *simplex map* and define it explicitly in terms of k *vertex maps*.

Definition 1.4.2. Let P be an n-element poset and let $\pi = \pi_1 \cdots \pi_n$ be a linear extension of P having descent set $D(\pi) = \{\ell_1, \ldots, \ell_k\}$. For each descent ℓ_i in $D(\pi)$, define the *vertex map* $\psi_{\ell_i} : \mathcal{L}(P) \to \mathcal{L}_1(P)$ by

$$\psi_{\ell_i}(\pi) = (\overline{\pi_1 \cdots \pi_{\ell_i}}) \cdot (\overline{\pi_{\ell_i+1} \cdots \pi_n}).$$

Define the *simplex map* $\Psi : \mathcal{L}(P) \to \Xi(P)$ by

$$\Psi(\pi) = \{\psi_{\ell_1}(\pi), \ldots, \psi_{\ell_k}(\pi)\}.$$

To demonstrate that $\Xi(P)$ is a well defined simplicial complex, and thus complete the proof of Theorem 1.4.2, we observe that for each k-set σ which belongs to the collection, each $(k-1)$-subset of σ does as well.

Proposition 1.4.3. *For any poset P, the collection $\Xi(P)$ of subsets of $\mathcal{L}_1(P)$ is a simplicial complex.*

Proof. Let P be a poset on n elements. For each pair of indices (i, j) satisfying $1 \leq i < j \leq n$, define the map $\phi_{ij} : S_n \to S_n$ by

$$\pi \mapsto (\pi_1 \cdots \pi_i) \cdot (\overline{\pi_{i+1} \cdots \pi_j}) \cdot (\pi_{j+1} \cdots \pi_n), \tag{1.5}$$

where $\overline{\pi_{i+1} \cdots \pi_j}$ is the strictly increasing rearrangement of the word $\pi_{i+1} \cdots \pi_j$.

It is easy to see that if π is a linear extension of P, then $\phi_{ij}(\pi)$ is as well. Moreover, if π belongs to $\mathcal{L}_k(P)$ and satisfies $\Psi(\pi) = \{v_1, \ldots, v_k\}$, then $\phi_{\ell_{j-1}\ell_{j+1}}(\pi)$ belongs to $\mathcal{L}_{k-1}(P)$ and satisfies $\Psi(\phi_{\ell_{j-1}\ell_{j+1}}(\pi)) = \Psi(\pi) \setminus \{v_j\}$. □

Let $n\mathbf{1}$ be the n-element antichain $\mathbf{1} + \cdots + \mathbf{1}$ and define $\Xi_n = \Xi(n\mathbf{1})$. By Proposition 1.4.3, we see that that for each n-element poset P, the complex $\Xi(P)$ is a subcomplex of Ξ_n.

Example 1.4.3. Figure 1-3 shows the complex Ξ_4.

For any poset P, the complex $\Xi(P)$ is balanced. Each vertex may be colored according to its single descent. However, the complex is not in general an *order complex*, so Theorem 1.4.2 does not imply that for each distributive lattice $J(P)$ there exists a poset $Q = Q(P)$ which satisfies $f_Q = f_{\Xi(P)} = h_{J(P)}$. The fact that the complex Ξ_4 is not an order complex is apparent from Figure 1-3. While Ξ_4 contains the faces $\{2413, 2341\}$, $\{2413, 4123\}$, and $\{2341, 4123\}$, it is *missing* the face $\{2413, 2341, 4123\}$. Clearly no three poset elements can be pairwise comparable without forming a three-element chain.



Figure 1-3: The complex Ξ_4.

In order for a simplicial complex to be an order complex it is necessary that each maximal non-face (or *missing face*) have at most two elements. This condition characterizes the class of *flag complexes*, which contains the class of order complexes. (See [48, p. 100].) Other properties which the complexes $\{\Xi_n\}_{n>0}$ lack are connectivity and acyclicity. The problem of calculating the homology groups of these complexes is open. (See [20, Question 4.15].)

Returning to Theorems 1.4.1 and 1.4.2, we note that the hypothesis of Theorem 1.4.2 is much stronger than that of Theorem 1.4.1, and yet the conclusions of the two theorems are the same. In principle, a stronger hypothesis should allow us to assert a stronger conclusion. This observation suggests the following questions, which we will discuss further in Section 2.7.

Question 1.4.4. For which posets P is there a flag complex Γ such that $f_\Gamma = h_{J(P)}$?

Question 1.4.5. For which posets P is there a poset Q such that $f_Q = h_{J(P)}$?

1.5 Three open problems

The results and conjectures in Sections 1.1-1.4 point to some intriguing open problems, and to several strategies for solving them.

Theorem 1.4.1, which refers to the broad class of balanced Cohen-Macaulay complexes, offers many opportunities for proving stronger results in special cases. In the interest of studying the h-vectors of distributive lattices, we will consider the following problem.

Problem 1.5.1. Find the most general poset class $\mathcal{P}$ such that for every poset P in $\mathcal{P}$ there exists a poset Q satisfying $f_Q = h_{J(P)}$. More generally, find the most general class $\mathcal{S}$ of simplicial complexes such that for every complex Σ in $\mathcal{S}$, there exists a poset Q satisfying $f_Q = h_\Sigma$.

The discussion following Theorem 1.4.2 suggests solving Problem 1.5.1 by using an Eulerian statistic to associate an order complex to each poset P in $\mathcal{P}$. We will use this strategy to solve a special case of Problem 1.5.1 in Chapter 2.

The proof of Theorem 1.2.1 suggests that a good permutation statistic to be used in solving Problem 1.5.1 might be one for which we have an analogy of the descent set. Thus, we could reinterpret the f-vector and h-vector in terms of new interpretations of the numbers $\alpha(S)$ and $\beta(S)$ in the proof of the theorem. A close look at the definition of the Jordan-Hölder set in Section 1.1 suggests that Mahonian statistics and their relations to our chosen Eulerian statistic may also be of some use.

Problem 1.5.2. Find a simple Eulerian permutation statistic *stat* to which we can associate a set-valued function $F : S_n \to 2^{[n-1]}$ satisfying the equations

$$\begin{aligned} |F(\pi)| &= \mathrm{stat}(\pi), && \text{for all } \pi \in S_n, \\ \#\{\pi \in S_n \,|\, F(\pi) = T\} &= \#\{\pi \in S_n \,|\, D(\pi) = T\}, && \text{for all } T \subset [n-1]. \end{aligned}$$

Find equidistribution results for pairs such as $(\mathrm{stat}, \textsc{maj})$ and $(\mathrm{stat}, \textsc{inv})$, and modify the proof of Theorem 1.2.1 to use stat and F in place of des and D.

We will consider this problem in Chapter 3, and will state some new results involving joint distributions of permutation statistics.

If there were *any* method of interpreting the h-vector of one poset as the f-vector of second, then one would be tempted to prove the Neggers-Stanley conjectures by demonstrating that the chain polynomial of this second poset has only real zeros. This possibility suggests obtaining as much information as possible about the posets whose chain polynomials have only real zeros.

Problem 1.5.3. Describe poset classes $\mathcal{P}$ which have the property that $f_P(x)$ has only real zeros whenever the poset P belongs to $\mathcal{P}$. Give various characterizations of the posets in any such class.

In Chapter 4 we will give a simple characterization of the posets in one such class $\mathcal{Q}$, and will note that there are many distributive lattices $J(P)$ for which there exists a poset Q in $\mathcal{Q}$ satisfying $f_Q = h_{J(P)}$.

All three of the problems above provide many opportunities for new results, and perhaps significant progress on any of them would help solve the Neggers-Stanley conjectures.

Chapter 2

Dumont's statistic on words

2.1 Eulerian permutation statistics

Let S_n be the symmetric group on n letters. A *permutation statistic* is a function stat : $S_n \to \mathbb{N}$ which maps permutations to nonnegative integers. The *distribution* of a permutation statistic on S_n is simply a count of the number of permutations π in S_n for which $\mathrm{stat}(\pi) = k$, for all possible values of k.

Recall from Section 1.1 that *des* is a statistic counting the number of descents in a permutation. Writing $\pi = \pi_1 \cdots \pi_n$, we have

$$\mathrm{des}(\pi) = \#\{i \,|\, \pi_i > \pi_{i+1}\}.$$

A second statistic *exc* counts the number of excedances in a permutation. We call position i an *excedance* in π if $\pi_i > i$. Therefore,

$$\mathrm{exc}(\pi) = \#\{i \,|\, \pi_i > i\}.$$

It is well known that the number of permutations in S_n with k descents equals the number of permutations in S_n with k excedances. This number is often denoted $A(n, k+1)$. The numbers $\{A(n, k+1) \,|\, n \geq 1; k+1 = 1, \ldots, n\}$ are called the *Eulerian numbers* and satisfy the recurrence

$$A(n, k+1) = (k+1)A(n-1, k+1) + (n-k-1)A(n-1, k), \tag{2.1}$$

subject to the initial conditions

$$A(1, k+1) = \begin{cases} 1, & \text{for } k+1 = 1, \\ 0, & \text{otherwise.} \end{cases}$$

Example 2.1.1. The following table shows the Eulerian numbers $A(n, k+1)$ for $n = 1, \ldots, 6$.

$n\backslash k+1$	1	2	3	4	5	6
1	1					
2	1	1				
3	1	4	1			
4	1	11	11	1		
5	1	26	66	26	1	
6	1	57	302	302	57	1

The generating function

$$A_n(x) = \sum_{k+1=1}^{n} A(n,k+1)x^{k+1} = \sum_{\pi \in S_n} x^{1+\mathrm{des}(\pi)} = \sum_{\pi \in S_n} x^{1+\mathrm{exc}(\pi)}$$

is called the nth *Eulerian polynomial*, and any permutation statistic *stat* satisfying

$$A_n(x) = \sum_{\pi \in S_n} x^{1+\mathrm{stat}(\pi)}$$

is called *Eulerian*.

A third Eulerian statistic, essentially defined by Dumont [18], counts the number of distinct nonzero letters in the code of a permutation. We define code(π) to be the word $c_1 \cdots c_n$, where

$$c_i = \#\{j > i \,|\, \pi_j < \pi_i\}.$$

Denoting Dumont's statistic by *dmc*, we have

$$\mathrm{dmc}(\pi) = \#\{\ell \neq 0 \,|\, \ell \text{ appears in code}(\pi)\}.$$

Example 2.1.2.

$$\pi = 2\ 8\ 4\ 3\ 6\ 7\ 9\ 5\ 1,$$
$$\mathrm{code}(\pi) = 1\ 6\ 2\ 1\ 2\ 2\ 2\ 1\ 0.$$

The distinct nonzero letters in code(π) are $\{1,2,6\}$. Thus, dmc(π) = 3.

Dumont showed bijectively that the statistic dmc is Eulerian,

$$\#\{\pi \in S_n \,|\, \mathrm{dmc}(\pi) = k\} = A(n,k+1).$$

While few researchers have found an application for dmc since [18], Foata [24] proved the following equidistribution result involving the statistics INV (inversions) and MAJ (major index). These two statistics belong to the class of *Mahonian* statistics. (See [24] for further information.)

Theorem 2.1.1. *The Eulerian-Mahonian statistic pairs* (des, INV) *and* (dmc, MAJ) *are equally distributed on* S_n*, i.e.*

$$\#\{\pi \in S_n \,|\, \mathrm{des}(\pi) = k; \mathrm{INV}(\pi) = p\} = \#\{\pi \in S_n \,|\, \mathrm{dmc}(\pi) = k; \mathrm{MAJ}(\pi) = p\}.$$

Note that the statistics des, exc, and dmc are all defined on a permutation π to be the cardinalities of certain sets associated to π. To name these sets, we define the *descent set* and *excedance set* of π to be the sets of descents and excedances in π. We denote these by $D(\pi)$ and $E(\pi)$, respectively. Similarly, we define the *letter set* of a word to be the set of its nonzero letters, and we denote the letter set of code(π) by $LC(\pi)$. Thus,

$$\begin{aligned} \mathrm{des}(\pi) &= |D(\pi)|, \\ \mathrm{exc}(\pi) &= |E(\pi)|, \\ \mathrm{dmc}(\pi) &= |LC(\pi)|. \end{aligned}$$

It is easy to see that for every subset T of $[n-1]$ there are permutations π, σ, and ρ in S_n satisfying

$$T = D(\pi) = E(\sigma) = LC(\rho).$$

In fact, Dumont's original bijection [18] shows that for each such subset T we have

$$\#\{\pi \in S_n \,|\, E(\pi) = T\} = \#\{\pi \in S_n \,|\, LC(\pi) = T\}.$$

However, the analogous statement involving $D(\pi)$ is not true. In Chapter 3 we will define a fourth Eulerian statistic stc and a related set-valued function STC which satisfy the equations

$$\begin{aligned} \mathrm{stc}(\pi) = |STC(\pi)|, &\qquad \text{for all } \pi \in S_n, \\ \#\{\pi \in S_n \,|\, STC(\pi) = T\} = \#\{\pi \in S_n \,|\, D(\pi) = T\}, &\qquad \text{for all } T \subset [n-1]. \end{aligned}$$

Eulerian permutation statistics are related to the h-vectors of distributive lattices by Theorem 1.2.1. In particular, since S_n is the Jordan-Hölder set of the n-element antichain $n\mathbf{1} = \mathbf{1} + \cdots + \mathbf{1}$, we see that the Eulerian numbers $A(n,1), \ldots, A(n,n)$ are the h-vector of the Boolean algebra $B_n = J(n\mathbf{1})$. However, while the h-vector of a distributive lattice $J(P)$ counts linear extensions of P by descent, it is not in general true that this h-vector counts linear extensions of P by other statistics. Several interesting results assert the equidistribution of two permutation statistics on the Jordan-Hölder set of a poset P, provided that P belongs to some specific poset class $\mathcal{P}$. (See [7], for example.) We will prove a result of this type in the following two sections, and will use the result in Sections 2.4-2.6 to strengthen a special case of Theorem 1.4.1.

2.2 Word statistics

Generalizing permutations on n letters are *words* $w = w_1 \cdots w_m$ on n letters, where $m \geq n$. We will assume that each letter in $[n]$ appears at least once in w. Generalizing the symmetric group S_n, we define the *rearrangement class* of w by

$$R(w) = \{w_{\sigma^{-1}(1)} \cdots w_{\sigma^{-1}(m)} \,|\, \sigma \in S_m\}.$$

Each element of $R(w)$ is called a *rearrangement* of w.

Many definitions pertaining to S_n generalize immediately to the rearrangement class of any word. In particular, the definitions of descent, descent set, code, letter set of a code, and Dumont's statistic remain the same for words as for permutations. Generalization of excedances requires only a bit of effort.

For any word w, denote by $\bar{w} = \bar{w}_1 \cdots \bar{w}_m$ the unique *nondecreasing rearrangement* of w. We define position i to be an *excedance* in w if $w_i > \bar{w}_i$. Thus,

$$\mathrm{exc}(w) = \#\{i \,|\, w_i > \bar{w}_i\}.$$

If position i is an excedance in word w, we will refer to the letter w_i as the *value* of excedance i. One can see word excedances most easily by associating to the word w the *biword*

$$\binom{\bar{w}}{w} = \binom{\bar{w}_1 \cdots \bar{w}_m}{w_1 \cdots w_m}.$$

Example 2.2.1. Let $w = 312312311$. Then,

$$\binom{\bar{w}}{w} = \binom{1\ 1\ 1\ 1\ 2\ 2\ 3\ 3\ 3}{3\ 1\ 2\ 3\ 1\ 2\ 3\ 1\ 1}.$$

Thus, $E(w) = \{1, 3, 4\}$ and $\mathrm{exc}(w) = 3$. The corresponding excedance values are 3, 2, and 3.

We will use biwords not only to expose excedances, but to define and justify maps in Sections 2.4 and 2.5. In particular, if $u = u_1 \cdots u_m$ and $v = v_1 \cdots v_m$ are words and y is the biword

$$y = \binom{u}{v},$$

then we will define *biletters* $y_1, \ldots, y_m$ by

$$y_i = \binom{u_i}{v_i},$$

and will define the rearrangement class of y by

$$R(y) = \{y_{\sigma^{-1}(1)} \cdots y_{\sigma^{-1}(m)} \,|\, \sigma \in S_m\}.$$

It is well known that the *word statistics* des and exc are equally distributed on the rearrangement class of any word w, just as the permutation statistics des and exc are equally distributed on S_n. That is,

$$\#\{v \in R(w) \,|\, \mathrm{exc}(v) = k\} = \#\{v \in R(w) \,|\, \mathrm{des}(v) = k\}.$$

Analogously to the case of permutation statistics, a word statistic *stat* is called *Eulerian* if it satisfies

$$\#\{v \in R(w) \,|\, \mathrm{stat}(v) = k\} = \#\{v \in R(w) \,|\, \mathrm{des}(v) = k\}$$

for any word w and any nonnegative integer k.

2.3 Generalizing Dumont's statistic

As implied in Section 1.1, we define Dumont's statistic on a word w to be the number of distinct nonzero letters in code(w),

$$\mathrm{dmc}(w) = |LC(w)|.$$

This generalized statistic is Eulerian.

Theorem 2.3.1. *If $R(w)$ is the rearrangement class of an arbitrary word w and k is any nonnegative integer, then*

$$\#\{v \in R(w) \,|\, \mathrm{dmc}(v) = k\} = \#\{v \in R(w) \,|\, \mathrm{exc}(v) = k\}.$$

Our bijective proof of the theorem depends upon an encoding of a word which we call the *excedance table*.

Definition 2.3.1. Let $v = v_1 \cdots v_m$ be an arbitrary word and let let $c = c_1 \cdots c_m$ be its code. Define the excedance table of v to be the unique word $\mathrm{etab}(v) = e_1 \cdots e_m$ satisfying

1. If i is an excedance in v, then $e_i = i$.
2. If $c_i = 0$, then $e_i = 0$.
3. Otherwise, e_i is the c_ith excedance of v having value at least v_i.

Note that etab(v) is well defined for any word v. In particular, if i is not an excedance in v and if $c_i > 0$, then there are at least c_i excedances in v having value at least v_i. To see this, define

$$k = \#\{j \in [m] \mid v_j < v_i\}.$$

Since c_i of the letters $\bar{v}_1, \ldots, \bar{v}_k$ appear to the right of position i in v, then at least c_i of the letters $\bar{v}_{k+1}, \ldots, \bar{v}_m$ must appear in the first k positions of v. The positions of these letters are necessarily excedances in v. An important property of the excedance table is that the letter set of etab(v) is precisely the excedance set of v.

Example 2.3.2. Let $v = 514514532$, and define $c = \mathrm{code}(v)$. Using v, $\bar{v}$, and c, we calculate $e = \mathrm{etab}(v)$,

$$\begin{aligned} \bar{v} &= 1\ 1\ 2\ 3\ 4\ 4\ 5\ 5\ 5, \\ v &= 5\ 1\ 4\ 5\ 1\ 4\ 5\ 3\ 2, \\ c &= 6\ 0\ 3\ 4\ 0\ 2\ 2\ 1\ 0, \\ e &= 1\ 0\ 3\ 4\ 0\ 3\ 4\ 1\ 0. \end{aligned}$$

Calculation of $e_1, \ldots, e_5$ and e_9 is straightforward since the positions $i = 1, \ldots, 5$ and 9 are excedances in v or satisfy $c_i = 0$. We calculate e_6, e_7, and e_8 as follows. Since $c_6 = 2$, and the second excedance in v with value at least $v_6 = 4$ is 3, we set $e_6 = 3$. Since $c_7 = 2$, and the second excedance in v with value at least $v_7 = 5$ is 4, we set $e_7 = 4$. Since $c_8 = 1$, and the first excedance in v with value at least $v_8 = 3$ is 1, we set $e_8 = 1$.

We prove Theorem 2.3.1 with a bijection $\theta : R(w) \to R(w)$ which satisfies

$$E(v) = LC(\theta(v)), \tag{2.2}$$

and therefore

$$\mathrm{exc}(v) = \mathrm{dmc}(\theta(v)). \tag{2.3}$$

Definition 2.3.3. Let $w = w_1 \cdots w_m$ be any word. Define the map $\theta : R(w) \to R(w)$ by applying the following procedure to an arbitrary element v of $R(w)$.

1. Define the biword $z = \binom{v}{\mathrm{etab}(v)}$.
2. Let y be the unique rearrangement of z satisfying $y = \binom{u}{\mathrm{code}(u)}$.
3. Set $\theta(v) = u$.

Construction of y is quite straightforward. Let $e = e_1 \cdots e_m = \mathrm{etab}(v)$, and linearly order the biletters $z_1, \ldots, z_m$ by setting $z_i < z_j$ if

$$v_i < v_j, \text{ or}$$
$$v_i = v_j \text{ and } e_i > e_j.$$

Break ties arbitrarily. Considering the biletters according to this order, insert each biletter z_i into y to the left of e_i previously inserted biletters.

Example 2.3.4. Let v and e be as in Example 2.3.2. To compute $\theta(v)$, we define

$$z = \binom{v}{e} = \begin{pmatrix} 5\ 1\ 4\ 5\ 1\ 4\ 5\ 3\ 2 \\ 1\ 0\ 3\ 4\ 0\ 3\ 4\ 1\ 0 \end{pmatrix}.$$

We consider the biletters of z in the order

$$\binom{1}{0}, \binom{1}{0}, \binom{2}{0}, \binom{3}{1}, \binom{4}{3}, \binom{4}{3}, \binom{5}{4}, \binom{5}{4}, \binom{5}{1},$$

and insert them individually into y:

$$\binom{1}{0}, \begin{pmatrix} 1\ 1 \\ 0\ 0 \end{pmatrix}, \begin{pmatrix} 1\ 1\ 2 \\ 0\ 0\ 0 \end{pmatrix}, \begin{pmatrix} 1\ 1\ 3\ 2 \\ 0\ 0\ 1\ 0 \end{pmatrix}, \begin{pmatrix} 1\ 4\ 1\ 3\ 2 \\ 0\ 3\ 0\ 1\ 0 \end{pmatrix}, \ldots$$

Finally we obtain

$$y = \begin{pmatrix} u \\ \mathrm{code}(u) \end{pmatrix} = \begin{pmatrix} 1\ 4\ 5\ 5\ 4\ 1\ 3\ 5\ 2 \\ 0\ 3\ 4\ 4\ 3\ 0\ 1\ 1\ 0 \end{pmatrix}$$

and set $\theta(v) = 145541352$.

It is easy to see that any biword z has at most one rearrangement y satisfying Definition 2.3.3 (2). Such a rearrangement exists if and only if we have

$$e_i \leq \#\{j \in [m] \mid v_j < v_i\}, \text{ for } i = 1, \ldots, m, \tag{2.4}$$

or equivalently, if and only if

$$\bar{v}_{e_i} < v_i, \text{ for } i = 1, \ldots, m, \tag{2.5}$$

where we define $\bar{v}_0 = 0$ for convenience.

Observation 2.3.2. *Let $v = v_1 \cdots v_m$ be any word and let $e = \mathrm{etab}(v)$. Then we have*

$$e_i \leq \#\{j \in [m] \mid v_j < v_i\}, \text{ for } i = 1, \ldots, m.$$

Proof. If i is an excedance in v, then $e_i = i$ and $\bar{v}_1 \leq \cdots \leq \bar{v}_i < v_i$. If $c_i = 0$, then $e_i = 0$. Otherwise, define

$$k = \#\{j \in [m] \mid v_j < v_i\}.$$

By the discussion following Definition 2.3.1, at least c_i of the positions $1, \ldots, k$ are excedances in v with values at least v_i. The letter e_i, being one of these excedances, is therefore at most k. □

Thus the map θ is well defined and satisfies (2.2) and (2.3). We invert θ by applying the procedure in the following proposition.

Proposition 2.3.3. *Let $y = \binom{u}{c}$ be a biword satisfying $c = \mathrm{code}(u)$. The following procedure produces a rearrangement $z = \binom{v}{e}$ of y satisfying $e = \mathrm{etab}(v)$.*

1. *For each letter ℓ in $L(c)$, find the greatest index i satisfying $c_i = \ell$, and define $z_\ell = y_i$. Let S be the set of such greatest indices, and define $T = [m] \smallsetminus S$.*

2. *For each index $i \in T$, define*

$$d_i = \begin{cases} \#\{j \in S \mid c_j \leq c_i; u_j \geq u_i\}, & \text{if } c_i > 0, \\ 0, & \text{otherwise.} \end{cases}$$

3. *Let $(y_{\sigma^{-1}(i)})_{i \in T}$ be the unique rearrangement of $(y_i)_{i \in T}$ satisfying*

$$(d_{\sigma^{-1}(i)})_{i \in T} = \mathrm{code}((u_{\sigma^{-1}(i)})_{i \in T}).$$

4. *Insert the biletters $(y_{\sigma^{-1}(i)})_{i \in T}$ in order into the remaining positions of z.*

Proof. The procedure above is well defined. In particular, we may perform step 3 because the biword

$$\binom{u_i}{d_i}_{i \in T}$$

satisfies

$$d_i \leq \#\{j \in T \mid u_j < u_i\}, \text{ for each } i \in T,$$

as required by (2.4). To see that this is the case, let i be an index in T with $c_i > 0$. In step 1 we have placed d_i biletters y_j with $u_j > u_i$ into positions $1, \ldots, c_i$ of z. Thus, at least d_i biletters y_j with $u_j \leq \bar{u}_{c_i}$ have *not* been placed into these positions. The index j of any such biletter belongs to S only if $c_j > c_i$. However, since $\bar{u}_{c_j} < u_j \leq \bar{u}_{c_i} < u_i$, we have $c_j < c_i$. Thus, j belongs to T.

We begin to verify that $e = \mathrm{etab}(v)$ by calculating the excedance set of v. We claim that $E(v) = L(c)$.

The positions $L(c) = \{c_j \,|\, j \in S\}$ are excedances in v, because for each index j in S, we have $v_{c_j} = u_j > \bar{u}_{c_j} = \bar{v}_{c_j}$. These positions are in fact the only excedances in v. For each index j in T, denote by $\phi(j)$ the position of z into which we have placed y_j. Supposing that some indices $\{\phi(j) \,|\, j \in T\}$ are excedances in v, choose $i \in T$ so that $\phi(i)$ is the leftmost of these excedances and define

$$k = \#\{j \in [m] \,|\, u_j < u_i\}.$$

Then, we have

$$k > \#\{j \in S \,|\, c_j \leq k\} + \#\{j \in T \,|\, \sigma(j) < \sigma(i)\}. \tag{2.6}$$

Since $c_i \leq k$ by (2.4), we have

$$\#\{j \in S \,|\, c_j \leq k\} = \#\{j \in S \,|\, c_j \leq c_i\} + \#\{j \in S \,|\, c_i < c_j \leq k\},$$

and by the definition of σ, we have

$$\begin{aligned}\#\{j \in T \,|\, \sigma(j) < \sigma(i)\} &= \#\{j \in T \,|\, u_j < u_i\} - d_i \\ &= \#\{j \in T \,|\, u_j < u_i\} - \#\{j \in S \,|\, c_j \leq c_i; u_j \geq u_i\}.\end{aligned}$$

Using these identities to simplify (2.6), we obtain

$$\#\{j \in S \,|\, u_j < u_i; c_j > c_i\} > \#\{j \in S \,|\, c_i < c_j \leq k\}. \tag{2.7}$$

If j belongs to the set on the left hand side of (2.7) and satisfies $c_j > k$, then we have

$$u_j > \bar{u}_{c_j} \geq \bar{u}_k = u_i - 1,$$

a contradiction. On the other hand, if each index j in this set satisfies $c_j \leq k$, then we have the inclusion

$$\{j \in S \,|\, u_j < u_i; c_j > c_i\} \subset \{j \in S \,|\, c_i < c_j \leq k\},$$

which contradicts (2.7). We conlcude that the set $\{\phi(j) \,|\, j \in T\}$ is precisely the set of non-excedances in v, and that we have

$$E(v) = L(c) = \{c_j \,|\, j \in S\}.$$

Finally, we show that e has the defining properites of etab(v). For each index j in S, we have defined $e_{c_j} = c_j$ so that e satisfies condition (1) of Definition 2.3.1. Let c' be the code of v. We claim that for each index $i \in T$, we have

$$e_{\phi(i)} = c_i = \begin{cases} \text{the } c'_{\phi(i)}\text{th excedance in } v \text{ having value at least } u_i, & \text{if } c'_{\phi(i)} > 0, \\ 0, & \text{otherwise.} \end{cases}$$

By our definition of the sequence $(d_i)_{i \in T}$, it suffices to show that $c'_{\phi(i)} = d_i$ for each index i. The subword $v_{\phi(i)+1} \cdots v_m$ of v includes d_i letters $v_{\phi(j)}$ with $j \in T$ and $v_{\phi(j)} < v_{\phi(i)}$. On the other hand, any excedance in v to the right of $\phi(i)$ has value *greater* than $v_{\phi(i)}$. We conclude that $c'_{\phi(i)} = d_i$. $\square$

The above procedure inverts θ because the biword z it produces is the *unique* rearrangement of y having the desired properties.

Proposition 2.3.4. *Let $v = v_1 \cdots v_m$ be an arbitrary word, and define*

$$z = \begin{pmatrix} v \\ e \end{pmatrix} = \begin{pmatrix} v \\ \mathrm{etab}(v) \end{pmatrix}.$$

If there is any rearrangement z' of z satisfying

$$z' = \begin{pmatrix} v' \\ e' \end{pmatrix} = \begin{pmatrix} v' \\ \mathrm{etab}(v') \end{pmatrix},$$

then $z' = z$.

Proof. Let L be the letter set of e. By Definition 2.3.1, we must have $E(v) = E(v') = L$. Let i be an excedance of v and v'. By condition (1) of Definition 2.3.1 we must have $e_i = e'_i = i$, and by condition (3) the upper letters v_i and v'_i must be as large as possible. Thus, $(z_i)_{i \in L} = (z'_i)_{i \in L}$.

Let $T = [m] \setminus L$ be the set of non-excedance positions of v and v', and consider the corresponding subsequences of biletters $(z_i)_{i \in T}$ and $(z'_i)_{i \in T}$. By condition (3) of Definition 2.3.1, the codes of $(v_i)_{i \in T}$ and $(v'_i)_{i \in T}$ are determined by the excedances and excedance values in v and v'. Thus, the two codes must be identical. Applying the argument following Example 2.3.4, we conclude that $(z_i)_{i \in T} = (z'_i)_{i \in T}$. □

2.4 An application of Dumont's statistic

Recall from Theorems 1.4.1 and 1.4.2 that the h-vector of any distributive lattice $J(P)$ is also the f-vector of some balanced simplicial complex Γ. It would be interesting to determine when Γ can be chosen to be an *order complex* $\Delta(Q)$ for some poset Q. Then, $h_{J(P)}$ would be the f-vector of Q and one could study the polynomial $h_{J(P)}(x)$ by studying chains in Q. As an application of Dumont's (word) statistic, we will show that this is the case when $J(P)$ is a product of chains (or equivalently, when P is a disjoint sum of chains.)

Theorem 2.4.1. *Let the distributive lattice $J(P)$ be a product of chains. Then there is a poset Q such that the h-vector of $J(P)$ is the f-vector of Q.*

If $J(P)$ is a product of chains having cardinalities $(p_1 + 1), \ldots, (p_n + 1)$, then P is the disjoint *sum* of chains $(\mathbf{p_1} + \cdots + \mathbf{p_n})$. (See Section 4.1 and [44, Ch. 3].) It is not difficult to see that linear extensions of P are in bijective correspondence with rearrangements of the word $w = 1^{p_1} \cdots n^{p_n}$,

$$\mathcal{L}(P) \overset{1-1}{\longleftrightarrow} R(w).$$

Combining this observation with Theorem 2.3.1, we restate Theorem 2.4.1 in terms of Dumont's statistic.



Figure 2-1: A poset with a k-element chain for each k-letter code in $C(11223)$.

Proposition 2.4.2. *Let w be any word and define the vector $h = (h_0, \ldots, h_d)$ by*

$$h_i = \#\{u \in R(w) \,|\, \mathrm{dmc}(u) = i\},$$

where d is the maximum cardinality of $LC(u)$ over all rearrangements u of w. Then, there is a poset Q whose f-vector is h.

Let $C(w)$ be the set of codes of all rearrangements of w,

$$C(w) = \{\mathrm{code}(u) \,|\, u \in R(w)\}.$$

Proposition 2.4.2 asserts that for any word w, there is a bijection between k-letter codes in $C(w)$ and k-element chains in some poset Q,

$$\{c \in C(w) \,|\, c \text{ a } k\text{-letter code}\} \stackrel{1-1}{\longleftrightarrow} \{(v_1 <_Q \cdots <_Q v_k) \in \Delta(Q)\}. \tag{2.8}$$

To prove the proposition, and therefore Theorem 2.4.1, we will construct such a poset $Q = Q(w)$ as follows.

Definition 2.4.1. For any word w, let Q be the subset of one-letter codes in $C(w)$, and let c and c' be codes in Q whose letters are ℓ and ℓ', respectively. Define $c <_Q c'$ if

1. $\ell < \ell'$.
2. The multiplicity of ℓ in c is strictly *greater* than that of ℓ' in c'.
3. For each position i such that $c'_i = \ell'$, we have $c_{i+\ell'-\ell} = \ell$.

Example 2.4.2. Figure 2-1 shows the poset Q corresponding to the word $w = 11223$. The f-vector f_Q counts words in $R(w)$ by Dumont's statistic. Equivalently, it counts linear extensions of the poset $P = \mathbf{2} + \mathbf{2} + \mathbf{1}$ by descent, and is equal to the h-vector of $J(P)$,

$$f_Q = h_{J(P)} = (1, 12, 15, 2).$$

In Sections 2.5 and 2.6 we will demonstrate that the poset Q satisfies the bijections of (2.8). Mimicking the proof of Theorem 1.4.2, we will give an explicit bijection $\Psi : C(w) \to \Delta(Q)$ which takes k-letter codes in $C(w)$ to k-element chains in Q.

2.5 The chain map Ψ

Fix a nondecreasing word $w = w_1 \cdots w_m$ on n letters, and define the poset Q as in Definition 2.4.1. We will define a *chain map* $\Psi : C(w) \to \Delta(Q)$ which will identify a code c with a chain

$$\Psi(c) = v_1 <_Q \cdots <_Q v_k,$$

of elements in Q. If c is a code on the k letters $\ell_1 < \cdots < \ell_k$, then each poset element v_i will be a code whose unique nonzero letter is ℓ_i. Specifically, we will determine v_i by applying a *vertex map* $\psi_{\ell_i} : C(w) \to Q$ to c,

$$v_i = \psi_{\ell_i}(c).$$

After proving that the vertex map satisfies

$$\psi_{\ell_i}(c) <_Q \psi_{\ell_j}(c)$$

whenever $\ell_i < \ell_j$, we will define the chain map to be a product of vertex maps,

$$\Psi(c) = \psi_{\ell_1}(c) <_Q \cdots <_Q \psi_{\ell_k}(c).$$

We begin by observing that several simple operations on codes in $C(w)$ yield other codes in $C(w)$.

Observation 2.5.1. *Let $u = u_1 \cdots u_m$ be a rearrangement of w and let $c = c_1 \cdots c_m$ be its code.*

1. *If $c_i > c_{i+1}$, then the word*

$$c' = c_1 \cdots c_{i-1} \cdot c_{i+1} \cdot (c_i - 1) \cdot c_{i+2} \cdots c_m$$

belongs to $C(w)$.

2. *If for some $r > i$, c_i is strictly less than $c_{i+1}, \ldots, c_r$, and $c_i > c_{r+1}$, then the word*

$$c'' = c_1 \cdots c_{i-1} \cdot c_{r+1} \cdot c_{i+1} \cdots c_r \cdot (c_i - 1) \cdot c_{r+2} \cdots c_m$$

belongs to $C(w)$.

3. *If $c_i < c_{i+1}$, or if $c_i = c_{i+1}$ and $u_i < u_{i+1}$, then the word*

$$c''' = c_i \cdots c_{i-1} \cdot (c_{i+1} + 1) \cdot c_i \cdot c_{i+2} \cdots c_m$$

belongs to $C(w)$.

Proof. Let u' be the word obtained from u by switching the letters in positions i and $i+1$, and let u'' be the word obtained by switching the letters in positions i and $r+1$,

$$u' = (u_1 \cdots u_{i-1} \cdot u_{i+1} \cdot u_i \cdot u_{i+2} \cdots u_m),$$
$$u'' = (u_1 \cdots u_{i-1} \cdot u_{r+1} \cdot u_{i+1} \cdots u_r \cdot u_i \cdot u_{r+2} \cdots u_m).$$

(1) We have $u_i > u_{i+1}$ and $c' = \mathrm{code}(u')$.
(2) We have $u_{r+1} < u_i < u_{i+1}, \cdots, u_r$ and $c'' = \mathrm{code}(u'')$.
(3) We have $u_i < u_{i+1}$ and $c''' = \mathrm{code}(u')$. □

Using this observation we will define two families of maps from $C(w)$ to itself, $\lambda_1, \ldots, \lambda_{m-1}$ and $\mu_1, \ldots, \mu_{m-1}$. Then, composing maps from these two families, we will define the family of vertex maps $\psi_1, \ldots, \psi_{m-1}$.

The map $\lambda_{\ell_i} : C(w) \to C(w)$ removes from a code c all letters ℓ_j which are greater than ℓ_i. It essentially changes each such letter ℓ_j to ℓ_i and moves it $\ell_j - \ell_i$ places to the right in c. If we identify c with the k-element chain $v_1 <_Q \cdots <_Q v_k$, then we will identify $\lambda_{\ell_i}(c)$ with the i-element subchain $v_1 <_Q \cdots <_Q v_i$.

Definition 2.5.1. Let ℓ be a nonzero letter. Define the map $\lambda_\ell : C(w) \to C(w)$ by performing the following procedure on a code c.

For $i = m, m-1, \ldots, 1$,

If $c_i > \ell$, then

1. Set $\delta = c_i - \ell$.
2. Redefine $c = c_1 \cdots c_{i-1} \cdot c_i \cdots c_{i+\delta} \cdot \ell \cdot c_{i+\delta+1} \cdots c_m$.

Analogous to λ_{ℓ_i}, the map $\mu_{\ell_i} : C(w) \to C(w)$ removes all letters which are *less* than ℓ_i. It simply changes each lower letter to 0. Thus, if we identify c with the k-element chain $v_1 <_Q \cdots <_Q v_k$, then we will identify $\mu_{\ell_i}(c)$ with the $(k-i+1)$-element subchain $v_i <_Q \cdots <_Q v_k$.

Definition 2.5.2. Let ℓ be a nonzero letter. Define the map $\mu_\ell : C(w) \to C(w)$ by $\mu_\ell(c) = a_1 \cdots a_m$, where

$$a_i = \begin{cases} 0, & \text{if } c_i \leq \ell, \\ c_i, & \text{otherwise.} \end{cases}$$

The maps $\lambda_1, \ldots, \lambda_{m-1}$, and $\mu_1, \ldots, \mu_{m-1}$ are well defined, for their definitions are merely repeated applications of Observation 2.5.1 (1) and (2). Note that the composition $\mu_\ell \lambda_\ell$ produces a code on the single letter ℓ. This code is an element of Q, and a *vertex* of $\Delta(Q)$.

Definition 2.5.3. Let ℓ be a nonzero letter. Define the *vertex map* $\psi_\ell : C(w) \to Q$ by

$$\psi_\ell = \mu_\ell \lambda_\ell.$$

It is easy to see that $\lambda_\ell^2 = \lambda_\ell$, and therefore that $\psi_\ell \lambda_\ell = \psi_\ell$. These and the following relations will be essential in establishing a bijection between $C(w)$ and $\Delta(Q)$.

Proposition 2.5.2. *Let ℓ and ℓ' be letters, $1 \leq \ell < \ell' \leq n$. The maps $\lambda_\ell, \lambda_{\ell'}, \psi_\ell$, and $\psi_{\ell'}$ satisfy the relations*

1. $\lambda_{\ell'} \lambda_\ell = \lambda_\ell \lambda_{\ell'} = \lambda_\ell$.
2. $\psi_\ell \lambda_{\ell'} = \psi_\ell$.
3. $\psi_\ell(c) <_Q \psi_{\ell'}(c)$, *if c contains both letters.*

Proof. (1) Let $c = \text{code}(u)$ be an element of $C(w)$. By the comments following Definition 2.5.2, we may interpret $\lambda_\ell(c)$ as follows. Define $b = b_1 \cdots b_m$ by

$$b_i = \begin{cases} \ell, & \text{if } c_i > \ell, \\ c_i, & \text{otherwise,} \end{cases}$$

and rearrange the biword $\binom{u}{b}$ as $\binom{u'}{b'}$ so that $b' = \text{code}(u')$. Then, $b' = \lambda_\ell(c)$.

It is not hard to see that there is a unique such rearrangement. Using this interpretation, it is easy to see that $\lambda_{\ell'} \lambda_\ell$, $\lambda_\ell \lambda_{\ell'}$, and λ_ℓ describe the same procedure.

(2) Using (1), we have $\psi_\ell \lambda_{\ell'} = \mu_\ell \lambda_\ell \lambda_{\ell'} = \mu_\ell \lambda_\ell = \psi_\ell$.

(3) We may assume that ℓ' is the greatest letter in c. (Otherwise, we define $d = \lambda_{\ell'}(c)$ and note that $\psi_\ell(c) = \psi_\ell(d)$ and $\psi_{\ell'}(c) = \psi_{\ell'}(d)$.) Let $e = \psi_\ell(c)$ and $e' = \psi_{\ell'}(c)$. Clearly, the multiplicity of ℓ in e is strictly greater than that of ℓ' in e', for

$$\#\{i \,|\, e_i = \ell\} = \#\{i \,|\, c_i \geq \ell\} > \#\{i \,|\, c_i \geq \ell'\} = \#\{i \,|\, e'_i = \ell'\}.$$

Next, we show that for any position i of e' satisfying $e'_i = \ell$, we must have $e_{i+\ell'-\ell} = \ell$. Since by assumption, ℓ' is the greatest letter in c, we have $e'_i = \ell'$ if and only if $c_i = \ell'$. To find e, we first calculate $\lambda_\ell(c)$ by the procedure of Definition 2.5.1. At each iteration i such that $c_i = \ell'$, we place the letter ℓ into position $i + \ell' - \ell$ of $\lambda_\ell(c)$. This position will not be altered by iterations $i - 1, \ldots, 1$, since all letters of c are no greater than ℓ'. Finally, since $e = \mu_\ell \lambda_\ell(c)$, and μ_ℓ changes only those letters less than ℓ, we see that $e_{i+\ell'-\ell} = \ell$ for every position i such that $e_i = \ell'$. □

Now we may define the map Ψ.

Definition 2.5.4. Define the *chain map* $\Psi : C(w) \to \Delta(Q)$ by

$$\Psi(c) = \psi_{\ell_1}(c) <_Q \cdots <_Q \psi_{\ell_k}(c),$$

where $\ell_1 < \cdots < \ell_k$ are the distinct nonzero letters in c.

2.6 Inverting the chain map

We will define a map $\Phi : \Delta(Q) \to C(w)$ which takes a k-element chain in Q to a k-letter code in $C(w)$. By demonstrating that Φ inverts Ψ, we will complete the proof of Proposition 2.4.2.

We begin by defining an operation $\vee : C(w) \times Q \to C(w)$ which joins a new letter to a code.

Definition 2.6.1. Let $d \in Q$ be a code whose unique nonzero letter is ℓ', and let $c \in C(w)$ be a code whose greatest letter is ℓ. Assume that $\psi_\ell(c) <_Q d$. Define the code $e = c \vee d$ by the following procedure.

1. For each i such that $d_i = \ell'$, set $e_i = \ell'$ and cross out the ℓ in position $i + \delta$ of c.
2. Fill the remaining positions of e with the remaining components of c, in order.

Note that $L(e) = L(c) \cup \{\ell\}$. Therefore, we may map a chain of k one-letter codes to a single k-letter code by iterating the join operation.

Definition 2.6.2. Let $v_1 <_Q \cdots <_Q v_k$ be a chain of one-letter codes on the letters $\ell_1 < \cdots < \ell_k$, respectively. Define the map $\Phi : \Delta(Q) \to C(w)$ by

$$\Phi(v_1 <_Q \cdots <_Q v_k) = (\cdots((v_1 \vee v_2) \vee v_3)\cdots) \vee v_k.$$

The following proposition shows that the join operation is well defined. It follows that Φ is well defined also.

Proposition 2.6.1. *If c and d are codes in $C(w)$ satisfying the hypotheses of Definition 2.6.1, then $c \vee d$ also belongs to $C(w)$.*

Proof. Let u and y be words in $R(w)$ whose codes $c = \mathrm{code}(u)$ and $d = \mathrm{code}(y)$ satisfy the conditions of Definition 2.6.1. Consider the leftmost position i in c such that $c_i = \ell$ and $d_{i-\delta} = \ell'$. By assumption, $c_{i-1} \leq c_i$. If $c_{i-1} < c_i$, or if $c_{i-1} = c_i$ and $u_{i-1} < u_i$, then we may apply Observation 2.5.1 (3) $\ell' - \ell$ times to obtain the word

$$c_1 \cdots c_{i-\delta-1} \cdot \ell' \cdot c_{i-\delta+1} \cdots \hat{c_i} \cdots c_m,$$

which belongs to $C(w)$. (Here, $\hat{c_i}$ means that the letter c_i is omitted.) Repeating this process for each such position i, we redefine the join operation. Therefore it suffices to show that for every position i satisfying $d_{i-\delta-1} = 0$, $d_{i-\delta} = \ell'$, and $c_{i-1} = c_i = \ell$, we have $u_{i-1} < u_i$.

Let i be such a position and suppose that $u_{i-1} = u_i$. Since $d_{i-\delta} = \ell'$ and $d_{i-\delta-1} = 0$, there are exactly $i - \delta - 1 + \ell' = i + \ell - 1$ letters in y which are strictly less than y_i. In particular, we have

$$w_{i+\ell-1} < w_{i+\ell}. \tag{2.9}$$

Let k be the number of positions preceeding i such that $u_{i-k} = u_{i-k+1} = \cdots = u_i$ and $c_{i-k} = c_{i-k+1} = \cdots = \ell$. Then there are exactly $i - k - 1 + \ell$ letters in u which are strictly less than u_i $(= u_{i-1} = \cdots = u_{i-k})$. In particular, we have

$$w_{i-k-1+\ell} < w_{i-k+\ell} = w_{i-k+1+\ell} = \cdots = w_{i+\ell},$$

which contradicts (2.9). We conclude that $u_{i-1} < u_i$, and therefore that $c \vee d$ belongs to $C(w)$. □

To begin demonstrating that Φ inverts Ψ we note the following relations satisfied by ψ, λ and $\vee$.

Proposition 2.6.2. *The pair of maps (ψ, λ) inverts the operation $\vee$ in the following sense.*

1. *Let $c \in C(w)$ be a code with greatest letter ℓ, and let $d \in Q$ be a code with letter $\ell' > \ell$ and satisfying $\psi_\ell(c) <_Q d$. Then we have*

$$\psi_{\ell'}(c \vee d) = d,$$
$$\lambda_\ell(c \vee d) = c.$$

2. *Let $c \in C(w)$ be a code whose greatest two letters are $\ell < \ell'$. Then we have*

$$\lambda_\ell(c) \vee \psi_{\ell'}(c) = c.$$

Proof. (1) Let S be the set of positions of d containing the letter ℓ', and let $\delta = \ell' - \ell$.

Define the words $e = c \vee d$, $d' = \psi_{\ell'}(c \vee d)$, and $c' = \lambda_\ell(c \vee d)$. Calculating e, we have

$$(e_i)_{i \in S} = \ell' \cdots \ell',$$
$$(e_i)_{i \notin S} = (c_i)_{i-\delta \notin S}.$$

Since e contains no letters greater than ℓ', we have $d' = \psi_{\ell'}(e) = \mu_{\ell'}(e)$. Thus, $d' = d$:

$$(d'_i)_{i \in S} = \ell' \cdots \ell',$$
$$(d'_i)_{i \notin S} = 0 \cdots 0.$$

Calculating $c' = \lambda_\ell(e)$, we change each occurrence of ℓ' in e to ℓ, and move it δ positions to the right. Since $\psi_\ell(c) <_Q d$, we see that $c' = c$:

$$(c'_i)_{i-\delta \in S} = \ell \cdots \ell = (c_i)_{i-\delta \in S},$$
$$(c'_i)_{i-\delta \notin S} = (c_i)_{i-\delta \notin S}.$$

(2) Similar. □

Completing the proof of Proposition 2.4.2, the following proposition shows that Ψ is bijective.

Proposition 2.6.3. *Let $c \in C(w)$ be a code on the letters $\ell_1 < \cdots < \ell_k$, and let $v_1 <_Q \cdots <_Q v_k$ be a k-element chain in Q, where the letter of v_i is ℓ_i for each i. The maps Ψ and Φ satisfy*

1. $\Psi\Phi(v_1 <_Q \cdots <_Q v_k) = v_1 <_Q \cdots <_Q v_k$.
2. $\Phi\Psi(c) = c$.

Proof. (1) By Definition 2.6.2, we have

$$\Phi(v_1 <_Q \cdots <_Q v_k) = (\cdots((v_1 \vee v_2) \vee v_3)\cdots) \vee v_k.$$

Applying $\Psi = \psi_{\ell_1} \times \cdots \times \psi_{\ell_k}$ to this code, we calculate $\psi_{\ell_i}((\cdots(v_1 \vee v_2) \vee \cdots) \vee v_k)$, for $i = 1, \ldots, k$. By Proposition 2.6.2 (1), we have

$$\begin{aligned}\psi_{\ell_i}((\cdots(v_1 \vee v_2) \vee \cdots) \vee v_k) &= \psi_{\ell_i}\lambda_{\ell_i}\lambda_{\ell_{i+1}}\cdots\lambda_{\ell_k}((\cdots(v_1 \vee v_2) \vee \cdots) \vee v_k) \\ &= \psi_{\ell_i}((\cdots(v_1 \vee v_2) \vee \cdots) \vee v_i) \\ &= v_i,\end{aligned}$$

as desired.

(2) By Definition 2.5.4, we have

$$\Psi(c) = \psi_{\ell_1}(c) <_Q \cdots <_Q \psi_{\ell_k}(c).$$

Applying Φ to this chain, we join vertices one at a time. Noting that $\psi_{\ell_1}(c) = \lambda_{\ell_1}(c)$, we use Proposition 2.6.2 (2) to calculate

$$\begin{aligned}\lambda_{\ell_i}(c) \vee \psi_{\ell_{i+1}}(c) &= \lambda_{\ell_i}\lambda_{\ell_{i+1}}(c) \vee \psi_{\ell_{i+1}}\lambda_{\ell_{i+1}}(c) \\ &= \lambda_{\ell_i}(\lambda_{\ell_{i+1}}(c)) \vee \psi_{\ell_{i+1}}(\lambda_{\ell_{i+1}}(c)) \\ &= \lambda_{\ell_{i+1}}(c).\end{aligned}$$

Thus, after $k-1$ join iterations, we recover c. □

2.7 Open questions

Since the class of balanced Cohen-Macaulay complexes contains so many widely studied classes of complexes, there are many possibilities to refine Theorem 1.4.1. In Theorem 2.4.1, we have required that Σ be an order complex of the form $\Delta(J(P))$, where P is a disjoint sum of chains. It is not known whether the theorem holds for more general classes of posets. (See [44], [48] for definitions in the questions that follow.)

Question 2.7.1. If P is a forest, then is there another poset Q such the h-vector of $J(P)$ is the f-vector of Q?

Question 2.7.2. If P is a series-parallel poset, then is there another poset Q such the h-vector of $J(P)$ is the f-vector of Q?

We conjecture that the answers to both questions are affirmative. In fact, we conjecture that the answer remains affirmative for *any* choice of a poset P.

Conjecture 2.7.1. *Let $J(P)$ be any distributive lattice. Then there is another poset Q such that the h-vector of $J(P)$ is the f-vector of Q.*

This conjecture has been tested by computer for all distributive lattices $J(P)$ arising from posets P having up to seven elements. Other open questions place requirements on Γ instead of on Σ.

Question 2.7.3. For which balanced Cohen-Macaulay complexes Σ is h_Σ the f-vector of a *graded* poset (or $(\mathbf{3}+\mathbf{1})$-free poset, or flag complex)?

To begin to answer Questions 2.7.1 - 2.7.3, it would be interesting to utilize any Eulerian permutation statistic *stat* to define posets such as Q in Definition 2.4.1 which satisfy the following two conditions.

1. Each k-element chain in Q corresponds to a permutation π in $\mathcal{L}(P)$ with $\mathrm{stat}(\pi) = k$.

2. For each poset P in some class $\mathcal{P}$, the statistics stat and des are equidistributed on $\mathcal{L}(P)$, so that $h_{J(P)} = f_Q$.

One might also consider a variation of this method based upon objects other than permutations, such as Motzkin paths or either of the tree representations in [44, pp. 23-25].

A result similar to Theorem 2.3.1 (in the sense that word rearrangements correspond to linear extensions of certain posets) states that the statistics INV and MAJ are equally distributed on the linear extensions of posets known as postorder labelled forests [7]. Perhaps Theorem 2.3.1 could be extended similarly.

Question 2.7.4. For what conditions on a poset are the statistics des and dmc equidistributed on $\mathcal{L}(P)$?

One might apply another variation of the method above by defining a rule which maps each n-element poset P to a subset $\mathcal{K}(P)$ of S_n which is *not* a set of linear extensions of P. This subset should have the property that the elements π in $\mathcal{K}(P)$ satisfying $\mathrm{stat}(\pi) = k$ are in bijective correspondence with $\mathcal{L}_k(P)$. In hopes of solving some of these open questions, we will study another Eulerian statistic in the following section.

Chapter 3

Joint distributions of permutation statistics on S_n

3.1 Introduction

Closely related to the Jordan-Hölder set of a poset is the permutation statistic INV which counts *inversions*. We define a pair (i, j) to be an inversion in a permutation π if $i < j$ and $\pi_i > \pi_j$. Thus,

$$\text{INV}(\pi) = \#\{(i, j) \,|\, i < j \text{ and } \pi_i > \pi_j\}.$$

Let P be an n-element poset. Redefining the Jordan-Hölder set as

$$\mathcal{L}(P) = \{\pi \in S_n \,|\, (i, j) \text{ an inversion in } \pi \Rightarrow \pi_i \not\succ_P \pi_j\},$$

we see that the study of linear extensions of a poset by descent is essentially the study of the *pair* of permutation statistics (des, INV).

In the past few decades, many results about permutation statistics have involved *joint* distributions on S_n of k-tuples of statistics. In 1954, Carlitz [10] essentially found a generating function for the joint distribution of (des, MAJ) [11]. This distribution has some particularly interesting symmetries. Foata [24] demonstrated the equidistribution of the pairs (des, INV) and (dmc, MAJ), and the symmetry of the distribution of the pair (INV, MAJ). Extending joint distribution results beyond pairs of statistics, Gessel [33] found a generating function for the joint distribution of $(\text{des}, \text{MAJ}, \text{INV})$, and Garsia and Gessel [30] found a generating function for the joint distribution of $(\text{des}, \text{ides}, \text{MAJ}, \text{IMAJ})$.

The statistic INV arises often in combinatorics, and an equidistribution result conspicuously absent from those above is one which states that for some Eulerian statistic *stat*, the pairs $(\text{stat}, \text{INV})$ and (des, MAJ) are equidistributed on S_n. We solve this well known problem in the following sections and state some related conjectures which may be of use in solving Conjecture 1.3.1.

3.2 Eulerian-Mahonian statistic pairs

The generating function for the number of permutations π in S_n with k inversions is the q-analog of n-factorial,

$$\sum_{\pi \in S_n} q^{\mathrm{INV}(\pi)} = (1+q)(1+q+q^2)\cdots(1+q+q^2+\cdots+q^{n-1}).$$

The permutation statistics distributed this way on S_n have been named *Mahonian statistics* [24], in honor of MacMahon. Following Clarke and Steingrímsson [14], we will write Eulerian statistics with lowercase letters and Mahonian statistics with capitals. Another Mahonian statistic is the *major index*, denoted MAJ. It is defined to be the sum of the descents of a permutation,

$$\mathrm{MAJ}(\pi) = \sum_{i \in D(\pi)} i.$$

The following table gives the distributions of INV and MAJ on S_n for $n = 1, \ldots, 5$.

$n\backslash k$	0	1	2	3	4	5	6	7	8	9	10
1	1										
2	1	1									
3	1	2	2	1							
4	1	3	5	6	5	3	1				
5	1	4	9	15	20	22	20	15	9	4	1

The generating function for the joint distribution of $(\mathrm{des}, \mathrm{MAJ})$ on S_n is given by Carlitz's q-Eulerian polynomial $B_n(t,q)$ [10]. (We follow the notation of Foata and Zeilberger [26].)

$$\sum_{\pi \in S_n} t^{\mathrm{des}(\pi)} q^{\mathrm{MAJ}(\pi)} = B_n(t,q).$$

Writing $B_n(t,q)$ as

$$B_n(t) = \sum_{k=0}^{n-1} B_{n,k}(q) t^k,$$

we may view the q-Eulerian polynomial as a polynomial in t whose coefficients are polynomials in q. Carlitz [10] showed that these coefficients satisfy the recurrence relation

$$B_{n,k}(q) = [k+1]_q B_{n-1,k}(q) + q^k B_{n-1,k-1}(q),$$

with initial conditions $B_{0,k}(q) = \delta_{0,k}$.

Example 3.2.1. The following table shows the joint distribution of $(\mathrm{des}, \mathrm{MAJ})$ on S_5. Note the interesting symmetries.

des\MAJ	0	1	2	3	4	5	6	7	8	9	10
0	1										
1		4	9	9	4						
2				6	16	22	16	6			
3							4	9	9	4	
4											1

A second pair of statistics, $(\mathrm{exc}, \mathrm{DEN})$ was conjectured by Denert [17] and shown by Foata and Zeilberger [26] and Han [35] to have the same joint distribution as $(\mathrm{des}, \mathrm{MAJ})$.

Theorem 3.2.1. *The pairs of permutation statistics* (exc, DEN) *and* (des, MAJ) *are equidistributed on* S_n,

$$\sum_{\pi \in S_n} t^{\mathrm{exc}(\pi)} q^{\mathrm{DEN}(\pi)} = \sum_{\pi \in S_n} t^{\mathrm{des}(\pi)} q^{\mathrm{MAJ}(\pi)} = B_n(t, q).$$

In other words,

$$\#\{\pi \in S_n \,|\, \mathrm{exc}(\pi) = k; \mathrm{DEN}(\pi) = p\} = \#\{\pi \in S_n \,|\, \mathrm{des}(\pi) = k; \mathrm{MAJ}(\pi) = p\}.$$

Thus the table in Example 3.2.1 which counts permutations in S_5 by descents and major index also counts permutations by excedances and *Denert's statistic*, DEN.

Denert's statistic is a Mahonian statistic with a slightly peculiar definition. Given a permutation $\pi = \pi_1 \cdots \pi_n$, we define the *Denert table* of π to be the word

$$\mathrm{dentab}(\pi) = d_1 \cdots d_n,$$

where d_i counts the number of indices $j < i$ such that π_j appears in the subword

$$\pi_i \cdots i$$

of the cyclic permutation

$$\pi_i \cdots n \cdot 1 \cdots (\pi_i - 1).$$

(Note that each letter d_i in the Denert table is no greater than $i-1$. We will consider such words in greater detail in the next section.) We define DEN to be the function which maps a permutation π to the sum of the components of its Denert table,

$$\mathrm{DEN}(\pi) = \sum_{i=1}^{n} d_i.$$

Foata and Zeilberger gave an equivalent definition which avoids calculation of the Denert table [26]. Let $E(\pi)$ be the excedance set of π. Then $\text{DEN}(\pi)$ is the sum of the elements of $E(\pi)$ plus the number of inversions (i, j) in π such that both i and j are excedances or both i and j are not excedances,

$$\text{DEN}(\pi) = \sum_{i \in E(\pi)} i + \text{INV}((\pi_i)_{i \in E(\pi)}) + \text{INV}((\pi_i)_{i \notin E(\pi)}).$$

Example 3.2.2. Let $\pi = 72835146$. Computing the Denert table, we have

$$\begin{pmatrix} \pi \\ \text{dentab}(\pi) \end{pmatrix} = \begin{pmatrix} 7\,2\,8\,3\,5\,1\,4\,6 \\ 0\,0\,1\,0\,0\,3\,2\,2 \end{pmatrix}.$$

The nonzero components of $\text{dentab}(\pi)$ are

$$\begin{aligned} d_3 &= 1, \text{ because 2 occurs in the word 8123,} \\ d_6 &= 3, \text{ because } 2, 3, 5 \text{ occur in the word 123456,} \\ d_7 &= 2, \text{ because } 5, 7 \text{ occur in the word 4567,} \\ d_8 &= 2, \text{ because } 7, 8 \text{ occur in the word 678.} \end{aligned}$$

Thus, $\text{DEN}(\pi) = 1 + 3 + 2 + 2 = 8$. Alternatively, we may calculate $\text{DEN}(\pi)$ by adding the excedances of π to the numbers of inversions in the subwords 78 and 235146,

$$\text{DEN}(\pi) = (1 + 3) + 0 + 4 = 8.$$

Since the statistic INV arises so often in combinatorics, one might hope for a natural Eulerian statistic *stat* such that the pair $(\text{stat}, \text{INV})$ is equidistributed on S_n with the pairs (des, MAJ) and (exc, DEN). There is in fact a natural Eulerian statistic with this property. We will call it *stc* and will define it in Section 3.4.

3.3 Codes and major index tables

Denote by E_n the set of words $w = w_1 \cdots w_n$ on the letters $\{0, \ldots, n-1\}$ in which each letter w_i is at most $n - i$,

$$E_n = \{w_1 \cdots w_n \in \mathbb{N}^n \mid w_i \leq n - i; \text{ for } i = 1, \ldots, n\}.$$

We will refer to the strictly decreasing word

$$(n-1) \cdot (n-2) \cdots 0$$

as the *stair word* of length n, and to the elements of E_n as *sub-stair words* of length n. Clearly the cardinality of the set E_n is $n!$, and bijections between E_n and S_n arise frequently in combinatorics.

One such bijection sends a permutation to its *code*. Recall from Section 2.1 that code(π) is the word $c_1 \cdots c_n$, where

$$c_i = \#\{j \in [n] \,|\, j > i; \pi_j < \pi_i\}.$$

We will denote the bijection by γ,

$$\begin{aligned} \gamma : S_n &\to E_n \\ \pi &\mapsto \text{code}(\pi). \end{aligned}$$

An important property of the code of a permutation is that the components of code(π) sum to $\text{INV}(\pi)$.

We pause here to comment that the definition of Dumont's statistic in terms of the code (see Section 2.1) is somewhat arbitrary. One would arrive at different Eulerian statistics *dmd, dmi* by counting distinct nonzero letters in the Denert table or inversion table ($= \text{code}(\pi^{-1})$).

Another bijection from S_n to E_n sends a permutation π to its *major index table*, a word whose components sum to $\text{MAJ}(\pi)$. To define the major index table, we will denote by $\pi^{(i)}$ the restriction of π to the letters $i, \ldots, n$. (e.g. if $\pi = 284367951$, then $\pi^{(4)} = 846795$, the restriction of π to the letters $4, \ldots, 9$.)

Definition 3.3.1. Let π be a permuation in S_n. Define the major index table of π to be the word

$$\text{majtable}(\pi) = m_1 \cdots m_n,$$

where

$$m_i = \begin{cases} 0, & \text{if } i = n, \\ \text{MAJ}(\pi^{(i)}) - \text{MAJ}(\pi^{(i+1)}), & \text{otherwise.} \end{cases}$$

If we imagine building the permutation π by inserting the letters $[n]$ in decreasing order, then m_i is the amount by which the major index increases with the insertion of i.

Example 3.3.2. Let $\pi = 284367951$. To calculate the major index table we build π one letter at a time in the order $9, \ldots, 1$ and record each increase in the major index. (Slashes in the following table indicate descents.)

i	$\pi^{(i)}$	$\text{MAJ}(\pi^{(i)})$	m_i
9	9	0	0
8	89	0	0
7	8/79	1	1
6	8/679	1	0
5	8/679/5	5	4
4	8/4679/5	6	1
3	8/4/3679/5	9	3
2	28/4/3679/5	12	3
1	28/4/3679/5/1	20	8

Thus we have $\text{majtable}(\pi) = m_1 \cdots m_9 = 833140100$.

It is not hard to see that for each permutation π in S_n, the word $\text{majtable}(\pi)$ belongs to E_n. (See Lemma 3.3.2.) Furthermore, the map taking a permutation to its major index table is known to be a bijection. This fact has been proved several times in the literature [25], perhaps first by Carlitz [11].

Theorem 3.3.1. *The map $\mu : S_n \to E_n$ defined by $\mu(\pi) = \text{majtable}(\pi)$ is a bijection.*

Proof. To invert μ we apply the following procedure to a word $m = m_1 \cdots m_n$ in E_n.

1. Define $w^{(n)}$ to be the one-letter word n.
2. For $i = n-1, \ldots, 1$, let $w^{(i)}$ be the unique word obtained by inserting the letter i into the word $w^{(i+1)}$ in such a way that $\text{MAJ}(w^{(i)}) - \text{MAJ}(w^{(i+1)}) = m_i$.
3. Set $\pi = w^{(1)}$.

It is clear that if the word π exists, then it satisfies $\pi^{(i)} = w^{(i)}$ for $i = 1, \ldots, n$, so that $\text{majtable}(\pi) = m$. By the following lemma, π does exist and is unique. In particular, given any permutation w on the letters $\{i+1, \ldots, n\}$ and any integer ℓ in the interval $\{0, \ldots, n-i\}$, then there is a unique permutation w' obtained by inserting the letter i into w in such a way that $\text{MAJ}(w') - \text{MAJ}(w) = \ell$. □

Lemma 3.3.2. *Let $\pi = \pi_1 \cdots \pi_{n-i}$ be a permutation on the letters $\{i+1, \ldots, n\}$, and suppose that π has k descents. Let $d_{k-1} < \cdots < d_0$ be the positions of these k descents, let $d_k = 0$, and let $a_{k+1} < \cdots < a_{n-i} = n - i$ be the remaining positions of π.*



Figure 3-1: Interpretation of the major index table of one permutation as the code of another.

1. *Let ℓ be an integer satisfying $0 \leq \ell \leq k$ and define π' to be the permutation obtained by inserting the letter i into position $d_\ell + 1$ of π. Then,*

$$\begin{aligned} \mathrm{des}(\pi') &= \mathrm{des}(\pi), \\ \mathrm{MAJ}(\pi') &= \mathrm{MAJ}(\pi) + \ell. \end{aligned}$$

2. *Let ℓ be an integer satisfying $k < \ell \leq n - i$ and define π' to be the permutation obtained by inserting the letter i into position $a_\ell + 1$ of π. Then,*

$$\begin{aligned} \mathrm{des}(\pi') &= \mathrm{des}(\pi) + 1, \\ \mathrm{MAJ}(\pi') &= \mathrm{MAJ}(\pi) + \ell. \end{aligned}$$

Proof. (1) The descent set of π' is

$$D(\pi') = \{d_{k-1}, \ldots, d_\ell, d_{\ell-1} + 1, \ldots, d_0 + 1\}.$$

(2) Let p be the least number such that $d_p < a_\ell$. Then, the descent set of π' is

$$D(\pi') = \{d_{k-1}, \ldots, d_p, a_\ell, d_{p-1} + 1, \ldots, d_0 + 1\},$$

and $a_\ell = (\ell - k) + (k - p) = \ell - p$. □

Combining the bijections γ and μ, we prove the equidistribution on S_n of MAJ and INV as a corollary. (See Figure 3-1.)

Corollary 3.3.3. *The permutation statistics* INV *and* MAJ *are equally distributed on S_n.*

Proof. Define the bijection $\phi : S_n \to S_n$ by $\phi = \gamma^{-1}\mu$. Since ϕ satisfies

$$\text{majtable}(\pi) = \text{code}(\phi(\pi)),$$

it also satisfies

$$\text{MAJ}(\pi) = \text{INV}(\phi(\pi)).$$

□

Surprisingly, the letter order $n, n-1, \ldots, 1$ preceding Definition 3.3.1 is not crucial for the construction of the major index table. In fact *any* letter order $\sigma_1, \ldots, \sigma_n$ induces a bijection $S_n \to E_n$ as in Theorem 3.3.1 [34]. Let us reconsider Theorem 3.3.1, Lemma 3.3.2, and Corollary 3.3.3 in terms of a more general major index table.

Fix a permutation $\sigma = \sigma_1 \cdots \sigma_n$ in S_n and denote by $\pi^{(i)}$ the restriction of π to the letters $\sigma_i, \ldots, \sigma_n$. (e.g. if $\sigma = 852739461$ and $\pi = 284367951$, then $\pi^{(4)} = 436791$, the restriction of π to the letters $\{\sigma_4, \ldots, \sigma_9\} = \{1, 3, 4, 6, 7, 9\}$.) We will call the sequence $\pi^{(1)}, \ldots, \pi^{(n)}$ defined in this way the *sequence of restricted permutations corresponding to* σ.

Definition 3.3.3. Fix a permutation σ in S_n. For any permutation π in S_n, construct the sequence $\pi^{(1)}, \ldots, \pi^{(n)}$ of restricted permutations corresponding to σ and define the σ-major index table of π to be the word

$$\sigma\text{-majtable}(\pi) = m_1 \cdots m_n,$$

where

$$m_i = \begin{cases} 0, & \text{if } i = n, \\ \text{MAJ}(\pi^{(i)}) - \text{MAJ}(\pi^{(i+1)}), & \text{otherwise.} \end{cases}$$

Note that if σ is the permutation $1 \cdots n$, then the sequence $\pi^{(1)}, \ldots, \pi^{(n)}$ has the same meaning as preceding Definition 3.3.1, and the σ-major index table is just the major index table.

Theorem 3.3.4. *Fix a permutation σ in S_n. The map $\mu_\sigma : S_n \to E_n$ defined by $\mu_\sigma(\pi) = \sigma$-majtable(π) is a bijection.*

Proof. Similar to the proof of Theorem 3.3.1. Use the following lemma instead of Lemma 3.3.2. □

Lemma 3.3.5. *Fix a permutation σ in S_n. Let π be a word on the letters $\{\sigma_{i+1}, \ldots, \sigma_n\}$, and suppose that π has k descents. Let $d_{k-1} < \cdots < d_0$ be the positions of these k descents, let $d_k = 0$, and let $d_{-1} = n - i$. Define the positions $d'_k < \cdots < d'_0$ by*

$$d'_\ell = \begin{cases} d_\ell, & \text{if } \pi_{d_\ell+1} > \sigma_i, \\ \max\{j \in [n-i] \,|\, d_\ell < j \leq d_{\ell-1}; \pi_j < \sigma_i\}, & \text{otherwise.} \end{cases}$$

Let $a'_{k+1} < \cdots < a'_{n-i}$ be the positions

$$\{0, 1, \ldots, n-i\} \setminus \{d'_0, \ldots, d'_k\}.$$

1. *Let ℓ be an integer satisfying $0 \leq \ell \leq k$ and define π' to be the permutation obtained by inserting the letter σ_i into position $d'_\ell + 1$ of π. Then,*

$$\begin{aligned} \mathrm{des}(\pi') &= \mathrm{des}(\pi), \\ \mathrm{MAJ}(\pi') &= \mathrm{MAJ}(\pi) + \ell. \end{aligned}$$

2. *Let ℓ be an integer satisfying $k < \ell \leq n - i$ and define π' to be the permutation obtained by inserting the letter σ_i into position $a_\ell + 1$ of π. Then,*

$$\begin{aligned} \mathrm{des}(\pi') &= \mathrm{des}(\pi) + 1, \\ \mathrm{MAJ}(\pi') &= \mathrm{MAJ}(\pi) + \ell. \end{aligned}$$

Proof. Identical to the proof of Lemma 3.3.2. □

As a corollary of Theorem 3.3.4, we have $n!$ bijections of the form

$$\phi_\sigma = \gamma^{-1}\mu_\sigma : S_n \to S_n$$

which satisfy

$$\mathrm{MAJ}(\pi) = \mathrm{INV}(\phi_\sigma(\pi))$$

and therefore prove that

$$\#\{\pi \in S_n \,|\, \mathrm{INV}(\pi) = k\} = \#\{\pi \in S_n \,|\, \mathrm{MAJ}(\pi) = k\}.$$

We shall see that the bijections $\{\phi_\sigma \,|\, \sigma \in S_n\}$ prove other equidistribution results as well.

It is not difficult to show that if $\sigma = \sigma_1 \cdots \sigma_n$ and $\sigma' = \sigma'_1 \cdots \sigma'_n$ are two different permutations, then the bijections μ_σ and $\mu_{\sigma'}$ are identical if and only if $\sigma_n = \sigma'_{n-1}$ and $\sigma_{n-1} = \sigma'_n$. Thus, the cardinality of the set $\{\phi_\sigma \,|\, \sigma \in S_n\}$ is really $\frac{n!}{2}$ rather than $n!$.

3.4 The statistic stc

We will define a simple Eulerian statistic stc which satisfies

$$\#\{\pi \in S_n \,|\, \mathrm{stc}(\pi) = k; \mathrm{INV}(\pi) = p\} = \#\{\pi \in S_n \,|\, \mathrm{des}(\pi) = k; \mathrm{MAJ}(\pi) = p\},$$

and will give a bijective proof of this fact in Theorem 3.4.2. To begin, we introduce a function $st : E_n \to \mathbb{N}$ and relate this function to the Eulerian numbers.

Definition 3.4.1. Define $st : E_n \to \mathbb{N}$ to be the function which maps a sub-stair word $v = v_1 \cdots v_n$ to the greatest number ℓ such that v contains a subsequence $v_{i_1}, \ldots, v_{i_\ell}$ which is (componentwise) strictly greater than the stair word of length ℓ,

$$v_{i_1} \cdots v_{i_\ell} > (\ell - 1) \cdot (\ell - 2) \cdots 0.$$

While v may contain several such subsequences of maximum length, identifying one and calculating this maximum length is quite easy. Starting from the rightmost position of v and reading left, we circle the first letter which is at least one, the next which is at least two, etc., until we cannot continue. The number of circled positions of v is then $st(v)$.

Example 3.4.2. Let $v = 245223010$. Starting from the right, we circle the 1, 3, 5, and 4. Thus, $st(v) = 4$. Note that $v_2 v_3 v_5 v_8 = 4531$ is strictly greater than 3210, the stair word of length 4.

Representing words as histograms, we see that $st(v)$ is the number of nonzero stairs which we can place beneath the histogram of v, preserving the order of these stairs. Figure 3-2 shows four nonzero stairs beneath the histogram of the word 245223010.

Let us count the words v in E_n satisfying $st(v) = k$, and denote these numbers by $\alpha(n, k)$. To derive a recursive formula for $\alpha(n, k)$, we note that

$$st(v_1 \cdot v_2 \cdots v_n) = \begin{cases} st(v_2 \cdots v_n) + 1, & \text{if } v_1 > st(v_2 \cdots v_n), \\ st(v_2 \cdots v_n), & \text{if } v_1 \leq st(v_2 \cdots v_n). \end{cases} \tag{3.1}$$

Thus we have the recursion

$$\alpha(n, k) = (k + 1)\alpha(n - 1, k) + (n - k - 1)\alpha(n - 1, k - 1),$$

and initial conditions

$$\alpha(1, k) = \begin{cases} 1, & \text{for } k = 0, \\ 0, & \text{otherwise.} \end{cases}$$

We see that the numbers $\alpha(n, k)$ are in fact the Eulerian numbers. (By similar reasoning one can show that the statistic dmc is Eulerian.)



Figure 3-2: A histogram interpretation of $st(245223010)$.

Observation 3.4.1. *The number of words v in E_n satisfying $st(v) = k-1$ is given by the Eulerian number $A(n,k)$.*

Composing the function st with the bijections γ and μ_σ from Section 3.3, we create a family of permutation statistics.

Definition 3.4.3. Define the permutation statistics stc and stm_σ by

$$\begin{aligned} \text{stc}(\pi) &= st(\text{code}(\pi)), \\ \text{stm}_\sigma(\pi) &= st(\sigma\text{-majtable}(\pi)). \end{aligned}$$

Observation 3.4.1 shows that each of these statistics is Eulerian; however, our main theorem reproves this fact bijectively without appealing to the observation. Furthermore, it shows that the statistic stc satisfies

$$\#\{\pi \in S_n \,|\, \text{stc}(\pi) = k; \text{INV}(\pi) = p\} = \#\{\pi \in S_n \,|\, \text{des}(\pi) = k; \text{MAJ}(\pi) = p\}.$$

We prove the main theorem with the bijection $\phi : S_n \to S_n$ from Corollary 3.3.3, which satisfies

$$\begin{aligned} \text{MAJ}(\pi) &= \text{INV}(\phi(\pi)), \\ \text{des}(\pi) &= \text{stc}(\phi(\pi)). \end{aligned}$$

Theorem 3.4.2. *The pairs of permutation statistics* (des, MAJ) *and* (stc, INV) *are equally distributed on S_n.*

Proof. Fix a permutation σ in S_n, and let $\phi_\sigma : S_n \to S_n$ be the bijection defined in Theorem 3.3.1. For every permutation π in S_n, the bijection ϕ_σ satisfies

$$\sigma\text{-majtable}(\pi) = \text{code}(\phi_\sigma(\pi)),$$

and therefore also satisfies

$$\begin{aligned} \text{MAJ}(\pi) &= \text{INV}(\phi_\sigma(\pi)), \\ \text{stm}_\sigma(\pi) &= \text{stc}(\phi_\sigma(\pi)). \end{aligned}$$

We claim that $\text{stm}_\sigma(\pi) = \text{des}(\pi)$.

Let $m = m_1 \cdots m_n$ be the σ-major index table of π, and let $\pi^{(1)}, \ldots, \pi^{(n)}$ be the sequence of restricted permutations corresponding to σ. Fix $i < n$ and assume that $st(m_{i+1} \cdots m_n) = \text{des}(\pi^{(i+1)})$. By equation (3.1), we have

$$st(m_i \cdot m_{i+1} \cdots m_n) = \begin{cases} st(m_{i+1} \cdots m_n) + 1, & \text{if } m_i > st(m_{i+1} \cdots m_n), \\ st(m_{i+1} \cdots m_n), & \text{otherwise.} \end{cases}$$

By Lemma 3.3.5, we have

$$\text{des}(\pi^{(i)}) = \begin{cases} \text{des}(\pi^{(i+1)}) + 1, & \text{if } m_i > \text{des}(\pi^{(i+1)}), \\ \text{des}(\pi^{(i+1)}), & \text{otherwise.} \end{cases}$$

Thus, $st(m_i \cdots m_n) = \text{des}(\pi^{(i)})$. Proceding by induction, we obtain $\text{stm}_\sigma(\pi) = \text{des}(\pi)$, as desired. □

3.5 The stc-set of a permutation

Theorem 3.4.2 states an analogy between the statistic pairs (des, MAJ) and (stc, INV). We will extend this result by associating a set to any permutation in such a way that the relationship of this set to the statistics stc and INV is analogous to that of the descent set to the statistics des and MAJ. Let $D : S_n \to 2^{[n-1]}$ be the set-valued function which maps a permutation to its descent set. Recall that D satisfies

$$\begin{aligned} |D(\pi)| &= \text{des}(\pi), \\ \sum_{i \in D(\pi)} i &= \text{MAJ}(\pi). \end{aligned}$$

In Definition 3.5.2 we will define an analogous set-valued function $STC : S_n \to 2^{[n-1]}$ which satisfies

$$\begin{aligned} |STC(\pi)| &= \text{stc}(\pi), \\ \sum_{i \in STC(\pi)} i &= \text{INV}(\pi). \end{aligned}$$

We will call $STC(\pi)$ the *stc-set* of π, and we will show that the functions D and STC are equally distributed on S_n.

Our strategy in defining $STC(\pi)$ for any permutation π is to transform the word code(π), whose components sum to $\text{INV}(\pi)$, into another word in which all the nonzero letters are *distinct* and sum to $\text{INV}(\pi)$. We begin by defining a map $\omega : E_n \to E_n$ which transforms a sub-stair word v into another word whose component sum is the same as that of v, and whose nonzero letters are strictly decreasing. We give an algorithmic description of the map ω.

Definition 3.5.1. Define the map $\omega : E_n \to E_n$ by performing the following procedure on a sub-stair word $v = v_1 \cdots v_n$.

For $i = n-2$ to 1, do

For $j = i$ to $n-2$, do

If $v_j \leq v_{j+1}$, and these letters are not both zero, then replace them in positions j and $j+1$ by the letters ℓ and ℓ', where

$$(\ell, \ell') = \begin{cases} (v_{j+1}+1, v_j - 1), & \text{if } v_j > 0, \\ (v_{j+1}, v_j), & \text{otherwise.} \end{cases} \tag{3.2}$$

The map ω is essentially a "bubble sort" algorithm, modified so that each time it exchanges two nonzero letters, it increments the letter moving left and decrements the letter moving right. Figure 3-3 shows the computation of ωv for $v = 332110$. Each line in the table represents a single iteration of the algorithm, with positions j and $j+1$ marked below by $\times$ if the corresponding letters are exchanged, and by)(otherwise.

Using Definition 3.5.1, one easily verifies for any word v in E_n with $\text{stc}(v) = k$ that the word $v' = \omega v$ also belongs to E_n and satisfies

$$\begin{aligned} v'_1 > \cdots > v'_k > 0, \\ v'_{k+1} = \cdots = v'_n = 0, \\ v'_1 + \cdots + v'_k = v_1 + \cdots + v_n. \end{aligned}$$

(These properties also follow from Theorem 3.5.1.) The map ω therefore naturally associates a subset of $[n-1]$ to each sub-stair word v in E_n: the set of nonzero letters of ωv.

Definition 3.5.2. Define the st-set of a sub-stair word v to be the set

$$ST(v) = \{\ell > 0 \,|\, \ell \text{ appears in } \omega v\}.$$



Figure 3-3: Computation of $\omega(332110)$.

Applying this definition to the code and major index tables of a permutation π, define the *stc-set* and *σ-stm-set* of π to be the sets

$$STC(\pi) = ST(\text{code}(\pi)),$$
$$\sigma\text{-}STM(\pi) = ST(\sigma\text{-majtable}(\pi)).$$

We claim that for every subset T of $[n-1]$, the number of permutations in S_n with stc-set T equals the number of permutations in S_n with descent set T. We prove this result with the bijection ϕ from Corollary 3.3.3 which satisfies

$$D(\pi) = STC(\phi(\pi)).$$

Theorem 3.5.1. *For every subset T of $[n-1]$, we have*

$$\#\{\pi \in S_n \,|\, D(\pi) = T\} = \#\{\pi \in S_n \,|\, STC(\pi) = T\}.$$

Proof. Fix a permutation σ in S_n, and let $\phi_\sigma : S_n \to S_n$ be the bijection defined in Theorem 3.3.1. For every permutation π in S_n, the bijection ϕ_σ satisfies

$$\sigma\text{-majtable}(\pi) = \text{code}(\phi_\sigma(\pi)),$$

and therefore also satisfies

$$\sigma\text{-}STM(\pi) = STC(\phi_\sigma(\pi)).$$

We claim that $\sigma\text{-}STM(\pi) = D(\pi)$.

Let $m = m_1 \cdots m_n$ be the σ-major index table of π, and let $\pi^{(1)}, \ldots, \pi^{(n)}$ be the sequence of restricted permutations corresponding to σ. Fix $i < n$ and let $d_0 > \cdots > d_{k-1}$ be the descents of $\pi^{(i+1)}$. Assume that

$$\omega(m_{i+1} \cdots m_n) = d_0 \cdots d_{k-1} \cdot 0 \cdots 0$$

so that we have

$$ST(m_{i+1} \cdots m_n) = D(\pi^{(i+1)}).$$

It is easy to see from Definition 3.5.1 that the map ω satisfies

$$\omega(m_i \cdots m_n) = \omega(m_i \cdot \omega(m_{i+1} \cdots m_n)).$$

Thus, we have

$$\begin{aligned}\omega(m_i \cdots m_n) &= \omega(m_i \cdot d_0 \cdots d_{k-1} \cdot 0 \cdots 0) \\ &= \begin{cases}(d_0+1)\cdots(d_{m_i-1}+1)\cdot d_{m_i}\cdots d_{k-1}\cdot 0\cdots 0, & \text{if } m_i \le k,\\ (d_0+1)\cdots(d_{j-1}+1)\cdot(m_i-j)\cdot d_j\cdots d_{k-1}\cdot 0\cdots 0, & \text{otherwise,}\end{cases}\end{aligned}$$

where j is the least integer satisfying $m_i - j > d_j$.

By Lemma 3.3.2, the nonzero letters in $\omega(m_i \cdots m_n)$ are precisely the descent set of $\pi^{(i)}$. Thus, we have $ST(m_i \cdots m_n) = \mathrm{des}(\pi^{(i)})$. Proceding by induction, we obtain $\sigma\text{-}STM(\pi) = D(\pi)$, as desired. □

Corollary 3.5.2. *Let π be a permutation in S_n with $\mathrm{stc}(\pi) = k$ and $\mathrm{INV}(\pi) = p$. Then the stc-set of π is a k-subset of $[n-1]$ whose elements sum to p.*

We formulate a second, algebraic, definition of the map ω as follows. Let G be a set of $n-2$ operators $\eta_1, \ldots, \eta_{n-2}$ on E_n, and define the action of η_i on a word $v = v_1 \cdots v_n$ by

$$\eta_i(v) = v_1 \cdots v_{i-1} \cdot \ell \cdot \ell' \cdot v_{i+2} \cdots v_n,$$

where

$$(\ell, \ell') = \begin{cases}(v_i, v_{i+1}) & \text{if } v_i > v_{i+1} \text{ or } v_i = v_{i+1} = 0,\\ (v_{i+1}+1, v_i - 1) & \text{if } 0 < v_i \le v_{i+1},\\ (v_{i+1}, v_i) & \text{if } 0 = v_i < v_{i+1}.\end{cases}$$

It is not difficult to see that the operators in the set G satisfy the relations of the 0-*Hecke monoid* on $n-2$ generators:

$$\begin{aligned} \eta_i^2 &= \eta_i, && \text{for } i = 1, \ldots, n, \\ \eta_i\eta_{i+1}\eta_i &= \eta_{i+1}\eta_i\eta_{i+1}, && \text{for } i = 1, \ldots, n-1, \\ \eta_i\eta_j &= \eta_j\eta_i, && \text{for } |i-j| \geq 2. \end{aligned}$$

(See [27].) Thus, an alternative definition of ω is

$$\omega = (\eta_{n-2}\eta_{n-3}\cdots\eta_1)\cdot(\eta_{n-2}\eta_{n-3}\cdots\eta_2)\cdots(\eta_{n-2}\eta_{n-3})\cdot(\eta_{n-2}).$$

3.6 More equidistribution results

Since Dumont's statistic has not enjoyed many applications since its definition, the problem of finding a Mahonian partner X for dmc has not received much attention. We will define such a statistic X and two additional statistics Y and Z in order to prove several joint distribution results.

Let $c = c_1 \cdots c_n$ be the code of a permutation π, and for each letter ℓ in $L(c)$, define the number $\xi(\ell)$ by

$$\xi(\ell) = \ell + \#\{i \,|\, c_i \geq \ell; \ell \text{ appears in } c_1 \cdots c_{i-1}\}.$$

Define $\mathrm{X}(\pi)$ to be the sum

$$\mathrm{X}(\pi) = \sum_{\ell \in L(c)} \xi(\ell).$$

Example 3.6.1. Let $\pi = 427196385$. Then, $\mathrm{code}(\pi) = 314042010$. For $\ell = 1, \ldots, 4$, row ℓ of the table below shows the contribution of each component of $\mathrm{code}(\pi)$ to $\xi(\ell)$.

$\mathrm{code}(\pi)$	3	1	4	0	4	2	0	1	0
contributions to $\xi(1)$	0	1	1	0	1	1	0	1	0
contributions to $\xi(2)$	0	0	0	0	0	2	0	0	0
contributions to $\xi(3)$	3	0	1	0	1	0	0	0	0
contributions to $\xi(4)$	0	0	4	0	1	0	0	0	0

Summing the contributions, we have $\mathrm{X}(\pi) = 5 + 2 + 5 + 5 = 17$.

To show that X is a Mahonian partner for Dumont's statistic, we introduce a bijection $\nu : S_n \to E_n$. By interpreting $\nu(\pi)$ as the code of another permutation, we create another bijection

$$\theta' = \gamma^{-1}\nu : S_n \to S_n,$$

which satisfies

$$\begin{aligned} E(\pi) &= LC(\theta'(\pi)), \\ \mathrm{exc}(\pi) &= \mathrm{dmc}(\theta'(\pi)), \\ \mathrm{DEN}(\pi) &= \mathrm{X}(\theta'(\pi)). \end{aligned}$$

(As we have remarked in Section 2.1, the equidistribution of E and LC on S_n is essentially due to Dumont [18]. For more information about this distribution, see [21].)

Definition 3.6.2. Define the map $\nu : S_n \to E_n$ by sending a permutation π with excedance table $e = e_1 \cdots e_n$ to the word $a = a_1 \cdots a_n$, where

$$a_{n-\pi_i+1} = e_i.$$

Note that this corresponds to rearranging the biword

$$\binom{\pi}{\mathrm{etab}(\pi)}$$

so that the top letters are in reverse order, and simply reading the resulting bottom word. By equation (2.4) the word $\nu(\pi)$ belongs to E_n.

Example 3.6.3. To apply ν to the permutation $\pi = 72835146$, we calculate

$$\binom{\pi}{\mathrm{etab}(\pi)} = \binom{7\ 2\ 8\ 3\ 5\ 1\ 4\ 6}{1\ 1\ 3\ 1\ 3\ 0\ 0\ 0},$$

and rearrange this biword as

$$\binom{8\ 7\ 6\ 5\ 4\ 3\ 2\ 1}{3\ 1\ 0\ 3\ 0\ 1\ 1\ 0}.$$

Thus, we have $\nu(\pi) = 31030110$.

We invert θ' using the procedure in the following proposition.

Proposition 3.6.1. *Let u be the word $n \cdot (n-1) \cdots 1$, let c be an element of E_n, and define the biword $y = \binom{u}{c}$. The following procedure produces the unique rearrangement $z = \binom{\rho}{e}$ of y satisfying $e = \mathrm{etab}(\rho)$.*

1. *For each letter ℓ in $L(c)$, find the least index i satisfying $c_i = \ell$, and define $z_\ell = y_i$. Let S be the set of such least indices, and define $T = [n] \setminus S$.*

2. *For each index $i \in T$, define*

$$d_i = \begin{cases} \#\{j \in S \mid c_j \leq c_i; u_j \geq u_i\}, & \text{if } c_i > 0, \\ 0, & \text{otherwise.} \end{cases}$$

3. *Let $(y_{\sigma^{-1}(i)})_{i\in T}$ be the unique rearrangement of $(y_i)_{i\in T}$ satisfying*

$$(d_{\sigma^{-1}(i)})_{i\in T} = \mathrm{code}((u_{\sigma^{-1}(i)})_{i\in T}).$$

4. *Insert the biletters $(y_{\sigma^{-1}(i)})_{i\in T}$ in order into the remaining positions of z.*

Proof. Similar to the proof of Proposition 2.3.3. Uniqueness of the rearrangement z follows from Proposition 2.3.4. □

Proposition 3.6.2. *The pairs of statistics* (dmc, X) *and* (des, MAJ) *are equidistributed on S_n.*

Proof. By Theorem 3.2.1, it suffices to show the equidistribution of $(\mathrm{dmc}, \mathrm{X})$ with $(\mathrm{exc}, \mathrm{DEN})$ Let $\pi = \pi_1 \cdots \pi_n$ be a permutation in S_n. The map θ' satisfies

$$\mathrm{exc}(\pi) = \mathrm{dmc}(\theta'(\pi)).$$

We claim that it also satisfies

$$\mathrm{DEN}(\pi) = \mathrm{X}(\theta'(\pi)).$$

Using (3.1) and the definition of excedance table, we may express $\mathrm{DEN}(\pi)$ as

$$\begin{aligned} \mathrm{DEN}(\pi) &= \sum_{i\in E(\pi)} i + \mathrm{INV}((\pi_i)_{i\in E(\pi)}) + \mathrm{INV}((\pi_i)_{i\notin E(\pi)}) \\ &= \sum_{i\in E(\pi)} i + \sum_{i\in E(\pi)} \#\{j \in E(\pi) \mid \pi_i > \pi_j; i \leq j\} + \sum_{i\in E(\pi)} \#\{j \notin E(\pi) \mid \pi_i > \pi_j; i \leq e_j\} \\ &= \sum_{i\in E(\pi)} i + \sum_{i\in E(\pi)} \#\{j \mid \pi_i > \pi_j; i \leq e_j\}. \end{aligned}$$

Letting $a = a_1 \cdots a_n = \nu(\pi)$, we may express $\mathrm{X}(\theta'(\pi))$ as

$$\mathrm{X}(\theta'(\pi)) = \sum_{i\in L(a)} i + \sum_{i\in L(a)} \#\{j \mid i \leq a_j; i \text{ appears in } a_1 \cdots a_{j-1}\}.$$

Since $E(\pi) = L(a)$, it suffices to show that for each letter i in $E(\pi)$, we have

$$\#\{j \mid \pi_i > \pi_j; i \leq e_j\} = \#\{j \mid i \leq a_j; i \text{ appears in } a_1 \cdots a_{j-1}\}.$$

This is easily demonstrated by the bijection $j \mapsto n - \pi_j + 1$. Since $e_j = a_{n-\pi_j+1}$, we have $i \leq e_j$ if and only if $i \leq a_{n-\pi_i+1}$. Further, the letter i first appears in the word a in position $n - \pi_i + 1$, which precedes position $n - \pi_j + 1$ if and only if $\pi_i > \pi_j$. □

The Mahonian statistic Y and the Eulerian statistic Z might most honestly be defined as the preimages of the statistics MAJ and des with respect to the map θ',

$$\begin{aligned} \mathrm{Y}(\pi) &= \mathrm{MAJ}(\theta'(\pi)), \\ \mathrm{Z}(\pi) &= \mathrm{des}(\theta'(\pi)). \end{aligned}$$

Nevertheless, we define the two statistics without referring to θ', in case they might aid in the bijective proof of another equidistribution result.

Let $e = e_1 \cdots e_n$ be the excedance table of a permutation π, and for each letter i, let $\epsilon(i)$ be $e_{\pi^{-1}(i)}$, the letter in e whose position is that of i in π. Define $\mathrm{Y}(\pi)$ to be the sum

$$\mathrm{Y}(\pi) = \sum_{i:\epsilon(i+1)>\epsilon(i)} (n-i+1),$$

and define $\mathrm{Z}(\pi)$ to be the cardinality

$$\mathrm{Z}(\pi) = \#\{i \,|\, \epsilon(i+1) > \epsilon(i)\}.$$

Proposition 3.6.3. *The pairs of statistics* (exc, Y) *and* (des, INV) *are equidistributed on* S_n.

Proof. By Theorem 2.3.1 and Proposition 3.6.2, it suffices to show the equidistribution of (exc, Y) with (dmc, MAJ). Let $\pi = \pi_1 \cdots \pi_n$ be a permutation and let $a = a_1 \cdots a_n$ be $\nu(\pi)$ so that $a = \mathrm{code}(\theta'(\pi))$. We simply note that $\epsilon(i+1)$ is greater than $\epsilon(i)$ if and only if $n-i+1$ is a descent in a. This is equivalent to $n-i+1$ being a descent in $\theta'(\pi)$. □

The same argument proves the equidistribution of the pairs (exc, Z) and (dmc, des). Thus, as a corollary of Proposition 3.6.2, we have an equidistribution result for two quadruples of statistics.

Corollary 3.6.4. *The two quadruples of permutation statistics* (exc, DEN, Y, Z) *and* (dmc, X, MAJ, des) *are equidistributed on* S_n.

3.7 Open equidistribution questions

Experience with the statistic stc suggests that a result much stronger than that of Theorem 3.4.2 is true.

Conjecture 3.7.1. *The following quadruples of permutation statistics are equidistributed on* S_n.

1. (des, MAJ, INV, stc)

2. (stc, INV, MAJ, des)

3. (dmc, X, MAJ, des)

4. (des, MAJ, X, dmc)

This conjecture has been tested on the symmetric groups up to S_{10}. Perhaps the Mahonian statistic r-MAJ [41], which generalizes MAJ and INV, will be of some use in constructing a bijective proof.

Since the stc-set and the descent set are equidistributed on S_n, perhaps it would be possible to use the statistic stc to modify the proof of Theorem 1.2.1 or to construct a poset analogous to that in the proof of Proposition 2.4.2 for a larger class of posets.

Question 3.7.1. Can we create a poset using major index tables, in which each k-element chain corresponds to a major index table m with $st(m) = k$?

Another strategy for constructing such a complex is to look for new interpretations of Mahonian statistics as we have for Eulerian statistics. The relationship of linear extensions to INV depends on the fact that an inversion of a permutation is a *pair* of numbers. The *inversion set* of a permutation, a subset of $\binom{[n]}{2}$, has no obvious analog for the other Mahonian statistics we have discussed.

Question 3.7.2. Is there a way to associate a MAJ-set (or X-set or DEN-set) to a permutation which is analogous to the inversion set?

If these sets could be defined easily, they might help associate a set $\mathcal{K}(P)$ of permutations to a poset P in the manner we have described at the end of Section 2.7.

Chapter 4

Posets whose chain polynomials have only real zeros

4.1 Interval characterizations

For nonnegative integers a and b, we denote by $\mathbf{a}+\mathbf{b}$ the poset which is the disjoint sum of an a-element chain and a b-element chain. A poset is called $(\mathbf{a}+\mathbf{b})$-*free* if it contains no induced subposet isomorphic to $\mathbf{a}+\mathbf{b}$. (See [44, Ch. 3] for basic definitions.) For example, the first two posets in Figure 4-1 are $\mathbf{2}+\mathbf{2}$ and $\mathbf{3}+\mathbf{1}$. The third poset P is $(\mathbf{2}+\mathbf{2})$-free but not $(\mathbf{3}+\mathbf{1})$-free, because the subposet induced by the elements $\{2,3,4,6\}$ is isomorphic to $\mathbf{3}+\mathbf{1}$.

Fishburn [22] characterized $(\mathbf{2}+\mathbf{2})$-free posets by showing that any such poset P may be represented as a set of closed intervals of real numbers $[c_i, d_i]$, ordered by

$$[c_i, d_i] <_P [c_j, d_j], \text{ if } d_i < c_j.$$

He also showed that for posets free of *both* $\mathbf{3}+\mathbf{1}$ and $\mathbf{2}+\mathbf{2}$, the intervals may be chosen to have unit length. In honor of these results, $(\mathbf{2}+\mathbf{2})$-free posets are often called *interval orders* and $(\mathbf{2}+\mathbf{2})$-free, $(\mathbf{3}+\mathbf{1})$-free posets are often called *unit interval orders*.

Unfortunately, little is known about posets free *only* of $\mathbf{3}+\mathbf{1}$. Characterization of $(\mathbf{3}+\mathbf{1})$-free posets is desirable, because several results and conjectures about posets



Figure 4-1: $\mathbf{2}+\mathbf{2}$, $\mathbf{3}+\mathbf{1}$, and a poset P which is not $(\mathbf{3}+\mathbf{1})$-free.

require avoidance only of $\mathbf{3}+\mathbf{1}$. For instance, Stanley's generalization of the chromatic polynomial [46] is known to be s-positive for the incomparability graphs of $(\mathbf{3}+\mathbf{1})$-free posets [31], and is conjectured to be e-positive for these graphs as well [13, 46, 49, 51, 54]. Further, the chain polynomial of a $(\mathbf{3}+\mathbf{1})$-free poset is known to have only real zeros. (See [49] and Corollary 4.6.1.)

4.2 Antiadjacency matrices of unit interval orders

Unit interval orders have many well known properties, and have several characterizations which avoid intervals and forbidden induced subposets. (See [16], [23], [55], [57].) One such characterization uses natural labellings and antiadjacency matrices. (See [55].) Let P be a labelled poset on n elements. Recall that any bijective function $\phi : P \to [n]$ is called a *labelling* of P, and is called *natural* if it satisfies $\phi(x) < \phi(y)$ whenever $x <_P y$. If the elements x and y are labelled as i and j, we will often write $i <_P j$ to mean $x <_P y$. For any labelled poset P, we define the *antiadjacency matrix* $A = [a_{ij}]$ of P by

$$a_{ij} = \begin{cases} 0, & \text{if } i <_P j, \\ 1, & \text{otherwise.} \end{cases}$$

Note that the entries on the diagonal of A are equal to one for any labelling of P. The entries below the diagonal are *all* equal to one if and only if P is naturally labelled.

Theorem 4.2.1. *Let P be a poset on n elements. P is a unit interval order if and only if it may be naturally labelled so that the corresponding antiadjacency matrix A satisfies*

$$a_{jk} \geq a_{i\ell}, \text{ for } 1 \leq i \leq j \leq n \text{ and } 1 \leq k \leq \ell \leq n. \tag{4.1}$$

Since A is a 0-1 matrix, (4.1) implies that the entries of A which are zero form a Ferrers shape in the upper right corner of A. From this fact, it is easy to see that unit interval orders are counted by Catalan numbers.

Example 4.2.1. Corresponding to any natural labelling of the unit interval order P in Figure 4-2 is the antiadjacency matrix

$$A = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix}.$$



P

Figure 4-2: a unit interval order.

To prove that a unit interval order may be labelled as in the theorem, we will begin with an arbitrary labelling of P and will *relabel* P to satisfy the theorem. This is equivalent to performing simultaneous row and column permutation on the antiadjacency matrix A which corresponds to the arbitrary labelling. To verify that a poset labelling satisfies the theorem, we will find it convenient to compare matrix columns and rows with vector inequalities. Let $a = (a_1, \ldots, a_n)$ and $b = (b_1, \ldots, b_n)$ be any two vectors in $\mathbb{R}^n$. We define the inequalities $a \leq b$ and $a \lneqq b$ by

$$\begin{aligned} a &\leq b \quad \text{if } a_i \leq b_i \text{ for } i = 1, \ldots, n. \\ a &\lneqq b \quad \text{if } a \leq b, \text{ and } a_i < b_i \text{ for some } i. \end{aligned}$$

Let us denote the ith row and ith column of any matrix M by M_{i*} and M_{*i}.

Observation 4.2.2. *The following two conditions on any real matrix M are equivalent.*

1. *It is possible to simlutaneously permute the columns and rows of M to obtain a matrix $M' = [m'_{ij}]$ such that the entries of M' weakly increase to the left in rows and downward in columns, i.e.,*

$$m'_{jk} \geq m'_{i\ell}, \text{ for } 1 \leq i \leq j \leq n \text{ and } 1 \leq k \leq \ell \leq n.$$

2. *The rows and columns of M corresponding to any pair of indices i and j satisfy one of the following pairs of vector inequalities.*

   (a) $M_{i*} \geq M_{j*}$ *and* $M_{*i} \leq M_{*j}$.

   (b) $M_{i*} \leq M_{j*}$ *and* $M_{*i} \geq M_{*j}$.

The first statement simply says that we may sort the columns of M in weakly decreasing order while simultaneously sorting the rows in weakly increasing order. With a moment's thought, we see that this is possible if and only if the conditions in the second statement are true.

We prove Theorem 4.2.1 with two propositions. Since the condition (4.1) fails to hold for nonnatural poset labellings, it will suffice to consider arbitrary labellings.

Proposition 4.2.3. *Let P be a labelled poset with antiadjacency matrix A. P is $(\mathbf{2}+\mathbf{2})$-free if and only if the columns of A are pairwise comparable as vectors.*

Proof. Suppose that two columns of A are incomparable as vectors. That is,

$$A_{*i} \not\geq A_{*j} \text{ and } A_{*i} \not\leq A_{*j},$$

for some indices $i \neq j$. Then for some indices $k \neq \ell$ we have

$$1 = a_{ik} > a_{i\ell} = 0 \text{ and } 0 = a_{jk} < a_{j\ell} = 1. \tag{4.2}$$

This implies that $i <_P k$, $j <_P \ell$, $i \not<_P \ell$ and $j \not<_P k$. The four elements must be distinct, and they induce a subposet of P which is isomorphic to $\mathbf{2}+\mathbf{2}$.

Conversely, if four elements i, j, k, and ℓ of P are related as above, then any labelling of P induces an antiadjacency matrix whose entries satisfy (4.2). The columns of this matrix cannot be pairwise comparable. □

Proposition 4.2.4. *Let P be a labelled $(\mathbf{2}+\mathbf{2})$-free poset with antiadjacency matrix A. P is $(\mathbf{3}+\mathbf{1})$-free if and only if the rows and columns of A corresponding to any pair of indices i and j satisfy one of the following pairs of vector inequalities.*

1. *$A_{i*} \geq A_{j*}$ and $A_{*i} \leq A_{*j}$.*
2. *$A_{i*} \leq A_{j*}$ and $A_{*i} \geq A_{*j}$.*

Proof. Suppose that have an incorrect pair of comparisons of the form

$$A_{i*} \gneqq A_{j*} \text{ and } A_{*i} \gneqq A_{*j}.$$

That is, P contains elements k and ℓ satisfying

$$1 = a_{i\ell} > a_{j\ell} = 0, \tag{4.3}$$
$$1 = a_{ki} > a_{kj} = 0. \tag{4.4}$$

Thus, we have $k <_P j <_P \ell$, and i must be incomparable to all three of these elements. The subposet induced by these elements is isomorphic to $\mathbf{3}+\mathbf{1}$.

Conversely, if four elements i, j, k, and ℓ are related in P as immediately above, then any labelling of P induces an antiadjacency matrix whose entries satisfy (4.3) and (4.4). The columns of this matrix do not satisfy our hypotheses. □

An alternative statement of the condition (4.1) relates unit interval orders to totally positive matrices. A real matrix, finite or infinite, is called *totally positive* (or sometimes *totally nonnegative*) if each $k \times k$ minor is nonnegative. (See [2].) Totally positive matrices have many interesting properties and arise frequently in combinatorics. (See [1], [9], [15] [28], [29], [50], [51], [52].) One important property of a finite square totally positive matrix is that all of its eigenvalues are nonnegative and real. (See [2, Thm. 1.1]).

An infinite matrix which is well known to be totally positive is

$$C = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdot & \cdot & \cdot \\ 1 & 1 & 0 & 0 & \cdot & \cdot & \cdot \\ 1 & 1 & 1 & 0 & \cdot & \cdot & \cdot \\ 1 & 1 & 1 & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & & & \\ \cdot & \cdot & \cdot & \cdot & & & \\ \cdot & \cdot & \cdot & \cdot & & & \end{bmatrix}. \tag{4.5}$$

(See [1].) It is easy to see that the matrices satisfying the conditions of Theorem 4.2.1 are finite generalized submatrices of C. That is, each such matrix is determined by a finite multiset of columns and a finite multiset of rows of C. The matrix A in Example 4.2.1 is the generalized submatrix of C corresponding to rows $(2, 2, 4, 4, 5)$ and columns $(1, 1, 3, 3, 5)$. Since any generalized submatrix of a totally positive matrix is itself totally positive, we see that any unit interval order has a labelling for which the corresponding antiadjacency matrix A is totally positive. It follows that the antiadjacency matrix corresponding to *any* labelling of a unit interval order has only positive real eigenvalues.

4.3 The squared antiadjacency matrix

If A is the antiadjacency matrix of a finite labelled poset P on n elements, then the *squared antiadjacency matrix* $B = A^2$ has a simple combinatorial interpretation. Let $G = (P, E)$ be the graph whose *adjacency* matrix is A. The vertex set of G consists of the elements $\{1, \ldots, n\}$ of P, and the edge set consists of the ordered pairs $(i, j) \in P \times P$ such that $i \not<_P j$. (See Figure 4-3.) Clearly, $B = [b_{ij}]$ counts paths of length two in G. That is, b_{ij} is the number of ordered triples (i, x, j), where (i, x) and (x, j) belong to E.

Observation 4.3.1. *Assume that P is $(\mathbf{3}+\mathbf{1})$-free, and let i, j, k, and ℓ be elements of P.*

1. *If $i <_P j <_P k$, then $b_{ik} = 0$.*
2. *If $b_{ik} > b_{i\ell}$, then there is an element $x <_P \ell$, such that $i \not<_P x \not<_P k$*
3. *If $b_{ik} > b_{jk}$, then there is an element $y >_P j$, such that $k \not>_P y \not>_P i$.*



Figure 4-3: A poset P and a graph G whose adjacency matrix is the *antiadjacency matrix* of P

Proof. (1) Assume $b_{ik} > 0$. Then, for some element x of P, (i, x) and (x, k) belong to E, implying that $i \not<_P x \not<_P k$. In fact, x must be incomparable to i and k, for if $x <_P i$, then we have $x <_P i <_P j <_P k$, and if $k <_P x$, then we have $i <_P j <_P k <_P x$, both impossible. Similarly, x cannot be comparable to j. Thus, the subposet of P induced by $\{i, j, k, x\}$ is isomorphic to $\mathbf{3} + \mathbf{1}$, contradicting our assumption that P is $(\mathbf{3} + \mathbf{1})$-free.

(2) If $b_{ik} > b_{i\ell}$, then there are more paths of length two in G from i to k than from i to ℓ. It follows that for some element x of P, the pairs (i, x) and (x, k) belong to E, and the pair (x, ℓ) does not.

(3) Apply the argument of (2) to the dual poset of P. □

Elements such as x in Observation 4.3.1 (2) are central to the proof of Lemma 4.3.2. To simplify notation, we introduce the following definition.

Definition 4.3.1. Let i, k, ℓ, and x be elements of P. Call x a (k, ℓ)-*advantage for* i if (i, x) and (x, k) are edges in the graph G and (x, ℓ) is not.

We use the word *advantage*, imagining that x helps us to travel from i to k, but not from i to ℓ. In terms of the poset relation $<_P$, x is a (k, ℓ)-advantage for i if $x <_P \ell$, and $i \not<_P x \not<_P k$. Note that in Figure 4-3, the vertex x is a (k, ℓ)-advantage for i, although it is not a (k, ℓ)-advantage for j.

Lemma 4.3.2. *Assume that P is $(\mathbf{3} + \mathbf{1})$-free, and let i, j, k, and ℓ be elements of P. If $b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell}$, then one of the following is true:*

1. *P contains an element x such that $j <_P x <_P \ell$ and $b_{j\ell} = 0$.*
2. *P contains an element y such that $i <_P y <_P k$ and $b_{ik} = 0$.*

Proof. Let us denote by $\alpha(k, \ell, i)$ the number of elements of P which are (k, ℓ)-advantages for i. Then we have

$$b_{ik} - b_{i\ell} = \alpha(k, \ell, i) - \alpha(\ell, k, i).$$

Assuming that $b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell}$, we have

$$\alpha(k, \ell, i) + \alpha(\ell, k, j) > \alpha(k, \ell, j) + \alpha(\ell, k, i),$$

and at least one of the following two inequalities must be true.

$$\alpha(k, \ell, i) > \alpha(k, \ell, j),$$
$$\alpha(\ell, k, j) > \alpha(\ell, k, i).$$

Suppose that $\alpha(k, \ell, i) > \alpha(k, \ell, j)$. Then P contains an element x which is a (k, ℓ)-advantage for i and not a (k, ℓ)-advantage for j. By Definition 4.3.1 the pairs (i, x) and (x, k) belong to E and the pairs (x, ℓ) and (j, x) do not. Thus, $j <_P x <_P \ell$ and by Observation 4.3.1 (1) we have $b_{j\ell} = 0$. Similarly, if $\alpha(\ell, k, j) > \alpha(\ell, k, i)$, then P contains an element y such that the pairs (y, k) and (i, y) do not belong to E. Thus, we have $i <_P y <_P k$ and $b_{ik} = 0$.

□

4.4 A characterization of $(\mathbf{3} + \mathbf{1})$-free posets

To characterize $(\mathbf{3} + \mathbf{1})$-free posets in a manner analogous to Theorem 4.2.1, we will use squared antiadjacency matrices.

Example 4.4.1. Corresponding to any natural labelling of the $(\mathbf{3} + \mathbf{1})$-free poset P in Figure 4-3 is the squared antiadjacency matrix

$$B = \begin{bmatrix} 2 & 2 & 0 & 0 & 0 \\ 2 & 2 & 0 & 0 & 0 \\ 4 & 4 & 2 & 2 & 0 \\ 4 & 4 & 2 & 2 & 0 \\ 5 & 5 & 3 & 3 & 1 \end{bmatrix}.$$

Theorem 4.4.1. *Let P be a poset on n elements. P is $(\mathbf{3} + \mathbf{1})$-free if and only if it may be naturally labelled so that the squared antiadjacency matrix B satisfies the following two conditions for all integers $1 \leq i \leq j \leq n$ and $1 \leq k \leq \ell \leq n$.*

1. *$b_{jk} \geq b_{i\ell}$.*
2. *If $b_{ik} - b_{i\ell} \neq b_{jk} - b_{j\ell}$, then $b_{i\ell} = 0$ and $b_{ik} < b_{jk} - b_{j\ell}$.*

Condition (1) of the theorem says that entries of the squared antiadjacency matrix increase to the left in rows and downward in columns. Following the proof of Theorem 4.2.1, we will prove that a $(\mathbf{3}+\mathbf{1})$-free poset P may be labelled to satisfy this condition by choosing an arbitrary labelling of P and by performing simultaneous row and column permutation on the corresponding squared antiadjacency matrix B.

Proposition 4.4.2. *Any $(\mathbf{3}+\mathbf{1})$-free poset may be naturally labelled so that the entries of its squared antiadjacency matrix weakly increase to the left in rows and downward in columns.*

Proof. Let P be a $(\mathbf{3}+\mathbf{1})$-free poset and assume that for each labelling of P, the corresponding antiadjacency matrix fails to satisfy condition (2) of Observation 4.2.2. (Trivially, condition (2) fails to hold for each non-natural labelling of P.) We consider two cases for a fixed labelling of P and the corresponding squared antiadjacency matrix B.

Case 1: Two columns of B are incomparable as vectors. That is,

$$B_{*i} \not\geq B_{*j} \text{ and } B_{*i} \not\leq B_{*j},$$

for some indices $i \neq j$. Then for some indices $k \neq \ell$ we have $b_{ik} > b_{i\ell}$ and $b_{jk} < b_{j\ell}$, implying that

$$b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell}.$$

Applying Lemma 4.3.2 to this inequality, we have $b_{j\ell} = 0$ or $b_{ik} = 0$, both contradictions. The argument for incomparable rows is identical.

Case 2: All columns of B are pairwise comparable as vectors, as are all rows, but for some indices $i \neq j$, we have an incorrect pair of comparisons of the form

$$B_{i*} \gneqq B_{j*} \text{ and } B_{*i} \gneqq B_{*j}.$$

That is, there are elements k and ℓ in P, not necessarily distinct, satisfying $b_{i\ell} > b_{j\ell}$ and $b_{ki} > b_{kj}$.

By Observation 4.3.1 (3), P contains an element $x <_P j$, such that $k \not<_P x \not<_P i$. By Observation 4.3.1 (2), P contains an element $y >_P j$, such that $i \not<_P y \not<_P \ell$. Thus, $x <_P j <_P y$ is a chain, and each of these three elements is incomparable to i. This contradicts our assumption that P is $(\mathbf{3}+\mathbf{1})$-free. □

It happens that any labelling of a $(\mathbf{3}+\mathbf{1})$-free poset which satisfies the first condition of Theorem 4.4.1 also satisfies the second condition.

Proposition 4.4.3. *Let P be a $(\mathbf{3}+\mathbf{1})$-free poset, naturally labelled so that its squared antiadjacency matrix B weakly increases to the left in rows and downward in columns.*

Let i, j, k, and ℓ be numbers satisfying $1 \leq i < j \leq n$ and $1 \leq k < \ell \leq n$. Then the 2×2 submatrix

$$\begin{bmatrix} b_{ik} & b_{i\ell} \\ b_{jk} & b_{j\ell} \end{bmatrix}$$

satisfies one of the following two conditions:

1. $b_{ik} - b_{i\ell} = b_{jk} - b_{j\ell}$.
2. $b_{i\ell} = 0$ *and* $b_{ik} < b_{jk} - b_{j\ell}$.

Proof. Suppose that condition (1) is not satisfied.
Case 1: ($b_{ik} - b_{i\ell} > b_{jk} - b_{j\ell}$.) We apply Lemma 4.3.2 to this inequality. If $b_{j\ell} = 0$, then $b_{i\ell} = 0$ and $b_{ik} > b_{jk}$, contradicting our assumptions about weakly increasing entries of B. If instead $b_{ik} = 0$, then $b_{i\ell} = 0$ and $b_{j\ell} > b_{jk}$, another contradiction.
Case 2: ($b_{ik} - b_{i\ell} < b_{jk} - b_{j\ell}$.) Again we apply Lemma 4.3.2. If $b_{jk} = 0$, then all four numbers are zero, a contradiction. We conclude that $b_{i\ell} = 0$ and that condition (2) is satisfied. □

Finally, we show that the only posets satisfying the conditions of Theorem 4.4.1 are those which are $(\mathbf{3}+\mathbf{1})$-free.

Proposition 4.4.4. *Let P be a labelled poset containing $\mathbf{3}+\mathbf{1}$ as an induced subposet, and let B be its squared antiadjacency matrix. Then there are two distinct elements i and k such that*

$$b_{ik} \neq 0 \text{ and } b_{ii} - b_{ik} \neq b_{ki} - b_{kk}.$$

Proof. Let $1, 2, 3$, and 4 be four elements of P such that $1 <_P 2 <_P 3$ is a chain, and 4 is incomparable to $1, 2$, and 3. Let $G = (P, E)$ be the graph defined in Section 4.3.

Clearly, $b_{13} \neq 0$, since $(1, 4)$ and $(4, 3)$ are edges in G. We claim that

$$b_{11} - b_{13} \neq b_{31} - b_{33}.$$

Define the sets

$$X = \{x \in P \,|\, (1, x) \in E, (x, 1) \in E, (x, 3) \notin E\},$$
$$Y = \{x \in P \,|\, (x, 1) \in E, (x, 3) \notin E\},$$

and note that

$$|X| = b_{11} - b_{13},$$
$$|Y| = b_{31} - b_{33}.$$

Certainly X is a subset of Y. Moreover it is a proper subset, since the element 2 belongs to Y and not to X. Thus we have $b_{11} - b_{13} < b_{31} - b_{33}$. □

As we have considered Theorem 4.2.1 in terms of totally positive matrices, we now consider Theorem 4.4.1 in terms of totally positive matrices. The infinite matrix D,

$$D = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdot & \cdot & \cdot \\ 2 & 1 & 0 & 0 & \cdot & \cdot & \cdot \\ 3 & 2 & 1 & 0 & \cdot & \cdot & \cdot \\ 4 & 3 & 2 & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & & & \\ \cdot & \cdot & \cdot & \cdot & & & \\ \cdot & \cdot & \cdot & \cdot & & & \end{bmatrix},$$

is in some sense the square of the infinite matrix C from (4.5). It too is totally positive. It is easy to see that the matrices satisfying the conditions of Theorem 4.4.1 are generalized submatrices of D: each is determined by a finite multiset of columns and a finite multiset of rows. The matrix B in Example 4.4.1 is the generalized submatrix of D corresponding to rows $(2, 2, 4, 4, 5)$ and columns $(1, 1, 3, 3, 5)$. Since any generalized submatrix of a totally positive matrix is itself totally positive, we see that any $(\mathbf{3}+\mathbf{1})$-free poset has a labelling for which the corresponding squared antiadjacency matrix B is totally positive. It follows that the squared antiadjacency matrix corresponding to *any* labelling of a $(\mathbf{3}+\mathbf{1})$-free poset has only positive real eigenvalues, and that the antiadjacency matrix has only real eigenvalues.

Since Theorem 4.2.1 implies that the Catalan numbers count unit interval orders, it is natural to ask if the observation above will yield an analogous result for $(\mathbf{3}+\mathbf{1})$-free posets. No such formula is currently known.

Question 4.4.2. Does Theorem 4.4.1 imply a simple formula for the number of $(\mathbf{3}+\mathbf{1})$-free posets on n elements?

4.5 The altitude sequence

For each $(\mathbf{3}+\mathbf{1})$-free poset P, it is simple to construct a labelling ϕ which satisfies the conditions of Theorem 4.4.1, or of Theorem 4.2.1 if P is a unit interval order. We do so using the principal order ideals and dual principal order ideals of the elements of P.

For any element x in a poset P, we will denote the corresponding principal order ideal and principal dual order ideal by Λ_x and V_x, respectively.

$$\Lambda_x = \{y \in P \mid y \leq_P x\},$$
$$V_x = \{y \in P \mid y \geq_P x\}.$$

It is easy to verify that if P is $(\mathbf{3}+\mathbf{1})$-free, then its order ideals satisfy the following two properties.

1. If $|\Lambda_x| > |\Lambda_y|$, then $|V_x| \leq |V_y|$.
2. If $|V_x| > |V_y|$, then $|\Lambda_x| \leq |\Lambda_y|$.

For each element x of P, we will define its *altitude* to be the difference in cardinality between its principal order ideal and principal dual order ideal.

$$\alpha(x) = |\Lambda_x| - |V_x|.$$

If the element x is labelled as $\phi(x) = i$, we will write Λ_i, V_i, and $\alpha(i)$ in place of Λ_x, V_x, and $\alpha(x)$.

Example 4.5.1. Let P be the poset in Figure 4-3. Then we have

$$\begin{aligned} \alpha(j) = \alpha(k) &= -3, \\ \alpha(i) = \alpha(x) &= 1, \\ \alpha(\ell) &= 4. \end{aligned}$$

We will say that a labelling ϕ of P *respects altitude* if it satisfies $\phi(x) < \phi(y)$ whenever $\alpha(x) <_P \alpha(y)$. A labelling which respects altitude is necessarily natural.

By the following proposition, the labellings of unit interval orders which satisfy Theorem 4.2.1 are precisely those which respect altitude.

Proposition 4.5.1. *Let P be a unit interval order labelled by ϕ and let A be the corresponding antiadjacency matrix. The matrix A satisfies the conditions of Theorem 4.2.1 if and only if the labelling ϕ respects altitude.*

Proof. Suppose that P has been labelled as in Theorem 4.2.1, and let x and y be two elements satisfying $\phi(x) < \phi(y)$. Then we have

$$A_{x*} \leq A_{y*} \text{ and } A_{*x} \geq A_{*y}.$$

Equivalently, we have

$$|V_x| \geq |V_y| \text{ and } |\Lambda_x| \leq |\Lambda_y|. \tag{4.6}$$

Combining these inequalities, we obtain $\alpha(x) \leq \alpha(y)$.

Now suppose that ϕ is a natural labelling of P such that the corresponding antiadjacency matrix A does not satisfy (4.1). Then for some labels $i < j$ we have

$$A_{i*} \gneqq A_{j*} \text{ or } A_{*i} \lneqq A_{*j}.$$

Equivalently, we have

$$|V_i| < |V_j| \text{ or } |\Lambda_i| > |\Lambda_j|.$$

If $|V_i| < |V_j|$, then we must have $|\Lambda_i| \geq |\Lambda_j|$. Therefore,

$$\alpha(i) - \alpha(j) = |\Lambda_i| - |V_i| - |\Lambda_j| + |V_j| > 0,$$

and ϕ does not respect altitude. Similarly if $|\Lambda_i| > |\Lambda_j|$, then we must have $|V_i| \leq |V_j|$. Again, ϕ does not respect altitude. □

A result analogous to Proposition 4.5.1 holds for $(\mathbf{3}+\mathbf{1})$-free posets.

Proposition 4.5.2. *Let P be $(\mathbf{3}+\mathbf{1})$-free poset labelled by ϕ, and let B be the corresponding squared antiadjacency matrix. The matrix B satisfies the conditions of Theorem 4.4.1 if and only if the labelling ϕ respects altitude.*

Proof. Suppose that P has been labelled as in Theorem 4.4.1, and let x and y be two elements satisfying $\phi(x) < \phi(y)$. Then we have

$$B_{x*} \leq B_{y*} \text{ and } B_{*x} \geq B_{*y}.$$

In particular, if w is any minimal element of P and z is any maximal element, then we have

$$b_{xw} \leq b_{yw} \text{ and } b_{zx} \geq b_{zy}.$$

This implies that

$$|V_x| \geq |V_y| \text{ and } |\Lambda_x| \leq |\Lambda_y|. \tag{4.7}$$

Combining these inequalities, we obtain $\alpha(x) \leq \alpha(y)$.

Now suppose that ϕ is a natural labelling of P such that the corresponding squared antiadjacency matrix B does not satisfy the conditions of Theorem 4.4.1. Then for some labels $i < j$ we have

$$B_{i*} \gneqq B_{j*} \text{ or } B_{*i} \lneqq B_{*j}.$$

If $B_{*i} \lneqq B_{*j}$, then for some element z of P, we have $b_{zi} < b_{zj}$. By Theorem 4.4.1 (2), we may assume that z is maximal in P. Thus, we have $|\Lambda_i| > |\Lambda_j|$. It follows that $|V_i| \leq |V_j|$. Therefore,

$$\alpha(i) - \alpha(j) = |\Lambda_i| - |V_i| - |\Lambda_j| + |V_j| > 0,$$

and ϕ does not respect altitude. Similarly, if $B_{i*} \gneqq B_{j*}$, then we have $|V_i| < |V_j|$ and $|\Lambda_i| \geq |\Lambda_j|$. Again, ϕ does not respect altitude. □

Thus, to demonstrate that a poset is a unit interval order (is $(\mathbf{3}+\mathbf{1})$-free), it suffices to examine the (squared) antiadjacency matrix corresponding to any altitude respecting labelling of P.

Let us define the *altitude sequence* of any poset P to be the sequence of n altitudes, written in weakly increasing order. Is not difficult to show that a unit interval order is uniquely identified by its altitude sequence, although this fact does not hold for posets in general. Furthermore, we conjecture the analogous stronger result for $(\mathbf{3}+\mathbf{1})$-free posets.

Conjecture 4.5.3. *Two $(\mathbf{3}+\mathbf{1})$-free posets P and Q have the same altitude sequence if and only if they are isomorphic.*

If this conjecture is true, one might use altitude sequences to count $(\mathbf{3}+\mathbf{1})$-free posets.

Question 4.5.2. Is there a simple characterization of the altitude sequences arising from $(\mathbf{3}+\mathbf{1})$-free posets on n elements, or a simple formula counting such sequences?

Another conjecture relates the altitude sequences of unit interval orders to pairs of permutations. The following statement of the conjecture, due to Postnikov [40], differs somewhat from its original form due to Kostant [38].

Conjecture 4.5.4. *Let P be a unit interval order on n elements. Then there are two permutations π, σ in S_n such that the vector $(\pi_1-\sigma_1,\ldots,\pi_n-\sigma_n)$ is the altitude sequence of P.*

This conjecture is trivially true for the class of posets of dimension 2 (see [55]), and thus holds for all series-parallel posets and many unit interval orders. The analogous statement involving $(\mathbf{3}+\mathbf{1})$-free posets is not true.

4.6 $(\mathbf{3}+\mathbf{1})$-free posets and distributive lattices

While $(\mathbf{3}+\mathbf{1})$-free posets and distributive lattices have little in common by definition, the following corollary of Theorem 4.4.1 suggests that there may be a connection between the two classes.

By a result of Stanley [47], we may use the antiadjacency matrix of a poset to determine whether or not the chain polynomial has only real zeros. Let P be an arbitrarily labelled poset with antiadjacency matrix A. Then we have

$$f_P(x) = \det(I + xA). \tag{4.8}$$

From this formula we see that $f_P(x)$ has only real zeros if and only if A has only real eigenvalues. Applying the discussion following Proposition 4.4.4 to Theorem 4.4.1, we obtain the following corollary. (This result was originally proved in [49, Cor. 2.9], using facts about symmetric functions [31], [49, Thm. 2.8].)

Corollary 4.6.1. *Let P be a $(\mathbf{3}+\mathbf{1})$-free poset. Then the chain polynomial $f_P(x)$ has only real zeros.*

Recall that Stanley and Neggers have conjectured (Conjecture 1.3.1) the similar statement that the chain polynomial of any distributive lattice has only real zeros. Unfortunately, it is not in general possible to label a distributive lattice so that some power of its antiadjacency matrix is totally positive. However, if this were the case for some subclass of distributive lattices, we would have a combinatorial proof that the zeros of the corresponding chain polynomials were roots of unity.

Question 4.6.1. Let A be the antiadjacency matrix of a labelled distributive lattice $J(P)$. Under what conditions is there a power $k > 0$ for which A^k is totally positive?

Noting that two posets which are not isomorphic can have the same f-vector, we arrive at another possiblility of applying Corollary 4.6.1 to Conjecture 1.3.1.

Question 4.6.2. For which distributive lattices $J(P)$ is there a $(\mathbf{3}+\mathbf{1})$-free poset Q such that $f_{J(P)} = f_Q$ or $h_{J(P)} = f_Q$?

There is some evidence in favor of an affirmative answer to the second equality in Question 4.6.2, and therefore in favor of Conjecture 2.7.1. Namely, for all posets P having seven or fewer elements, there is at least one poset Q satisfying

$$f_Q = h_{J(P)}. \tag{4.9}$$

This remains true if we require Q to be a $(\mathbf{3}+\mathbf{1})$-poset, or even a unit interval order. In fact, for a given distributive lattice $J(P)$ with h-vector $h_{J(P)} = (h_0, h_1, \ldots, h_n)$, the number of h_1-element unit interval orders Q satisfying (4.9) is often quite large. In the following table we partition the set

$$\{J(P) \,|\, \#P \leq 7\}$$

into blocks, according to the value of h_1 in each h-vector $h_{J(P)}$. Column two of the table gives the number of unit interval orders Q which satisfy (4.9), averaged over the distinct h-vectors in each block.

h_1	average number of unit interval orders Q satisfying $f_Q = h_{J(P)}$
1	1
2	1
3	1.33
4	2
5	2.9
6	3.53
7	6
8	8
9	13.43
10	18.51
11	25.97
12	40.36
13	58.53
14	98.53
15	111.97
16	237.72
17	263.30
18	507.75
19	493.10
20	1088.03

Note that the average number of unit interval orders Q which satisfy (4.9) is a lower bound for the average number of $(\mathbf{3}+\mathbf{1})$-posets which satisfy (4.9), and a very weak lower bound for the average number of *abritrary* posets which satisfy (4.9). Interestingly, there is no equally strong evidence supporting the first equality in Question 4.6.2. Many distributive lattices $J(P)$ exist for which $f_{J(P)}$ is not the f-vector of a unit interval order, although $f_{J(P)}(x)$ does have only real zeros.

This raises some questions about the generality of chain polynomials among polynomials which have only real zeros. Let $\mathcal{F}$ be the class of polynomials $f(x)$ in $\mathbb{P}[x]$ which have only real zeros and which satisfy $f(0) = 1$,

$$\mathcal{F} = \{ f(x) = 1 + a_0 x + \cdots + a_{n-1} x^n \in \mathbb{P}[x] \mid \text{ all zeros of } f(x) \text{ are real } \}.$$

Let $\mathcal{F}_1$ be the subset of $\mathcal{F}$ for which $(1, a_0, \ldots, a_{n-1})$ is the f-vector of some poset. Let $\mathcal{F}_2$ be the subset of $\mathcal{F}_1$ for which $(1, a_0, \ldots, a_{n-1})$ is the f-vector of some $(\mathbf{3}+\mathbf{1})$-free poset.

Question 4.6.3. Which, if any, of the inclusions $\mathcal{F}_2 \subset \mathcal{F}_1 \subset \mathcal{F}$ are strict?

Note that if we had the equality $\mathcal{F}_2 = \mathcal{F}_1$, then this would imply a partial converse to Corollary 4.6.1. If on the other hand we had the equality $\mathcal{F}_1 = \mathcal{F}$, then Conjecture 1.3.1 would imply Conjecture 2.7.1.

Bibliography

[1] M. Aissen, I. J. Schoenberg, and A. Whitney. On generating functions of totally positive sequences. *J. Anal. Math.* **2**, 93–103 (1952).

[2] T. Ando. Totally positive matrices. *Linear Algebra Its Appl.* **90**, 165–219 (1987).

[3] M. M. Bayer and L. J. Billera. Counting faces and chains in polytopes and posets. *Contemp. Math.* **34**, 207–252 (1984).

[4] L. J. Billera and A. Björner. Face numbers of polytopes and complexes. In J. E. Goodman and J. O'Rourke, eds., *Handbook of Discrete and Computational Geometry*, pp. 291–310. CRC Press, Boca Raton/New York (1997).

[5] L. J. Billera and G. Hetyei. Linear inequalities for flags in graded partially ordered sets (1998). to appear.

[6] A. Björner, P. Frankl, and R. Stanley. The number of faces of balanced Cohen-Macaulay complexes and a generalized Macaulay theorem. *Combinatorica* **7**, 23–34 (1987).

[7] A. Björner and M. Wachs. Permutation Statistics and Linear Extensions of Posets. *J. Combin. Theory Ser. A* **58**(1), 85–114 (1991).

[8] F. Brenti. *Unimodal, Log-Concave, and Pólya Frequency Sequences in Combinatorics*. No. 413 in Mem. Amer. Math. Soc. American Mathematical Society, Providence, RI (1989).

[9] F. Brenti. Combinatorics and total positivity. *J. Combin. Theory Ser. A* **71**(2), 175–218 (1995).

[10] L. Carlitz. q-Bernoulli and Eulerian numbers. *Trans. of the AMS* **76**, 332–350 (1954).

[11] L. Carlitz. A combinatorial property of q-Eulerian numbers. *American Math. Monthly* **82**, 51–54 (1975).

[12] M. Chari. Matroid inequalities. *Disc. Math.* **147**, 283–286 (1995).

[13] T. Chow. Descents, quasi-symmetric functions, Robinson-Schensted for posets, and the chromatic symmetric function. *J. Alg. Combin.* **10**, 227–240 (1999).

[14] R. J. Clarke, E. Steingrímsson, and J. Zeng. New Euler-Mahonian statistics on permutations and words. *Adv. Appl. Math.* **18**, 237–270 (1997).

[15] C. W. Cryer. Some properties of totally positive matrices. *Lin. Alg. Its Appl.* **15**, 1–25 (1976).

[16] R. A. Dean and G. Keller. Natural partial orders. *Canad. J. Math* **20**, 535–554 (1968).

[17] M. Denert. The genus zeta function of hereditary orders in central simple algebras over global fields. *Math. Comp.* **54**, 449–465 (1990).

[18] D. Dumont. Interprétations combinatoires des nombres de Genocchi. *Duke Math. J.* **41**, 305–318 (1974).

[19] A. Duval. A combinatorial decomposition of simplicial complexes. *Israel J. Math.* **87**, 77–87 (1994).

[20] P. Edelman and V. Reiner. H-shellings and h-complexes. *Adv. Math.* **106**, 36–64 (1994).

[21] R. Ehrenborg and E. Steingrímsson. The excedance set of a permutation. *preprint* (2000).

[22] P. C. Fishburn. Intransitive indifference with unequal indifference intervals. *J. Math. Psych.* **7**, 144–149 (1970).

[23] P. C. Fishburn. *Interval Graphs and Interval Orders*. Wiley, New York (1985).

[24] D. Foata. Distributions Eulériennes et Mahoniennes sur le groupe des permutations. In M. Aigner, ed., *Higher Combinatorics*, vol. 19, pp. 27–49. D. Reidel Publishing Combany, Dordrecht-Holland (1977).

[25] D. Foata (1999). Personal communication.

[26] D. Foata and D. Zeilberger. Denert's permutation statistic is indeed Euler-Mahonian. *Studies in Appl. Math.* **83**, 31–59 (1990).

[27] S. Fomin and C. Greene. Noncommutative Schur functions and their applications. *Disc. Math.* pp. 179–200 (1998).

[28] S. Fomin and A. Zelevinsky. Double bruhat cells and total positivity. *J. Amer. Math. Soc.* **12**, 335–380 (1999).

[29] S. Fomin and A. Zelevinsky. Total positivity: Tests and parametrizations. *Math. Intelligencer* (To appear).

[30] A. M. Garsia and I. Gessel. Permutation statistics and partitions. *Adv. Math.* **31**(3), 288–305 (1979).

[31] V. Gasharov. Incomparability graphs of $(\mathbf{3}+\mathbf{1})$-free posets are s-positive. *Disc. Math.* **157**, 211–215 (1996).

[32] V. Gasharov. On the Neggers-Stanley conjecture and the Eulerian polynomials. *J. Combin. Theory Ser. A* **82**(2), 134–146 (1998).

[33] I. Gessel. *Generating functions and enumeration of sequences*. Ph.D. thesis, MIT, Cambridge, MA (1977).

[34] I. Gessel (1999). Personal communication.

[35] G. N. Han. Une transformation fondamentale sur les réarrangements de mots. *Adv. Math.* **105**, 26–41 (1994).

[36] T. Hibi. Linear and nonlinear inequalities concerning a certain combinatorial sequence which arises from counting the number of chains of a finite distributive lattice. *Disc. Appl. Math.* **34**, 145–150 (1991).

[37] D. M. Jackson and I. P. Goulden. A formal calculus for the enumerative system of sequences-i. combinatorial theorems. *Studies in Applied Mathematics* **61**, 141–178 (1979).

[38] B. Kostant (2000). Personal communication.

[39] J. Neggers. Representations of finite partially ordered sets. *J. Combin. Inform. System Sci.* **3**, 113–133 (1978).

[40] A. Postnikov (2000). Personal communication.

[41] D. Rawlings. The r-major index. *J. Combin. Theory Ser. A* **31**, 175–183 (1981).

[42] R. Simion. A multiindexed Sturm sequence of polynomials and unimodality of certain combinatorial sequences. *J. Combin. Theory Ser. A* **36**, 15–22 (1984).

[43] R. Stanley. Balanced Cohen-Macaulay complexes. *Trans. of the AMS* **241**(1), 139–157 (1979).

[44] R. Stanley. *Enumerative Combinatorics*, vol. 1. Wadsworth & Brooks/Cole, Belmont, CA (1986).

[45] R. Stanley. Log-concave and unimodal sequences in algebra, combinatorics, and geometry. In *Graph Theory and its Applications: East and West*, vol. 576, pp. 500–535. Annals of the New York Academy of Sciences (December 29, 1989).

[46] R. Stanley. A symmetric function generalization of the chromatic polynomial of a graph. *Adv. Math.* **111**, 166–194 (1995).

[47] R. Stanley. A matrix for counting paths in acyclic digraphs. *J. Combin. Theory Ser. A* **74**, 169–172 (1996).

[48] R. Stanley. *Combinatorics and Commutative Algebra*. Birkhäuser, Boston, MA (1996).

[49] R. Stanley. Graph colorings And Related Symmetric functions: Ideas and Applications. *Discrete Math.* **193**, 267–286 (1998).

[50] R. Stanley. Positivity problems and conjectures. In *Mathematics: Frontiers and Perspectives*. International Mathematics Union (to appear).

[51] R. Stanley and J. R. Stembridge. On immanants of Jacobi-Trudi matrices and permutations with restricted positions. *J. Combin. Theory Ser. A* **62**, 261–279 (1993).

[52] J. Stembridge. Immanants of totally positive matrices are nonnegative. *Bull. London Math. Soc.* **23**, 422–428 (1991).

[53] J. Stembridge (1999). Personal communication.

[54] T. S. Sundquist, D. G. Wagner, and J.West. A Robinson-Schensted algorithm for a class of partial orders. *J. Combin. Theory Ser. A* **79**, 36–52 (1997).

[55] W. T. Trotter. *Combinatorics and Partially Ordered Sets: Dimension Theory*. Johns Hopkins University Press (1992).

[56] D. Wagner. Enumeration of functions from posets to chains. *European J. Combin.* **13**, 313–324 (1992).

[57] R. L. Wine and J. E. Freund. On the enumeration of decision patterns involving n means. *Ann. Math. Stat.* **28**, 256–259 (1957).

Index

adjacency matrix, 67
advantage, 68
$\alpha(S)$, 15
altitude, 73
    respecting labelling, 73
    sequence, 75
antiadjacency matrix, 64
    of a distributive lattice, 76
    relation to chain polynomial, 75
    squared, 67
antichain, 15

balanced complex, 20, 22
$\beta(S)$, 15
biletter, 27
biword, 27
Boolean algebra, 26
Brenti, 18
bubble sort, 55

$C(w)$, 34
Carlitz, 43, 44, 48
Catalan numbers, 64
chain, 12, 16, 63
chain map, 38
chain polynomial, 12, 23
    of a distributive lattice, 18
    of a $(\mathbf{3}+\mathbf{1})$-free poset, 64
    real zeros of, 13, 23
chromatic polynomial, 64
Clarke, 44
code, 25, 47
Cohen-Macaulay complex, 20, 22, 42

D, *see* descent set
DEN, 45–46
Denert, 45
Denert table, 45, 47
Denert's statistic, 45–46
dentab, *see* Denert table
des, 14, 23, 24
descent set, 14, 20, 23, 26, 44, 49, 54
descents, 14, 24, 45, 47
disjoint sum of chains, 33
distribution of a permutation statistic, 24
distributive lattice, 15, 20
    order complex of, 20
Distributive Lattice Conjecture, 17, 18
dmc, 25, 52
dmd, 47
dmi, 47
Dumont, 25, 26
Dumont's statistic, 24–42, 47, 52
    application, 33
    definition, 25
    definition for words, 28

E, *see* excedence set
E-polynomial, 19
e-positivity, 64
eigenvalues, 67
E_n, 46
equidistribution, 43
    of MAJ and INV, 43, 49
    of stm-set and descent set, 56
etab, 29
Eulerian numbers, 24
Eulerian polynomial, 25
Eulerian statistic, 23, 44, 47
    definition, 25
    examples, *see* des, dmc, dmd, dmi, exc, ides, stc, stm, z
Eulerian word statistic, 28
Eulerian-Mahonian statistic pairs, 26
exc, 24, 45

excedance, 24, 45
in a word, 27
value, 27
excedance set, 26, 46
excedance table, 28

f-vector, 11, 12
of a distributive lattice, 16, 18, 23, 76
of a poset, 12, 14, 22, 23
of a simplicial complex, 19
Ferrers poset, 18
Ferrers shape, 64
Fishburn, 63
flag complex, 22, 42
flag f-vector, 15, 19, 23
flag h-vector, 15, 19, 23
Foata, 43–46
forest, 41

Garsia, 43
Gasharov, 18
Gaussian poset, 18
generalized submatrix, 67, 72
Gessel, 43
graded poset, 42

h-polynomial, 13
real zeros of, 13
h-vector, 12
combinatorial interpretation of, 15
of a distributive lattice, 16, 22, 23, 76
of a poset, 12, 14, 23
of a simplicial complex, 19
Han, 45
Hecke monoid, 58

ides, 43
IMAJ, 43
incomparability graph, 64
increasing rearrangement, 21
induced subposet, 63
infinite Toeplitz matrix, 67
interval order, 63
INV, 23, 43, 46
inversion set, 62
inversion table, 47
inversions, 25, 43

$J(P)$, 15
join operation, 38
joint distribution, 43
of (des, ides, MAJ, IMAJ), 43
of (des, INV), 43
of (des, MAJ), 43–45, 54
of (des, MAJ, INV), 43
of (des, MAJ, INV, stc), 62
of (dmc, des), 61
of (dmc, MAJ), 43
of (dmc, X, MAJ, des), 61, 62
of (exc, DEN), 45
of (exc, DEN, Y, Z), 61
of (exc, Z), 61
of (INV, MAJ), 43
of (stc, INV), 54
symmetric, 43, 62
Jordan-Hölder set, 14, 23, 26, 43

Katona, 11
Kostant, 75
Kruskal, 11

$\mathcal{L}(P)$, 14, 20
labelled poset, 14
labelling, 64
altitude respecting, 73
LC, *see* letter set
letter set, 26
linear extensions, 14, 15, 20, 43
by descent, 12, 14, 43
$\mathcal{L}_k(P)$, 14, 17, 20
log-concavity, 11, 18

MacMahon, 44
Mahonian statistic, 23, 44, 45
examples, *see* DEN, IMAJ, INV, MAJ, r-MAJ, X, Y
MAJ, 23, 44, 47
major index, 25, 43, 45, 47
major index table, 47–57
corresponding to σ, 50

majtable, 47
missing face, 22
modular lattice, 18
Motzkin paths, 42

natural labelling, 14, 17, 18, 64
Neggers, 11, 17–19, 23, 76
nondecreasing rearrangement, 27

order complex, 19, 23, 33
    of a distributive lattice, 20
order ideal, 15, 72
    dual, 72
    principal, 72

permutation statistic, 23, 24, 43
    classes, *see* Eulerian statistic, Mahonian statistic
    distribution of, 24
Poset Conjecture, 18
Postnikov, 75
product of chains, 33

q-analog of n-factorial, 44
q-Eulerian polynomial, 44

$R(w)$, 27
rank, 16, 18
rearrangement, 27
rearrangement class, 27
    of a biword, 28
    set of codes of, 34
r-MAJ, 62

s-positivity, 64
Schützenberger, 11
sequence of restricted permutations, 47
    corresponding to σ, 50
series-parallel poset, 18, 41, 75
σ-major index table, 50
σ-stm-set (σ-STM), 56
simplex map, 20
squared antiadjacency matrix, 67–72
    combinatorial interpretation of, 67
st, 52
st-set (ST), 55
stair word, 46
stairs beneath the code, 52
Stanley, 11, 17–19, 23, 64, 75, 76
stc, 26, 46, 52, 53
    histrogram interpretation of, 52
stc-set (STC), 26, 55, 56
    algebraic description of, 57
    algorithmic description of, 55
Steingrímsson, 44
Stembridge, 18
stm_σ, 53
strictly decreasing word, 46
sub-stair words, 46
symmetric distribution
    of $(\mathrm{des}, \mathrm{dmc})$, 62
    of $(\mathrm{des}, \mathrm{stc})$, 62
    of (MAJ, INV), 43, 62
    of (MAJ, X), 62
symmetric functions, 76

$(\mathbf{3}+\mathbf{1})$-free poset, 63
    characterization of, 63, 70
Toeplitz matrix, 72
totally positive matrix, 67, 72
tree representation, 42
$(\mathbf{2}+\mathbf{2})$-free poset, 63

unimodality, 11, 18
unit interval order, 63
    antiadjacency matrix of, 64
    characterization of, 64

vector inequalities, 65
vertex map, 20, 37

W-polynomial, 19
word statistic, 27
    Eulerian, 28
words, 27

X (Mahonian statistic), 58

Y (Mahonian statistic), 61

z (Eulerian statistic), 61
Zeilberger, 44–46